



Supplementary Material: Aqueous Extract of *Psiloxylon mauritianum*, Rich in Gallic Acid, Prevents Obesity and Associated Deleterious Effects in Zebrafish


Batoul Ghaddar ¹, Laura Gence ^{1,†}, Bryan Veeren ^{1,†}, Matthieu Bringart ¹, Jean-Loup Bascands ¹, Olivier Meilhac ^{1,2} and Nicolas Diotel ^{1,*}

¹ Université de La Réunion, INSERM, UMR 1188, Diabète Athérombose Thérapies Réunion Océan Indien (DéTROi), Saint-Denis, 97400 La Réunion, France; batoul.ghaddar@univ-reunion.fr (B.G.); laura.gence@univ-reunion.fr (L.G.); bryan.veeren@univ-reunion.fr (B.V.); matthieu.bringart@inserm.fr (M.B.); jean-loup.bascands@inserm.fr (J.-L.B.); olivier.meilhac@univ-reunion.fr (O.M.)

² CHU de La Réunion, Saint-Denis, 97400, La Réunion, France


* Correspondence: nicolas.diotel@univ-reunion.fr

† These authors contribute equally to this work.



Supplementary Figure S1: Body weight of CTRL and *P. mauritianum* treated fish

After three days of treatment with water (CTRL) or *P. mauritianum* (0.25 g/L), no significant change in the body weight of CTRL and *P. mauritianum* treated fish was observed.



Supplementary Figure S2: Bran cell proliferation in the main neurogenic niches

In the therapeutic protocol, neurogenesis was not different between control, DIO and DIO + *P. mauritianum* at week 2 of the experiment for CTRL, DIO, DIO+ *P. mauritianum* treated fish in the main neurogenic niches studied.