

## Supporting Information

# *Scutellaria petiolata* Hemsl. ex Lace & Prain (Lamiaceae).: A New Insight in Biomedical Therapies


Sidra Mubin<sup>1</sup>, Najeeb Ur Rehman<sup>2</sup>, Waheed Murad<sup>3</sup>, Muddaser Shah<sup>2,3\*</sup>, Ahmed Al-Harrasi<sup>2</sup> and Rabia Afza<sup>1\*</sup>

<sup>1</sup>Department of Botany, Hazara University Mansehra, Mansehra 21310, Pakistan; shahhu123@gmail.com (SM); ethnokp@gmail.com(RA).

<sup>2</sup>Natural and Medical Sciences Research Center, University of Nizwa, P.O. Box 33, Birkat Al Mauz, Nizwa 616, Oman; najeeb@unizwa.edu.om (N.U.R)

<sup>3</sup>Department of Botany, Abdul Wali Khan University Mardan, Mardan 23200, Pakistan; waheedmurad@awkum.edu.pk (W.M)


\*Correspondance: muddasershah@awkum.edu.pk (MS) aharrasi@unizwa.edu.om(A.A.H), rabiaafza@hu.edu.pk(RA),



A)

B)

**Figure S1.** Full-scan ESI-LC-MS mass spectrum of the SPM extract of *S. petiolata*. (A) negative ionization mode and (B) positive ionization mode.






Figure S2: Mass spectrum of the respective tentatively identified chemical ingredients via ESI-LC-MS analysis in the active MeOH extract of *S. petiolata*.