

Article

Hplc-Dad Phenolic Characterization and Antioxidant Activities of Ripe and Unripe Sweet Orange Peels

Olufunmilayo Sade Omoba ^{1,*}, Rebecca Olajumoke Obafaye ^{1,†}, Sule Ola Salawu ^{2,†},
Aline Augusti Boligon ^{3,†} and Margareth Linde Athayde ^{3,†}

¹ Department of Food Science and Technology, Federal University of Technology, P.M.B. 704, Akure 340284, Nigeria; E-Mail: ajufaye@gmail.com

² Department of Biochemistry, Federal University of Technology, P.M.B. 704, Akure 340284, Nigeria; E-Mail: sosalawu@futa.edu.ng

³ Phytochemical Research Laboratory, Department of Industrial Pharmacy, Federal University of Santa Maria, Build 26, Room 1115, Santa Maria, CEP 97105-900, Brazil; E-Mails: alineboligon@yahoo.com.br (A.A.B.); marga@ccs.ufsm.br (M.L.A.)

[†] These authors contributed equally to this work.

* Author to whom correspondence should be addressed; E-Mail: omobasade@gmail.com; Tel.: +234-706-483-0497.

Academic Editors: Antonio Segura-Carretero and David Arráez-Román

Received: 5 April 2015 / Accepted: 19 June 2015 / Published: 9 July 2015

Abstract: Phenolic compounds of unripe and ripe sweet orange peels were determined using a high-performance liquid chromatography separation method with diode array detector (HPLC-DAD). The *in vitro* antioxidant properties and the EC₅₀ (concentration required to obtain a 50% antioxidant effect) values were also determined. The predominant phenolic compounds were quercitrin, rutin, and quercetin with values of 18.77 ± 0.01 mg/mL, 18.65 ± 0.03 mg/mL, and 10.39 ± 0.01 mg/mL respectively in unripe orange peel and 22.61 ± 0.01 mg/mL, 17.93 ± 0.03 mg/mL, and 14.03 ± 0.02 mg/mL respectively in ripe orange peel. The antioxidant properties revealed 2,2'-azino-bis(3-ethyl benzothiazoline-6-sulfonic acid) diammonium salt (ABTS) scavenging ability of both unripe and ripe orange peels respectively as 14.68 ± 0.01 and 16.89 ± 0.02 mmol TEAC/g, the Ferric Reducing Antioxidant Properties (FRAP) as 70.69 ± 0.01 and 91.38 ± 0.01 mg gallic acid equivalents/100g, total phenol content as 5.27 ± 0.03 and 9.40 ± 0.01 mg gallic acid equivalents/g and total flavonoid content as 3.30 ± 0.30 and 4.20 ± 0.02 mg quercetin equivalent/g. The antioxidant assays showed enhanced potency of extract from ripe orange peel with EC₅₀ values of 2.71 ± 0.03 mg/mL for 2,2-diphenyl-1-picrylhydrazyl (DPPH),

0.67 ± 0.03 mg/mL for hydroxyl radicals (OH^*), 0.57 ± 0.02 mg/mL for Fe^{2+} chelation, and 0.63 ± 0.06 mg/mL for malondialdehyde (MDA), and was more potent than unripe orange peel.

Keywords: orange peel; phenolic profile; *in vitro* antioxidant

1. Introduction

In today's consumer perception of agriculture and food production, aspects like health, safety, and quality have become the key words [1]. The uncontrolled production of oxygen-derived free radicals is implicated in the onset of many diseases such as cancer, rheumatoid arthritis, and atherosclerosis, as well as in the degenerative processes associated with aging [2]. Antioxidants are abundant compounds primarily found in fresh fruits and vegetables, and their roles in the prevention of degenerative diseases are emerging. Two basic categories of antioxidants exist, namely, synthetic and natural antioxidants. Synthetic antioxidants are compounds with phenolic structures of various degrees of alkyl substitution, while natural antioxidants can be phenolic compounds (tocopherols, flavonoids, and phenolic acids), nitrogen compounds (alkaloids, chlorophyll derivatives, amino acids, and amines), or carotenoids as well as ascorbic acid [3]. However, the most commonly used synthetic antioxidants such as butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), have been alleged of being responsible for liver damage and carcinogenesis [4], thus, the growing interest in natural antioxidants.

Citrus sinensis, sweet orange is a fruit with a small evergreen tree (8–15 ft), or shrub in the *Rutaceae* (citrus family). It originated from southern China, where it has been cultivated for millennia. Orange is now grown commercially worldwide in tropical, semi-tropical, and some warm temperate regions, and has become the most widely planted fruit tree in the world [5]. Orange is the world's most popular fruit, and is eaten fresh and used for juice [6]; the fruit is made up of peel, pulp, seeds, and juice. In contrast with other types of fruits, around 34% of the fruit is used for juice production, yielding about 44% of peels as byproducts [7]. The peel and seeds results in a considerable amount of by-products which might be a source of environmental pollution since they are prone to microbial spoilage. Citrus by-products could be of use as functional ingredients in the production of functional foods, since they are good sources of dietary fiber and bioactive compounds [8]. Peel is a good source of phenolic compounds which may potentially be used in food formulations or when extracted can be used as natural antioxidants to prevent oxidation of selected foods [9]. The citrus peel and seeds are very rich in phenolic compounds, such as phenolic acids and flavonoids; the peel is richer in flavonoids than seeds [10]. Flavonoids are a group of natural compounds with variable phenolic structures and are found in plants, they are oxidized by radicals, resulting in a more stable, less-reactive radical. They can be divided into a variety of classes such as flavones (e.g., flavone, apigenin, and luteolin), flavonols (e.g., quercetin, kaempferol, myricetin, and fisetin), flavanones (e.g., flavanone, hesperetin, and naringenin), and others. Several flavonoids such as catechin, apigenin, quercetin, naringenin, rutin, and venoruton are reported for their hepatoprotective activities [11]. Selected flavonoids can directly scavenge superoxides, whereas other flavonoids can scavenge the highly reactive oxygen-derived radical called peroxynitrite. Epicatechin and rutin are also powerful radical scavengers [12]. The scavenging ability of rutin may be

due to its inhibitory activity on the enzyme xanthine oxidase. Rutin is also known for its anti-inflammatory and vasoactive properties, as well as for its capability to diminish capillary permeability and to reduce the risk of arteriosclerosis, whereby reducing coronary heart disease, possibly through the diminishing of platelet aggregation [13,14]. There are also studies that show a dose-response effect of rutin in inhibiting low-density lipoprotein (LDL) peroxidation [15] and the antioxidant activity of rutin in Fenton reaction [16]. Rutin was suggested as one of the most potent natural inhibitors of advanced glycation endproducts (AGEs) accumulation in human body, suggesting that its anti-glycation activity may mainly be due to its radical scavenging activity [17]. Specific flavonoids are known to chelate iron [18] thereby removing a causal factor for the development of free radicals. Quercetin in particular is known for its iron-chelating and iron-stabilizing properties.

Several authors describe the use of high-performance liquid chromatography with diode array detector (HPLC-DAD) for characterization and quantification of phenolic composition [18–22]. The presence of compounds with antioxidant properties in *Citrus sinensis* has been described using HPLC-DAD and UV–Vis spectra [22]. UV detectors are popular among all the detectors because they offer high sensitivity and also because majority of naturally occurring compounds encountered have some UV absorbance, phenolics are frequently identified using UV-VIS detector at wavelengths 190–380 nm [17]. Furthermore, the high sensitivity of UV detection is very important for the compounds of interest that are present in small amounts in a sample [20].

Despite all of the possible uses listed above, citrus peel remains underutilized, and it is treated as waste or as feed to ruminant producing meat in Nigeria. Some earlier reports on the phenol constituents and antioxidant properties of citrus peel are mainly determined on the soluble free phenolic [9,22] which is on the basis of the solvent soluble extraction. Therefore, this study sought to identify and quantify the phenolic compounds in both unripe and ripe orange peels using HPLC-DAD and investigate the *in vitro* antioxidant properties of both peels.

2. Experimental Section

2.1. Chemicals and Reagents

All chemicals used were of analytical grade for both HPLC analysis and spectrophotometric analysis. Deoxy-D-ribose, 2,2'-Azino-bis(3-ethylbenzothiazoline-6-sulfonic acid)diammonium salt, 2,2-diphenyl-1-picrylhydrazyl were obtained from Sigma Chemical Company (St. Louis, MO, USA). Methanol, formic acid, caffeic acid, catechin and epicatechin purchased from Merck (Darmstadt, Germany). Naringin (7-[[2-O-(6-deoxy-L-mannopyranosyl)-D-glucopyranosyl]oxy]-2,3-dihydro-5-hydroxy-2-(4-hydroxyphenyl)-4H-1-benzopyran-4-one), quercetin (3,3',4,5,7-pentahydroxyflavone), quercitrin (2-(3,4-dihydroxyphenyl)-5,7-dihydro-3-(3,4,5-trihydroxy-6-methyloxan-2-yl)oxychromen-4-one), kaempferol (3,4,5,7-tetrahydroxyflavone), luteolin (3,4,5,7-tetrahydroxyflavone) and rutin (2-(3,4-dihydroxyphenyl)-5,7-dihydroxy-4-oxo-4H-chromen-3-yl-6-O-(6-deoxy- α -L-mannopyranosyl)- β -D-glucopyranoside) were acquired from Sigma Chemical Company. All other chemicals used were of analytical grade. Distilled water was used for the *in vitro* antioxidant analysis.

2.2. Orange Peel Powder Production

Matured sweet oranges (Navel variety) were harvested from a tree in a commercial farm, aged 10 years, the fertilizers used were 2 kg of NPK (Nitrogen, Phosphorus and Potassium, at ratio of 15:15:15) and 600 g K₂O (Potassium Oxide) per tree, twice in the year, June and September. Irrigation was done during the long dry season of November to February, ten liters of water was delivered at the base of the trees twice a week. The commercial farm, was located at Aba Oyo Village (Federal University of Technology, Akure, Nigeria), with topographical elevation between 260 and 470 m above sea level. The climatic condition of the area during the study was peak dry condition between early December and late February 2013. The unripe orange fruits (physiologically matured with green peel) with 9° Brix and ripe orange fruits (commercially matured with orange yellow peel) with 13° Brix, were picked early in the morning by 9 am. Sweet oranges with an average weight of 175.0 ± 2 g each were used; they were washed with distilled water, and then manually peeled. The obtained peels were air dried in a ventilated oven at 40 °C for 48 h, milled using a laboratory hammer mill (Mikro Samplmill Lab Hammer Mill, ¾ HP motor) and passed through a 24 mesh sieve size to produce the orange peel powder (OPP) [23]. OPP produced was packed in polythene and kept in a sealed container at 4 °C until a month later when it was analyzed.

2.3. Decoction

Dried OPP (5 g) was boiled in water (100 mL) for 10 min. After cooling, the sample was centrifuged (5000 rpm for 10 min) and the clear solution was recovered in a conical flask, and then rinsed to 100 mL with water. This sample was analyzed as such by HPLC-DAD.

2.4. HPLC-DAD Analysis

High performance liquid chromatography (HPLC-DAD) was performed with a Shimadzu Prominence Auto Sampler (SIL-20A) HPLC system (Shimadzu, Kyoto, Japan), equipped with Shimadzu LC-20AT reciprocating pumps connected to a DGU 20A5 degasser with a CBM 20A integrator, SPD-M20A diode array detector and LC solution 1.22 SP1 software.

Quantification of Compounds

Reverse phase chromatographic analyses were carried out under gradient conditions using C₁₈ column (4.6 mm × 150 mm, 5 µm diameter particles, Dikma Technologies Inc., Beijing, China); the mobile phase was water containing 1% formic acid (A) and methanol (B), and the composition gradient was: 12% of B until 10 min and changed to obtain 20%, 30%, 50%, 60%, 70%, 20% and 10% B at 20, 30, 40, 50, 60, 70 and 80 min, respectively, following the method described by [24] with slight modifications. Orange peel (ripe and unripe) extracts and mobile phase were filtered through 0.45 µm membrane filter (Millipore) and then degassed by ultrasonic bath (25 °C; 10 min; 120Watts). Prior to use, the OPP extract was analyzed; at a concentration of 15 mg/mL. The flow rate was 0.6 mL/min, injection volume 50 µL and the wavelength were 280 nm for catechin and epicatechin, 327 nm for caffeic acid, and 365 nm for quercetin, quercitrin, kaempferol, luteolin, naringin and rutin. All the samples and mobile phase were filtered through 0.45 µm membrane filter (Millipore) and then degassed by ultrasonic

bath prior to use. Stock solutions of standards references were prepared in the HPLC mobile phase at a concentration range of 0.025–0.250 mg/mL for quercetin, quercitrin, kaempferol, luteolin, naringin and rutin; and 0.030–0.350 mg/mL for caffeic acid, epicatechin and catechin. Chromatography peaks were confirmed by comparing its retention time with those of reference standards and by DAD spectra (200 to 500 nm). Calibration curve for catechin: $Y = 13067 X + 1281.9$ ($r = 0.9997$); epicatechin: $Y = 11983 X + 1162.5$ ($r = 0.9999$); caffeic acid: $Y = 11961 X + 1187.0$ ($r = 0.9998$); naringin: $Y = 12647 X + 1346.9$ ($r = 0.9994$); quercitrin: $Y = 11893 X + 1187.6$ ($r = 0.9997$); rutin: $Y = 12706 X + 1384.5$ ($r = 0.9999$), kaempferol: $Y = 13509 X + 1305.8$ ($r = 0.9991$), luteolin: $Y = 11865 X + 1259.6$ ($r = 0.9998$) and quercetin: $Y = 13507 X + 1271.9$ ($r = 0.9997$). All chromatography operations were carried out at ambient temperature and in triplicate.

The limit of detection (LOD) and limit of quantification (LOQ) were calculated based on the standard deviation of the responses and the slope using three independent analytical curves. LOD and LOQ were calculated as 3.3 and $10 \sigma/S$, respectively, where σ is the standard deviation of the response and S is the slope of the calibration curve [25].

2.5. In vitro Antioxidant Studies of Unripe and Ripe Orange Peel

2.5.1. Determination of Total Phenol Content

The total phenol content was determined as suggested by Pereira *et al.* [26]. Briefly, appropriate dilutions of the extracts were oxidized with 2.5 mL 10% Folin-Ciocalteu's reagent (v/v) and neutralized by 2.0 mL of 7.5% sodium carbonate. The reaction mixture was incubated for 40 min at 45 °C and the absorbance was measured at 765 nm using a Visible Spectrophotometer (Model 721 Visible Spectrophotometer, Axiom Mediral LMD, UK). The total phenol content was subsequently calculated as gallic acid equivalent.

2.5.2. Determination of Total Flavonoid Content

The total flavonoid content of both extracts was determined using Folin Ciocalteu reagent as described by Singleton *et al.* [27]. About 0.5 mL of appropriately diluted sample was mixed with 0.5 mL methanol, 50 μ L 10% $AlCl_3$, 50 μ L 1M potassium acetate and 1.4 mL water, and incubated at room temperature for 30 min. The absorbance of the reaction mixture was subsequently measured at 415 nm using a Visible Spectrophotometer (Model 721 Visible Spectrophotometer); the total flavonoid content was then calculated. The non-flavonoid polyphenols were taken as the difference between the total phenol and total flavonoid content.

2.5.3. Determination of Ferric Reducing Antioxidant Property

The ferric reducing property of the extracts was determined by assessing the ability of the extract to reduce $FeCl_3$ solution as described by Oyaizu [28]. About 2.5 mL aliquot was mixed with 2.5 mL 200 mM sodium phosphate buffer (pH 6.6) and 2.5 mL 1% potassium ferricyanide. The mixture was incubated at 50 °C for 20 min. and then 2.5 mL 10% trichloroacetic acid was added. This mixture was centrifuged at 650 rpm for 10 min and 5 mL of the supernatant was mixed with an equal volume of water and 1 mL of 0.1% ferric chloride. The absorbance was measured at 700 nm using a Visible

Spectrophotometer (Model 721 Visible Spectrophotometer), and the ferric reducing antioxidant property was subsequently calculated.

2.5.4. ABTS* Scavenging Ability

The ABTS* scavenging ability of both extracts were determined according to the method described by Re *et al.* [29]. The ABTS* was generated by reacting an (7 mmol/L) ABTS aqueous solution with $K_2S_2O_8$ (2.45 mmol/L, final concentration) in the dark for 16 h and adjusting the Abs at 734 nm to 0.700 with ethanol. A measurement of 0.2 mL of both solvent extracts at different concentrations was added to 2.0 mL ABTS* solution and reaction mixture was allowed to stand at 30 °C the absorbances were measured using a Visible Spectrophotometer (Model 721 Visible Spectrophotometer) at 734 nm after 15 min. The trolox equivalent antioxidant capacity was subsequently calculated.

2.5.5. DPPH Free Radical Scavenging Ability

The free radical scavenging ability of the extracts against DPPH (1,1-diphenyl-2 picrylhydrazyl) free radical was evaluated as described by Gyamfi *et al.* [30]. Appropriate dilution of the extracts (1 mL) was mixed with 1 mL; 0.4 mM methanolic solution containing DPPH radicals, the mixture was left in the dark for 30 min and the absorbance was measured using a Visible Spectrophotometer (Model 721 Visible Spectrophotometer) at 516 nm. The DPPH free radical scavenging ability was subsequently calculated.

2.5.6. Fenton Reaction (Degradation of Deoxyribose)

The ability of the extract to prevent Fe^{2+}/H_2O_2 induced decomposition of deoxyribose was determined as described by Halliwell *et al.* [31]. The extract 0–100 μ L was added to a reaction mixture containing 120 μ L of 20 mM deoxyribose, 400 μ L of 0.1 mM phosphate buffer, 40 μ L of 500 mM of $FeSO_4$, and the volumes were made up to 800 μ L with distilled water. The reaction mixture was incubated at 37 °C for 30 min and the reaction was then stopped by the addition of 0.5 mL of 28% trichloro acetic acid. This was followed by addition of 0.4 mL of 0.6% thiobarbituric acid solution. The tubes were subsequently incubated in boiling water for 20 min. The absorbance was measured at 532nm using a Visible Spectrophotometer (Model 721 Visible Spectrophotometer).

2.5.7. Fe^{2+} Chelation Assay

The Fe^{2+} chelating ability of both extracts were determined using a modified method of Minotti and Aust, [32]. Freshly prepared 500 μ M $FeSO_4$ (150 μ L) was added to a reaction mixture containing 168 μ L 0.1 M Tris-HCl (pH 7.4), 218 μ L saline and the extracts (0–25 μ L). The reaction mixture was incubated for 5 min, before the addition of 13 μ L 0.25% 1,10-phenanthroline (w/v). The absorbance was subsequently measured at 510 nm using a Visible Spectrophotometer (Model 721 Visible Spectrophotometer). The $Fe(II)$ chelating ability was subsequently calculated.

2.5.8. Lipid Peroxidation Assay

The rats were decapitated under mild diethyl ether anaesthesia and the pancreas was rapidly isolated and placed on ice and weighed. This tissue was subsequently homogenized in cold saline (1/10, w/v) with about 10-up-and-down strokes at approximately 1200 r/min in a Teflon glass homogenizer (Glas-Col Co., CA, USA). The homogenate was centrifuged for 10 min at 3000× g to yield a pellet that was discarded, and a low-speed supernatant (S1) was kept for lipid peroxidation assay [33].

The lipid peroxidation assay was carried out using the modified method of Ohkawa *et al.* [34], 100 µL S1 fraction was mixed with a reaction mixture containing 30 µL of 0.1 M pH 7.4 Tris-HCl buffer, extract (0–100 µL) and 30 µL of 250 µM freshly prepared FeSO₄. The volume was made up to 300 µL by water before incubation at 37 °C for 1 h. The color reaction was developed by adding 300 µL of 8.1% SDS (sodium dodecyl sulphate) to the reaction mixture containing S1; this was subsequently followed by the addition of 600 µL of acetic acid/hydrochloric acid at ratio 8:3 and then adjusted to pH 3.4. Added to the mixture was 600 µL of 0.8% TBA (thiobarbituric acid), this mixture was incubated at 100 °C for 1 h. TBARS (thiobarbituric acid reactive species) produced were measured at 532 nm and the absorbance was compared with that of standard curve using MDA (malondialdehyde).

2.6. Statistical Analysis

All samples for the phenolic profile were analyzed in triplicate and the results are expressed as mean ± standard deviations (SD) of three determinations. Averages followed by different letters in the column differ by Tukey test at $p < 0.05$. Values with different alphabet along the same row are significantly different ($p < 0.05$) by one-way analysis of variance (ANOVA).

Data for the *in vitro* antioxidant activities were analyzed and expressed as mean ± standard error mean (S.E.M.). The statistical analysis was carried out using one-way analysis of variance (ANOVA) and means were separated using Duncan multiple range test (DMRT), $p < 0.05$ were considered as significant. The EC₅₀ was determined using linear regression analysis for the antioxidant assay.

3. Results and Discussion

3.1. Identification and Quantification of Phenolics Using HPLC-DAD

HPLC-DAD phenolic compounds analysis of orange peel (unripe and ripe) extracts revealed the presence of catechin ($t_R = 14.37$ min; peak 1), caffeic acid ($t_R = 24.79$ min; peak 2), naringin ($t_R = 31.25$ min; peak 3), epicatechin ($t_R = 34.97$; peak 4), rutin ($t_R = 43.81$ min; peak 5), quercitrin ($t_R = 49.68$ min; peak 6), quercetin ($t_R = 52.41$ min; peak 7), kaempferol ($t_R = 56.11$ min; peak 8) and luteolin ($t_R = 58.39$ min; peak 9) (Table 1 and Figure 1). A total of nine phenolic compounds which included four flavonoids, were identified and quantified as shown in Table 1. The chromatographic parameters of the method used for quantification of the compounds are shown in Table 2. Among the phenolic compounds, quercitrin and rutin showed highest concentration in both unripe and ripe peel with 18.79 mg/g and 18.6 mg/g obtained for unripe orange peel while 22.61 mg/g and 17.93 mg/g were obtained for ripe orange peel. Rutin, also called rutoside, quercetin-3-*O*-rutoside, and sophorin, was the glycoside between the flavonol quercetin and the disaccharide rutinose α -L-rhamnopyranosyl-

(1→6)-β-D-glucopyranose. Rutin, a bioflavonoid glycoside with strong antioxidant properties was found in many plants fruits and fruit rinds (especially the citrus fruits; orange, grapefruit, lemon, and lime), and apple and berries. Rutin (quercetin rutinoside), like quercitrin, is a glycoside of the flavonoid quercetin. As such, the chemical structures of both are very similar, with the difference existing in the hydroxyl functional group. Both quercetin and rutin are used in many countries as medications for blood vessel protection, and are ingredients of numerous multivitamin preparations and herbal remedies. Quercitrin has been reported to have antioxidant and anti-carcinogenic qualities via its inhibition of neoplastic transformation by blocking activation of the Mitogen-Activated Protein Kinase (MAPK) pathway and stimulation of cellular protection signaling [35]. It was observed from the present investigation that quercitrin and rutin concentration in ripe orange peel powder was higher than that of unripe orange peel powder. Reports on olive fruits showed that the phenolic contents have direct effect on the chemical composition of olive products which was dependent on the cultivar, the pedoclimatic production conditions, the agronomic techniques, and fruit ripening [36–39]. This result was in agreement with the report of [2], who observed that ripe raspberries (100% maturity) had stronger antioxidant activities and higher total anthocyanin content when compared with the pink stage (50% maturity). On the contrary, the report of [40] in tomato phenolics revealed lower quercitrin and rutin in ripe tomato. However, the observation suggested that OPP powder especially the ripe one might be used in the promotion of cardiovascular health.

Table 1. Phenolic acids and flavonoids composition of orange peel powder (ripe and unripe) extracts.

Compounds	Unripe Orange Peel (mg/g)	Ripe Orange Peel (mg/g)
Catechin	5.91 ± 0.03 ^{c,B}	12.49 ± 0.01 ^{c,A}
Caffeic acid	3.45 ± 0.01 ^{d,A}	3.57 ± 0.02 ^{c,A}
Naringin	3.58 ± 0.02 ^{d,B}	5.71 ± 0.02 ^{d,A}
Epicatechin	3.21 ± 0.01 ^{d,B}	6.08 ± 0.01 ^{d,A}
Rutin	18.65 ± 0.03 ^{a,A}	17.93 ± 0.03 ^{b,B}
Quercitrin	18.77 ± 0.01 ^{a,B}	22.61 ± 0.01 ^{a,A}
Quercetin	10.39 ± 0.01 ^{b,B}	14.03 ± 0.02 ^{b,A}
Kaempferol	5.86 ± 0.02 ^{c,A}	3.76 ± 0.03 ^{c,B}
Luteolin	3.19 ± 0.03 ^{d,B}	5.83 ± 0.01 ^{d,A}

Results are expressed as mean ± standard deviations (SD) of three determinations. Values with different alphabet along the same column (lowercase letters) are significantly different by Tukey test ($p < 0.05$). Values with different alphabet along the same row (capital letters) are significantly different ($p < 0.05$) by one-way analysis of variance (ANOVA).




Figure 1. Representative high performance liquid chromatography profile of unripe orange peel powder (A) and ripe orange peel powder (B). Catechin (peak 1), caffeic acid (peak 2), naringin (peak 3), epicatechin (peak 4), rutin (peak 5), quercitrin (peak 6), quercetin (peak 7), kaempferol (peak 8) and luteolin (peak 9).

Table 2. Chromatographic parameters of the method used for quantification of compounds in orange peel powder (ripe and unripe) extracts.

Compounds	Concentration Range mg/mL	Standard Curves	Correlation Coefficient (r)	LOD µg/mL	LOQ µg/mL
Catechin	0.030–0.350	$Y = 13067 X + 1281.9$	0.9997	0.017	0.056
Caffeic acid	0.030–0.350	$Y = 11961 X + 1187.0$	0.9998	0.026	0.090
Naringin	0.025–0.250	$Y = 12647 X + 1346.9$	0.9994	0.011	0.037
Epicatechin	0.030–0.350	$Y = 11983 X + 1162.5$	0.9999	0.032	0.105
Rutin	0.025–0.250	$Y = 12706 X + 1384.5$	0.9999	0.028	0.096
Quercitrin	0.025–0.250	$Y = 11893 X + 1187.6$	0.9997	0.009	0.034
Quercetin	0.025–0.250	$Y = 13507 X + 1271.9$	0.9997	0.019	0.063
Kaempferol	0.025–0.250	$Y = 13509 X + 1305.8$	0.9991	0.021	0.069
Luteolin	0.025–0.250	$Y = 11865 X + 1259.6$	0.9998	0.035	0.118

LOD, limit of detection; LOQ, limit of quantification.

3.2. In vitro Antioxidant Studies of Unripe and Ripe Orange Peel

Table 3 reveals the antioxidant properties (total phenols, total flavonoids, ABTS*, and FRAP) in OPP. The total phenolic and total flavonoid contents of ripe orange peel (9.40 mg GAE/g and 4.20 mg QE/g) were significantly higher than values obtained for the unripe orange peel of 5.27 mg GAE/g and 3.20 mg QE/g respectively. Phenolic compounds are important fruit constituents because they exhibit antioxidant activity by inactivating lipid free radicals or preventing decomposition of hydroperoxides into free radicals [41]. The phenolic contents obtained in the peels are higher than values obtained for orange juice [42] and some commonly consumed leafy vegetables in Nigeria [43,44]. Polyphenols such as flavanol contributed significantly to the ferric reducing capacity and radical scavenging capacity of orange peel during ripening, as ripe orange peel exhibited higher ABTS* (16.89 mmol/TEAC g) and FRAP (91.38 mg/GAE100 g) compared to the unripe orange peel. Peel extract gave a remarkable effect in protecting cells from oxidative damage [45], this could be due to the additional phenolic compounds present in peel compared to juice and pulp residue [46]. It implies therefore that the antioxidant capacities

of orange peel increased significantly during ripening as the ripe peel exhibited higher antioxidant activities and contents than the unripe peel.

Table 3. Antioxidant properties of unripe and ripe orange peel powder.

ANTIOXIDANT PROPERTIES	UNRIPE	RIPE
Total phenol content (mgGAE/g)	5.27 ± 0.03 ^b	9.40 ± 0.01 ^a
Total flavonoid content (mgQE/g)	3.30 ± 0.30 ^b	4.20 ± 0.02 ^a
ABTS* scavenging ability (mmol/TEAC g)	14.68 ± 0.01 ^b	16.89 ± 0.02 ^a
FRAP (mg/GAE100 g)	70.69 ± 0.01 ^b	91.38 ± 0.01 ^a

Values represent means ± standard deviation of triplicate ($n = 3$). Values with different alphabet along the same row (for the same antioxidant properties) are significantly different ($p > 0.05$).

Table 4 shows the EC_{50} of some antioxidant properties. EC_{50} (implies concentration required to obtain a 50% antioxidant effect) it is a typically used parameter to express the antioxidant capacity and to compare the activity of different compounds. A lower value of EC_{50} corresponds to a higher antioxidant activity of the orange peel extract. Free radical scavenging expressed as EC_{50} were 2.71 (unripe) and 2.23 mg/mL DPPH for ripe, the most effective being ripe orange peel. The trend in the results agree with the phenolic distribution in the citrus peels (Table 1) and other earlier reports, where correlations were reported between phenolic content and antioxidant capacity of some plant foods [42,45,46]. DPPH free radical scavenging ability is frequently used in the determination of free radical scavenging ability; however it has the limitation of color interference and sample solubility [47]. Ripe orange peel showed the highest scavenging activity against hydroxyl radicals (OH^*), Fe^{2+} chelation, and malondialdehyde with EC_{50} values of 0.57, 0.49 and 0.50 mg/mL, respectively when compared with the unripe orange peel. The Fe^{2+} chelating ability of the ripe orange peel ($EC_{50} = 0.49$ mg/mL) is higher than their corresponding OH^* scavenging abilities ($EC_{50} = 0.57$ mg/mL) as presented in Table 4; these findings agree with some earlier reports on the OH^* and Fe^{2+} chelating abilities of phenolics of some plant foods such as leafy vegetables, peppers, and spices that have stronger Fe^{2+} chelating than OH^* scavenging abilities [48,49]. The Fe^{2+} chelating ability of the phenolics could be attributed to the presence of two or more of the following functional groups namely, OH, -SH, -COOH, PO_3H_2 , C=O, -NR₂, -S- and -O- in a favourable structure-function configuration [50–52]. MDA formation is used as the general indicator of the extent of lipid peroxidation resulting from oxidative stress. It increases when Fe^{2+} induces lipid peroxidation by catalyzing the decomposition of hydrogen peroxide to generate hydroxyl radical via the Fenton reaction [53]. The ripe peel significantly ($p < 0.05$) inhibited MDA production in the pancreas in a dose-dependent manner, ripe peel had the highest inhibitory effect on MDA production in the pancreas (*in vitro*), while the unripe had the least inhibitory effect (Table 4). The possible mechanism through which the ripe peel of orange protects the pancreas could be by Fe^{2+} chelation [54] and the scavenging of OH^* [49,54,55] abilities. Since the phenolic extracts had higher Fe^{2+} chelating ability than OH^* scavenging ability, Fe^{2+} chelation could be the domineering mechanism through which the phenolics protect pancreas membrane from Fe^{2+} induced lipid peroxidation in the pancreas.

Table 4. EC₅₀ (mg/mL) of ripe and unripe orange peel on different antioxidant parameters.

ANTIOXIDANT PARAMETERS	UNRIPE	RIPE
DPPH*	2.71 ± 0.03 ^a	2.23 ± 0.03 ^b
OH*	0.67 ± 0.02 ^a	0.57 ± 0.03 ^b
Fe ²⁺ Chelation	0.57 ± 0.02 ^a	0.49 ± 0.01 ^b
Malondialdehyde (MDA)	0.63 ± 0.06 ^a	0.50 ± 0.01 ^b

Values represent means ± standard deviation of triplicate ($n = 3$). Values with the different alphabet along the same row (for the same parameter) are significantly different ($p > 0.05$).

4. Conclusions

According to HPLC-DAD analysis for the identification and quantification of phenolics, quercitrin and rutin were the most predominant phenolics. The *in vitro* antioxidant activities of the extracts of sweet orange peel revealed the radical scavenging activity of the unripe and ripe orange peel, which is strongly associated with the presence of flavonoids, phenolic acids, and their derivatives, thereby supporting the relevance of sweet orange peel as important dietary source of antioxidant compounds. Orange peel powder, especially the ripe orange peel may therefore be exploited as natural antioxidant in food application as well as for health supplements or functional food, to alleviate oxidative stress.

Author Contributions

Olufunmilayo Sade Omoba, Rebecca Olajumoke Obafaye and Sule Ola Salawu conceived and designed the experiments. Rebecca Olajumoke Obafaye and Sule Ola Salawu performed the experiments. Olufunmilayo Sade Omoba and Rebecca Olajumoke Obafaye wrote the paper. Aline Augusti Boligon and Margareth Linde Athayde participated in and facilitated the HPLC-DAD analysis. Rebecca Olajumoke Obafaye, Aline Augusti Boligon and Margareth Linde Athayde analyzed the data. All of the authors have read and approved the final manuscript.

Conflicts of Interest

The authors declare no conflict of interest.

References

1. Iqbal, S.; Haleem, S.; Akhtar, M.; Zia-ul-Haq, M.; Akbar, J. Efficiency of pomegranate peel extracts in stabilization of sunflower oil under accelerated conditions. *Food Res. Int.* **2008**, *41*, 194–200.
2. Wang, Z.; Luo, D. Antioxidant activities of different fractions of polysaccharide purified from *Gynostemma pentaphyllum* Makino. *Carbohydr. Polym.* **2007**, *68*, 54–58.
3. Hall, C.A.; Cuppet, S.L. Structure activities of natural antioxidants. In *Antioxidant Methodology in vivo and in vitro Concepts*; Aruoma, O.I., Cuppet, S.L., Eds.; Amer Oil Chemists Society: Champaign city, IL, USA, 1997; pp. 2–29.
4. Qi, H.M.; Zhang, Q.B.; Zhao, T.T.; Chenc, R.; Zhang, H.; Niu, X.Z. Antioxidant activity of different sulfate content derivatives of polysaccharide extracted from *Ulva pertusa* (Chlorophyta) *in vitro*. *Int. J. Biol. Macromol.* **2005**, *37*, 195–199.

5. Ehler, S.A. Citrus and its benefits. *J. Bot.* **2011**, *5*, 201–207.
6. Topuz, A.; Topakci, M.; Canakci, M.; Akinci, I.; Ozdemir, F. Physical and nutritional properties of four orange varieties. *J. Food Eng.* **2005**, *66*, 519–523.
7. Li, S.; Lo, C.; Ho, C. Hydroxylated Polymethoxyflavones and Methylated Flavonoids in Sweet Orange (*Citrus sinensis*) Peel. *J. Agric. Food Chem.* **2006**, *54*, 4176–4185.
8. Marin, F.R.; Martinez, M.T.; Uribesalgo, S.C.; Frutos, M.J. Changes in nutraceutical composition of lemon juices according to different industrial extraction systems. *Food Chem.* **2002**, *78*, 319–324.
9. Albishi, T.; John, J.A.; Al-Khalifa, A.S.; Shahidi, F. Phenolic content and antioxidant activities of selected potato varieties and their processing by-products. *J. Funct. Foods.* **2013**, *5*, 590–600.
10. Sawalha, S.M.S.; Arráez-Román, D.; Segura-Carretero, A. Quantification of main phenolic compounds in sweet and bitter orange peel using CE–MS/MS. *Food Chem.* **2009**, *116*, 567–574.
11. Tapas, A.R.; Sakarkar, D.M.; Kakde, R.B. Flavonoids as nutraceuticals: A review. *Trop. J. Pharm. Res.* **2008**, *7*, 1089–1099.
12. Hanasaki, Y.; Ogawa, S.; Fukui, S. The correlation between active oxygen scavenging and antioxidative effects of flavonoids. *Free Radic. Biol. Med.* **1994**, *16*, 845–850.
13. La Casa, C.; Villegas, I.; de la Lastra, C.; Motilva, V.; Calero, M.J. Evidence for protective and antioxidant properties of rutin, a natural flavone, against ethanol induced gastric lesions. *J. Ethnopharm.* **2000**, *71*, 45–53.
14. Jiang, P.; Burczynski, F.; Campbell, C.; Pierce, G.; Austria, J.A.; Briggs, C.J. Rutin and flavonoid contents in three buckwheat species *Fagopyrum esculentum*, *F. tataricum*, and *F. Homotropicum* and their protective effects against lipid peroxidation. *Food Res. Int.* **2007**, *40*, 356–364.
15. Cailet, S.; Yu, H.; Lessard, S.; Lamoureux, G.; Ajdukovic, D.; Lacroix, M. Fenton reaction applied for screening natural antioxidants. *Food Chem.* **2007**, *100*, 542–552.
16. Peng, X.; Zheng, Z.; Cheng, K.W.; Shan, F.; Ren, G.X.; Chen, F. Inhibitory effect of mung bean extract and its constituents vitexin and isovitexin on the formation of advanced glycation endproducts. *Food Chem.* **2008**, *106*, 475–481.
17. Ferrali, M.; Signorini, C.; Caciotti, B. Protection against oxidative damage of erythrocyte membrane by the flavonoid quercetin and its relation to iron chelating activity. *FEBS Lett.* **1997**, *416*, 123–129.
18. Boligon, A.A.; Athayde, M.L. Importance of HPLC in analysis of plants extracts. *Austin Chromatogr.* **2014**, *1*, 1–2.
19. Giuffrè, A.M. HPLC-DAD detection of changes in phenol content of red berry skins during grape ripening. *Eur. Food Res. Technol.* **2013**, *237*, 555–564.
20. Giuffrè, A.M. High performance liquid chromatography-diode array detector (HPLC-DAD) detection of *trans*-resveratrol: Evolution during ripening in grape berry skins. *Afr. J. Agric. Res.* **2013**, *8*, 224–229.
21. Albano, C.; Negro, C.; Tommasi, N.; Gerardi, C.; Mita, G.; Miceli, A.; de Bellis, L.; Blando F. Betalains, phenols and antioxidant capacity in Cactus Pear [*Opuntia ficus-indica* (L.) Mill.] fruits from Apulia (South Italy) Genotypes. *Antioxidants* **2015**, *4*, 269–280.
22. Di Stefano, R.; Cravero, M.C. The separation of hydroxycinnamates in wine. *Sci. Aliment.* **1992**, *12*, 139–144.

23. Chen, Z.T.; Chu, H.L.; Chyau, C.C.; Chu, C.C.; Duh, P.D. Protective effects of sweet orange (*Citrus sinensis*) peel and their bioactive compounds on oxidative stress. *Food Chem.* **2012**, *135*, 2119–2127.
24. Oboh, G.; Ademosun A.O. Comparative studies on the ability of crude polyphenols from some Nigerian citrus peels to prevent lipid peroxidation—*In vitro*. *Asian J. Biochem.* **2006**, *1*, 169–177.
25. Van-Acker, S.A.; van Den, D.J.; Tromp, M.N.; Bast, A. Structural aspects of antioxidant activity of flavanoids. *Free Radic. Biol. Med.* **2011**, *35*, 331–342.
26. Pereira, R.P.; Boligon, A.A.; Appel, A.S.; Fachineto, R.; Ceron, C.S.; Tanus-Santos, J.E.; Athayde, M.L.; Rocha, J.B.T. Chemical composition, antioxidant and anticholinesterase activity of *Melissa officinalis*. *Ind. Crop Prod.* **2014**, *53*, 34–45.
27. Singleton, V.L.; Orthofer, R.; Lamuela-Raventos, R.M. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Method Enzymol.* **1999**, *299*, 152–178.
28. Oyaizu, M. Studies on products of browning reaction: Antioxidative activity of products of browning reaction prepared from glucosamine. *Jpn. J. Nutr.* **1986**, *44*, 307–315.
29. Re, R.; Pellegrini, N.; Proteggente, A.; Pannala, A.; Yang, M.; Rice-Evans, C. Antioxidant activity applying an improved ABTS radical cation decolorisation assay. *Free Radic. Biol. Med.* **1999**, *26*, 1231–1237.
30. Gyamfi, M.A.; Yonamine, M.; Aniya, Y. Free-radical scavenging action of Medicinal herbs from Ghana: *Thonningia sanguinea* on experimentally-induced liver injuries. *Gen. Pharmacol.* **1999**, *32*, 661–667.
31. Halliwell, B.; Gutteridge, J.M.C.; Aruoma, O.I. The deoxyribose method: A sample test tube assay for determination of rate constant for reaction of hydroxyl radicals. *Anal. Biochem.* **1987**, *165*, 215–219.
32. Minotti, G.; Aust, S.D. An investigation into the mechanism of citrate-Fe²⁺-dependent lipid peroxidation. *Free Radic. Biol. Med.* **1987**, *3*, 379–387.
33. Belle, N.A.V.; Dalmolin, G.D.; Fonini, G.; Rubim, M.A.; Rocha, J.B.T. Polyamines reduces lipid peroxidation induced by different prooxidant agents. *Brain Res.* **2004**, *1008*, 245–251.
34. Ohkawa, H.; Ohishi, N.; Yagi, K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal. Biochem.* **1979**, *95*, 351–358.
35. Ding, M.; Zhao, J.S.; Bowman, L.; Lu, Y.J.; Shi, X.L. Inhibition of AP-1 and MAPK signaling and activation of Nrf2/ARE pathway by quercitrin. *Int. J. Oncol.* **2010**, *36*, 59–67.
36. Ben, O.N.; Roblain, D.; Chammen, N.; Thonart, P.; Hamdi, M. Antioxidant phenolic compounds loss during the fermentation of Chétoui olives. *Food Chem.* **2009**, *116*, 662–669.
37. Hanen, F.; Riadh, K.; Samia, O.; Sylvain, G.; Christian, M.; Chedly, A. Inter specific variability of antioxidant activities and phenolic composition in *Mesembryanthemum* genus. *Food Chem. Toxicol.* **2009**, *47*, 2308–2313.
38. Romero, M.P.; Tovar, M.J.; Girona, J.; Motilva, M.J. Changes in the HPLC phenolic profile of virgin olive oil from young trees (*Olea europaea* L. Cv. Arbequina) grown under different deficit irrigation strategies. *J. Agric. Food Chem.* **2002**, *50*, 5349–5354.

39. Vinha, A.F.; Ferreres, F.; Silva, B.M.; Valentão, P.; Gonçalves, A.; Pereira, J.A.; Andrade, P.B. Phenolic profiles of Portuguese olive fruits (*Olea europaea* L.): Influences of cultivar and geographical origin. *Food Chem.* **2005**, *89*, 561–568.
40. Raffo, A.; Leonardi, C.; Fogliano, V.; Ambrosino, P.; Salucci, M.; Gennaro, L.; Bugianesi, R.; Giuffrida, F.; Quaglia, G. Nutritional value of cherry tomatoes (*Lycopersicon esculentum* Cv. Naomi F1) harvested at different ripening stages. *J. Agric. Food Chem.* **2002**, *50*, 6550–6556.
41. Pokorny, J.; Yanishlieva, N.; Gordon, M. *Antioxidants in Food, Practical Applications*; Woodhead Publishing Ltd.: Cambridge, UK, 2001, pp. 145–171.
42. Sun, J.; Chu, Y.F.; Wu, X.; Liu, R.H. Antioxidant and anti-proliferation activities of common fruits. *J. Agric. Food Chem.* **2002**, *50*, 7449–7454.
43. Oboh, G.; Akindahunsi, A.A. Change in ascorbic acid, total phenol and antioxidant activity of sundried commonly consumed green leafy vegetables in Nigeria. *J. Nutr. Health* **2004**, *18*, 29–36.
44. Oboh, G. Effect of blanching on the antioxidant properties of some tropical green leafy vegetables. *LWT—Food Sci. Tech.* **2005**, *38*, 513–517.
45. Kim, H.; Moon, J.Y.; Kim, H.; Lee, D.S.; Cho, M.; Choi, H.K.; Kim, Y.S.; Mosaddik, A.; Cho, S.K. Antioxidant and anti-proliferative activities of mango (*Mangifera indica* L.) flesh and peel. *Food Chem.* **2010**, *121*, 429–436.
46. Li, B.B.; Smith, B.; Hossain, M. Extraction of phenolics from citrus peels. I. Solvent extraction Method. *Sep. Purif. Technol.* **2006**, *48*, 182–188.
47. Oboh, G.; Raddatz, H.; Henle, T. Antioxidant properties of polar and non-polar extracts of some tropical green leafy vegetables. *J. Sci. Food Agric.* **2008**, *88*, 2486–2492.
48. Oboh, G.; Rocha, J.B.T. Polyphenols in red pepper (*Capsicum annuum* var aviculare (Tepin) and their protective effect on some pro-oxidants induced lipid peroxidation in brain and liver. *Eur. Food Res. Technol.* **2007**, *225*, 239–247.
49. Oboh, G.; Rocha, J.B.T. Antioxidant and neuroprotective properties of sour tea (*Hibiscus abdarriffa*, calyx) and green tea (*Camellia sinensis*) on some pro-oxidants induced lipid peroxidation in brain *in vitro*. *Food Biophys.* **2008**, *3*, 382–389.
50. Lindsay, R.C. Food additives. In *Food Chemistry*, 2nd ed.; Fennema, O.R., Ed.; Marcel Dekker Inc.: New York, NY, USA, 1996; pp. 778–780.
51. Yuan, Y.V.; Bone, D.E.; Carrington, M.F. Antioxidant activity of pulse (*Palmaria palmata*) extract evaluated *in vitro*. *Food Chem.* **2005**, *91*, 485–494.
52. Gülçin, I. Antioxidant activity of caffeic acid (3,4-dihydroxycinnamic acid). *Toxicology* **2006**, *217*, 213–220.
53. Bayır, H.; Kochanek, P.M.; Kagan, V.E. Oxidative stress in immature brain after traumatic brain injury. *Int. J. Dev. Neurosci.* **2006**, *28*, 420–431.
54. Oboh, G.; Rocha, J.B.T. Antioxidant in foods: A new challenge for food processors. In *Leading Edge Antioxidants Research*; Panglossi, H.V., Ed.; Nova Science Publishers Inc.: New York, NY, USA, 2007; pp. 35–64.

55. Puntel, R.L.; Nogueira, C.W.; Rocha, J.B.T. Krebs cycle intermediates modulate thiobarbituricReactive species (TBARS) production in rat brain *in vitro*. *Neurochem. Res.* **2005**, *30*, 225–235.

© 2015 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>).