

**Table S1.** Antibodies and dilutions used to flow cytometry staining.

<b>Antibody</b>	<b>Company</b>	<b>Dilution</b>
Anti CD163 (mouse IgG1)	Bio-Rad, United Kingdom	1:100
Anti CD172a (mouse IgG1)	Bio-Rad, United Kingdom	1:100
Anti CADM1 (chicken IgY)	MBL International, USA	1:500
Anti CD14 (mouse IgG2b)	Bio-Rad, United Kingdom	1:200
Anti CD3 (mouse IgG1)	Southern Biotech, USA	1:100
Anti CD21 (mouse IgG1)	Southern Biotech, USA	1:200
Anti CD8 $\alpha$ (mouse IgG2a)	Bio-Rad, United Kingdom	1:200
Anti CD4 $\alpha$ (mouse IgG2b)	Bio-Rad, United Kingdom	1:100
Anti DEC205 (mouse x pig IgG1)	Homemade	1:100
Streptavidin PE-conjugated	Southern Biotech, USA	1:100
BV421-conjugated rat anti-mouse IgG1	(BioLegend, USA)	1:100
A647-conjugated goat anti-chicken IgY	Invitrogen, USA	1:500
FITC-conjugated goat anti-mouse IgG2b	SouthernBiotech, USA	1:100
A647-conjugated goat anti-mouse IgG2b	Invitrogen, USA	1:200
PE-conjugated mouse anti-pig IFN- $\gamma$	BD Pharmingen, USA	-