



Supplementary material

Interaction of Bortezomib with Cell Membranes Regulates Its Toxicity and Resistance to Therapy

Maria João Ramalho ^{1,2,*}, Stéphanie Andrade ^{1,2}, Joana Angélica Loureiro ^{1,2} and Maria Carmo Pereira ^{1,2,*}

¹ LEPABE—Laboratory for Process Engineering, Environment, Biotechnology and Energy, Faculty of Engineering, University of Porto, Rua Dr. Roberto Frias, 4200-465 Porto, Portugal

² ALiCE—Associate Laboratory in Chemical Engineering, Faculty of Engineering, University of Porto, Rua Dr. Roberto Frias, 4200-465 Porto, Portugal

* Correspondence: mjr Ramalho@fe.up.pt (M.J.R.); mcsp@fe.up.pt (M.C.P.); Tel.: +351-22-508-2262 (M.J.R.); +351-22-508-1590 (M.C.P.)

Table S1. Physicochemical properties of the prepared LUVs. Results are given as mean \pm SD (n=3).

LUVs	pH	Mean size (nm)	PdI	Zeta Potential (mV)
DMCP	7.4	122 \pm 4	0.09 \pm 0.04	-0.3 \pm 0.6
DMPC:Chol:SM	7.4	126 \pm 9	0.08 \pm 0.02	1.0 \pm 0.3
DMPC:Chol:SM	6.5	123 \pm 11	0.07 \pm 0.01	0.5 \pm 0.1