

Supplementary Table S1. Comparison of symptoms in supplemented and non-supplemented outpatients study groups at baseline and follow-up at days 7 and 14

Variable	Base-line			7 days			14 days		
	Supplemented outpatients n=22	Non-supplemented outpatients n=20	p-value	Supplemented outpatients n=22	Non-supplemented outpatients n=20	p-value	Supplemented outpatients n=22	Non-supplemented outpatients n=20	p-value
Fever ^a	9 (40.9)	8 (40.0)	0.95	0 (0.0)	2 (10.0)	0.22	0 (0.0)	2 (10.0)	0.22
Headache ^a	14 (64.3)	13 (65.0)	0.93	4 (18.2)	3 (15.0)	1.00	4 (18.2)	3 (15.0)	1.00
Loss of smell ^a	10 (45.5)	11 (55.0)	0.54	3 (13.6)	1 (5.0)	0.61	3 (13.6)	1 (5.0)	0.61
Dry cough ^a	13 (59.1)	15(75.0)	0.27	4 (18.2)	4(20.0)	1.00	4 (18.2)	4 (20.0)	1.00
Sore throat ^a	6 (27.3)	7 (35.0)	0.58	1 (4.5)	0 (0.0)	1.00	1 (4.5)	0 (0.0)	1.00
Ageusia ^a	10 (45.5)	12 (60.0)	0.34	3 (13.6)	4 (20.0)	0.69	3 (13.6)	4 (20.0)	0.69
Runny nose ^a	3 (13.6)	5 (25.0)	0.44	0 (0.0)	1 (5.0)	0.48	0 (0.0)	0 (0.0)	---
Nausea or Vomiting ^a	3 (13.6)	6 (30.0)	0.27	0 (0.0)	3 (15.0)	0.09	0 (0.0)	3 (15.0)	0.09
Tiredness ^a	8 (36.4)	12 (60.0)	0.13	4 (18.2)	4 (20.0)	1.00	5 (22.7)	4 (20.0)	1.00
Diarrhoea ^a	5 (22.7)	2 (25.0)	1.00	0 (0.0)	1 (5.0)	0.48	0 (0.0)	0 (0.0)	---
Myalgia ^a	12 (54.5)	20 (60.0)	0.72	1 (4.5)	2 (10.0)	0.60	1 (4.5)	2 (10.0)	0.60
Arthralgia ^a	8 (36.4)	12 (60.0)	0.13	0 (0.0)	0 (0.0)	---	0 (0.0)	0 (0.0)	---
Shortness of breath ^a	2 (9.1)	0 (0.0)	0.48	0 (0.0)	3 (15.0)	0.09	0 (0.0)	0 (0.0)	---

a= Qualitative variables are expressed as frequency and percentages. *p*-values with statistical significance are highlighted in bold. Chi square test o Fisher exact test was performed for comparison of qualitative variables $p \leq 0.05$.