

Supplementary material

related to: **Still's disease mortality trends in France, 1979-2016: a multiple-cause-of-death study**

by: Caroline Borciuch¹, Mathieu Fauvernier², Mathieu Gerfaud-Valentin¹, Pascal Sève^{1,3}, and Yvan Jamilloux^{1,4,5,*}

Figure S1. Distribution of deaths according to age in individuals aged under 35. The distribution of systemic-onset juvenile idiopathic arthritis (SJIA)-related deaths (black bars) and that in the general population (white bars) according to age is presented.




Table S1. Second non-underlying causes of death when SD was listed as the underlying cause of death, for the period 1979-2016.

	Men n=67	Women n=87	Total n=154
Infections	1 (1.5%)	1 (1.1%)	2 (1.3%)
Cardiovascular diseases	3 (4.5%)	3 (3.4%)	6 (3.9%)
Gastrointestinal diseases	0 (0)	0 (0)	6 (0)
Respiratory diseases	2 (3.0%)	1 (1.1%)	3 (1.9%)
Blood disorders	0 (0.0%)	1 (1.1%)	1 (0.6%)
Solid malignant neoplasms	2 (3.0%)	1 (1.1%)	3 (1.9%)
Unclassified above	12 (17.9%)	10 (11.5%)	22 (14.3%)
No 2 nd NUCD notification	59 (88.1%)	80 (92.0%)	139 (90.3%)