

Study or Subgroup	log[Hazard Ratio]	SE	Weight	Hazard Ratio	
				IV, Random, 95% CI	IV, Random, 95% CI
Aizawa 2014	0.58	0.31	7.8%	1.79	[0.97, 3.28]
Fukami 2017	-0.16	0.86	1.1%	0.85	[0.16, 4.60]
Garancini 2012	0.83	0.52	3.0%	2.29	[0.83, 6.35]
Kawahara 2020	0.52	0.66	1.9%	1.68	[0.46, 6.13]
Kinoshita 2015	0.18	0.21	14.7%	1.20	[0.79, 1.81]
Komeda 2014	-0.03	0.52	3.0%	0.97	[0.35, 2.69]
Liu J 2012	1.3	0.41	4.7%	3.67	[1.64, 8.20]
Liu Q 2015	0.43	0.97	0.9%	1.54	[0.23, 10.29]
Miki 2012	0.34	0.36	6.0%	1.40	[0.69, 2.85]
Nonaka 2019	-0.13	0.71	1.7%	0.88	[0.22, 3.53]
Oguro 2016	0.76	0.25	11.2%	2.14	[1.31, 3.49]
Song 2017	0.41	0.17	19.9%	1.51	[1.08, 2.10]
Tatsubayashi 2016	2.4	1.35	0.5%	11.02	[0.78, 155.39]
Tsujimoto 2010	0.17	0.21	14.7%	1.19	[0.79, 1.79]
Wang W 2014	0.9	0.4	5.0%	2.46	[1.12, 5.39]
Wang YN 2012	-0.12	0.47	3.7%	0.89	[0.35, 2.23]
Total (95% CI)			100.0%	1.54	[1.28, 1.85]
Heterogeneity: $\tau^2 = 0.01$; $\chi^2 = 16.86$, $df = 15$ ($P = 0.33$); $I^2 = 11\%$					
Test for overall effect: $Z = 4.64$ ($P < 0.00001$)					

