

Table S1. Differences in individual RNFL and mGCLT sectors between optic disc cohorts based on HRT division. Variable-wise Bonferoni-corrected p values $\leq .05$ are highlighted in bold. Median thickness values for groups 1, 2, 3 are given in μm .

	<i>Small vs medium</i>	<i>Medium vs large</i>	<i>Small vs large</i>	<i>median thickness small</i>	<i>median thickness medium</i>	<i>median thickness large</i>
<i>RNFL</i>						
NS	0.62	0.30	1.00	108.00	115.00	112.00
N	0.11	0.05	1.00	80.50	85.00	82.00
NI	0.42	1.00	0.43	103.50	109.00	112.00
TI	0.03	0.63	0.48	147.00	153.00	151.00
T	0.75	0.39	1.00	69.50	72.00	71.00
TS	0.06	1.00	0.22	130.50	137.00	138.00
<i>mGCLT</i>						
C	1.00	0.63	0.67	15.00	16.00	16.00
IN	0.46	1.00	1.00	52.50	53.00	53.00
ON	0.06	0.04	1.00	37.00	39.00	38.00
II	0.04	1.00	0.42	52.00	54.00	54.00
OI	0.04	0.09	1.00	32.00	34.00	33.00
IT	1.00	1.00	1.00	48.50	48.00	49.00
OT	0.05	1.00	0.36	36.00	37.00	37.00
IS	0.02	0.35	1.00	52.00	54.00	53.00
OS	0.01	0.17	0.80	34.00	35.50	35.00

RNFL sectors: NS = nasal superior; N = nasal; NI = nasal inferior; TI = temporal inferior; T = temporal; TS = temporal superior; mGCLT sectors: C = Central area; IN = Inner nasal; ON = Outer nasal; II = Inner inferior; OI = Outer inferior; IT = Inner temporal; OT = Outer temporal; IS = Inner superior; OS = Outer superior.