


# Supplementary Materials

**Table S1.** Effect size coefficients by clinical outcomes.

	Primary Outcome ( <i>n</i> = 759)	Secondary Outcome ( <i>n</i> = 294)	Tertiary Outcome ( <i>n</i> = 64)
	Effect Size	Effect Size	Effect Size
Age (years) <sup>a</sup>	0.013	0.061	0.137
Male sex, <i>n</i> (%) <sup>b</sup>	0.033	0.069	0.048
Hypertension, <i>n</i> (%) <sup>b</sup>	0.030	0.020	0.191
Diabetes mellitus, <i>n</i> (%) <sup>b</sup>	0.035	0.029	0.125
Bystander CPR, <i>n</i> (%) <sup>b</sup>	0.055	0.040	0.100
Witnessed, <i>n</i> (%) <sup>b</sup>	0.122	0.049	0.112
Initial shockable rhythm, <i>n</i> (%) <sup>b</sup>	0.099	0.190	0.457
Out-of-hospital CPR time <sup>a</sup>	0.069	0.022	0.171
In-hospital CPR time <sup>a</sup>	0.255	0.031	0.002
Total CPR time (min) <sup>a</sup>	0.227	0.039	0.090
Total epinephrine dose (mg) <sup>a</sup>	0.247	0.026	0.003
LDH (U/L), reference value (<290 U/L) <sup>a</sup>	0.054	0.064	0.013

<sup>a</sup>Eta squared. <sup>b</sup>Cramer's *v*.



**Figure S1.** (A) Primary outcome (ROSC). (B) Secondary outcome (Survival discharge). (C) Tertiary outcome (Favorable neurologic outcome).