Supplementary Table S1. Number of days individuals engaged in five NSSI behaviours in the last month.

	Scratch n = 48				Bruise n = 24				Cut n = 37				Burn n = 5				Bite n = 14			
	ED-R		ED-BP		ED-R		ED-BP n = 20		ED-R		ED-BP		ED-R		ED-BP		ED-R		ED-BP	
	<i>n</i> = 1 % a	n	<i>n</i> = 3 % a	n	% a	4 п	<i>n</i> = % a	n	n = 1	n	<i>n</i> = 2	n	<i>n</i> = %	n	n = 5 % a	п	<i>n</i> = 6 % a	п	<i>n</i> = 8	n
1–5 days	54.5%	6	67.6%	25	75%	3	75%	15	70%	7	51.9%	14	a	0	100%	5	66.7%	4	75%	6
6–10 days	27.3%	3	8.1%	3	0%	0	5%	1	10%	1	22.2%	6		0	0%	0	16.7%	1	12.5%	1
11-15 days	18.2%	2	8.1%	3	0%	0	15%	3	10%	1	7.4%	2		0	0%	0	0%	0	12.5%	1
>15 days	0%	0	16.2%	6	25%	1	5%	1	10%	1	18.5%	5		0	0%	0	16.7%	1	0%	0

Note. ^a Percentages of NSSI frequency per method within ED category; ED-R = eating disorder of the restrictive type; ED-BP = eating disorder of the bingeing/purging type.