



## Supplementary data

## Population Age < 65 Age 65 to 74 Age ≥ 75 979 225 p Value *n* = *n* = 615 n = *n* = 139 100% 62.8% 22.6% 14.2% **Peroperative Complications** 181 18.5% 17.1% 20.4% 30 21.6% 0.321 105 46 87 8.9% 47 7.6% 22 9.8% 12.9% Hemorrhage/ Transfusion 18 Digestive wound 11 1.1% 8 1.3% 0 0.0% 3 2.2% Vascular wound 6 0.6% 4 0.7% 1 0.4%1 0.7% 9 0.9% 0 0.0% Diaphragmatic wound/Pneumothorax 8 1.3% 1 0.4%5 0.5% 2 2 Ureteral or bladder wound 0.3% 1 0.4%1.4%2 2 12 1.2% 8 1.3% 0.9% Tumor rupture 1.4%**Postoperative Complications** 202 20.6% 131 21.3% 47 20.9% 24 17.3% 0.566 18 1.8% 9 1.5% 2 0.9% 7 5.0% Hemorrhage/ Transfusion Digestive (fistula, peritonitis, occlusion, 53 5.4% 36 5.9% 11 4.9% 6 4.3% pancreatitis) Urinary 19 1.9% 11 1.8% 5 2.2% 3 2.2% Lymphocele 19 1.9% 0 0.0% 13 2.1% 6 2.7% Abdominal wall (disunion, abscess, 33 3.4% 16 2.6% 12 5.3% 5 3.6% evisceration) 39 4.9% 5 4 2.9% Respiratory 4.0% 30 2.2% Pneumonia 10 1.0% 1.0% 1 0.4% 3 2.2% 6 Pneumothorax 4 0.4% 2 0.3% 2 0.9% 0 0.0% 9 2 0 Pulmonary embolism 11 1.1% 1.5% 0.9% 0.0% Pleural effusion 14 1.4% 13 2.1% 0 0.0% 1 0.7% Thrombosis 11 1.1% 1.0% 5 2.2% 0 0.0% 6 22 2 Infection (sepsis, pelvic abscess) 2.2% 16 2.6% 4 1.8%1.4%Neurological deficit 7 0.7% 5 0.8% 1 0.4% 1 0.7% **Repeat surgery** 51 5.2% 34 5.5% 12 5.3% 5 3.6% 0.649 **Overall Complications** 428 56% 259 54% 104 58% 65 61% 0.379 No 37% Minor complications 263 34% 178 53 30% 32 30% 9% 9% Major complications 73 10% 41 22 12% 10

## Table S1. Perioperative Complications.