

Article



Assessment of the Yam Landraces (*Dioscorea* spp.) of DR Congo for Reactions to Pathological Diseases, Yield Potential, and Tuber Quality Characteristics

Idris I. Adejumobi ^{1,2}, Paterne A. Agre ^{2,*}, Didy O. Onautshu ¹, Joseph G. Adheka ¹, Inacio M. Cipriano ¹, Jean-Claude L. Monzenga ³ and Joseph L. Komoy ¹

- Department of Biotechnology, Faculty of Science, University of Kisangani, Kisangani 2012, Congo; i.adejumobi@cgiar.org (I.I.A.); didy.onautshu@unikis.ac.cd (D.O.O.); jadheka@yahoo.fr (J.G.A.); inacio.cipriano@uem.ac.mz (I.M.C.); josephkomoy@gmail.com (J.L.K.)
- ² International Institute of Tropical Agriculture, Ibadan 5320, Nigeria
- ³ Institut Facultaire des Sciences Agronomiques de Yangambi, Kisangani 1232, Congo; claumonz@yahoo.fr
- Correspondence: p.agre@cgiar.org

Abstract: Yams (Dioscorea spp.) possess the potential to contribute to food security and poverty alleviation in DR Congo; however, yam production is limited by several constraints, including the lack of yam improvement programs to address challenges relating to yield improvement, resistance to foliar diseases, and post-harvest tuber quality. Identification of a superior genotype for these traits and reservoirs of genes for improvement would guide yams' improvement. This study aims to evaluate and identify landraces with superior performance for farmers and consumers. We evaluated 191 accessions from six yam species, and significant variation in the performances was observed at p < 0.05. Accessions of *D. alata* were superior for tuber oxidative browning (-0.01), *D. cayenensis* for high yield potential (29 t/ha), D. bulbifera for yam mosaic virus (YMV) tolerance (AUDPC = 3.88), and D. rotundata for tuber dry matter content (37%). A high genotypic and phenotypic coefficient of variation (>40) was observed for tuber yield, number of tubers per plots, tuber flesh oxidative browning, and tuber flesh texture. High broad-sense heritability estimates (>60) were similarly observed for all the assessed parameters except number of tubers per plot. Tuber size was identified as the best predictor for tuber yield (b = 2.64, p < 0.001) and tuber dry matter content (b = 2.21, p < 0.001). The study identified twenty stable landrace accessions from three *Dioscorea* species (D. alata (7); D. cayenensis (2); D. rotundata (11)). These accessions combined high yield potential, high tuber dry matter, high tolerance to YMV and YAD, and low tuber flesh oxidation. The accessions could be considered for the establishment of a yam improvement program in DR Congo.

Keywords: D. alata; D. bulbifera; D. cayenensis; D. dumetorum; D. praehensilis; D. rotundata; landraces

1. Introduction

Root and tuber crops make a significant contribution to global dietary needs after cereal crops [1]. Yam is among the principal root and tuber crops, including cassava and potato, that are widely grown and consumed as subsistence staples [2]. Yam is a generic name for the Dioscorea species widely cultivated in the tropics and subtropics by smallholder farmers mainly for its starchy underground tuber and aerial bulbils [3,4]. Thus, yam is a group of economically important multi-species crops that serve as a valuable source of food across Africa, Asia, South America, the Caribbean, and the Pacific [5,6]. The global estimated mean annually for yam production and gross values are approximately 73 million tons and 14 billion US dollars, respectively [7,8]. The genus *Dioscorea* has over 600 species, of which 11 are economically significant [9].

In DR Congo, yam is a major staple of the rural community, whose major occupation is farming, and is a scarce food commodity in the major city markets due to insufficient



Citation: Adejumobi, I.I.; Agre, P.A.; Onautshu, D.O.; Adheka, J.G.; Cipriano, I.M.; Monzenga, J.-C.L.; Komoy, J.L. Assessment of the Yam Landraces (*Dioscorea* spp.) of DR Congo for Reactions to Pathological Diseases, Yield Potential, and Tuber Quality Characteristics. *Agriculture* 2022, *12*, 599. https://doi.org/ 10.3390/agriculture12050599

Academic Editor: Francesco Vinale

Received: 27 February 2022 Accepted: 21 April 2022 Published: 24 April 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). production capacity. Of the economically significant species, seven have been previously reported to play a major role in subsistence livelihoods: white guinea yam (*D. rotundata*), yellow guinea yam (*D. cayenensis*), water yam (*D. alata*), bitter yam (*D. dumetorum*), bush yam (*D. praehensilis*), wild yam (*D. burkilliana*), and aerial yam (*D. bulbifera*) [10–12]. Many of these species are being cultivated under wide agro-ecological zones, though with higher preference for *D. rotundata* and *D. alata* [10,13].

Despite the importance of yam in sustaining rural livelihoods, yam production is faced with lots of constraints, including, but not limited to, biotic (pests and diseases), tuber quality (oxidative browning, dry matter, and taste), and agronomic (yield) constraints [10,14,15]. Of the biotic constraints, pests (nematodes, beetles, etc.) and two major foliar diseases (yam anthracnose disease (YAD) and yam mosaic virus disease (YMV)) are the major contributors to production loss. These foliar diseases have been reported by the yam scientific community as major pathological problems to yam productivity and have resulted in the loss of many traditional cultivars (landraces) in many yam-producing countries [6,14,16]. In DR Congo, the extent of affliction has over the time been aggravated by the absence of improved (resistant/tolerant) varieties of yams and the inability of subsistence farmers to afford the cost of adequate control measures.

Agronomic attributes, such as yield potential, tuber shape, and tuber quality characteristics (e.g., tuber dry matter content and oxidative browning), in general, play a major role in the acceptance of yam varieties by farmers and consumers. Thus, these attributes have most often been regarded as farmers' and consumers' preference criteria, upon which research has been focused in recent decades [1,15]. As in every other yam-producing country, yam farmers in DR Congo also prefer yam varieties characterized by a combination of marketable yield, sweet tuber taste, zero to minimal tuber flesh oxidative browning, high tuber dry matter content, and tolerance to yam foliar diseases [10]. These attributes are mostly combined in improved yam genotypes following years of breeding efforts. Obtaining such varieties is an impossibility for most farmers as they depend on local varieties (landraces) for seasonal cultivation. Though ennoblement efforts by a few farmers has helped in identifying very few landraces with good agronomic and tuber quality attributes, the majority of the farmers still lack access to seeds of these landraces [10–12].

Yam production constraints in DRC have been aggravated by the lack of yam improvement programs to address challenges relating to yield improvement, resistance to foliar diseases, and post-harvest tuber quality improvement. In the absence of structured yam improvement programs to enhance the genetic potential of the existing traditional cultivars, as well as to develop new and improved yam cultivars, an alternative way to contribute to the improvement of farmers' productivity will be to assess the existing traditional cultivars for the criteria that are of the utmost importance to the farmers and consumers. This will allow the identification of landraces that combine good agronomic, tuber quality, and disease resistance attributes, and thus they can be recommended to the farmers for cultivation through the Ministry of Agriculture. Therefore, this study was carried out to (i) identify landraces (cultivated and semi-wild species) with superior performance for yam foliar disease resistance, agronomic, and tuber quality traits and (ii) estimate the components of variance and heritability for the traits considered in the study for selection purposes in future yam improvement programs.

2. Materials and Methods

2.1. Experimental Site, Planting Materials, Experimental Layout, and Planting

The study was carried out at two research places of the University of Kisangani (UNIKIS), namely Simi-Simi (longitude 0°33′05.9″ N, latitude 25°05′17.3″ E, altitude 396 m a.s.l, elevation 397 m a.s.l) and Akodali (longitude 0°35′46.4″ N, latitude 25°08′56.6″ E, altitude 419 m a.s.l, elevation 428 m a.s.l), Kisangani, DR Congo. The duration of the field evaluation lasted 11 months from April 2020. The evaluation sites are characterized by the dense humid forest vegetation with an irregularly distributed rainfall pattern throughout the year (3156 mm annual). The soil type in both locations is mostly oxisols (ferralsols ac-

cording to FAO classification) [17], and the mean temperature range is 21–35 °C minimum and maximum temperatures, respectively.

The planting materials consisted of a panel of 191 genotypes (188 landraces and three breeding lines) across six species of *Dioscorea* (Table 1). The morphotypes within each species vary in quantity in the following order: *D. rotundata* (108), *D. alata* (33), *D. dumetorum* (16), *D. praehensilis* (16), *D. cayenensis* (12), and *D. bulbifera* (6). The landraces were sourced from six territories (Kisangani, Isangi, Bumba, Lisala, Buta, and Bambesa), categorized within three provinces (Tshopo, Mongala, and Bas-Uele). The breeding lines included as standard checks were obtained from the yam breeding unit (yam improvement program) of the International Institute of Tropical Agriculture (IITA), Ibadan, Nigeria. These standard checks were of the *D. rotundata* (TDr9519177 and TDr8902665) and *D. alata* (TDa1100316) species with known pathological, agronomic, and tuber quality attributes potential.

Table 1. List of the panel of 191 yam accessions used for the trial evaluation.

s/n	Accession Identity	Landrace Name	Territory	s/n	Accession Identity	Landrace Name	Territory	s/n	Accession Identity	Landrace Name	Territory
1	TDr21_001	Libanza-1	Bumba	32	TDr21_043	Moindo-1	Bumba	63	TDp21_052	Ahala-28	Bumba
2	TDr21_096	Moenge-1	Bumba	33	TDr21_067	Ahala-12	Bumba	64	TDr21_170	Bozongo-4	Bumba
3	TDr21_025	Ahala-1	Bumba	34	TDr21_027	Ahala-13	Bumba	65	TDp21_026	Ahala-29	Bumba
4	TDr21_141	Libanza-2	Bumba	35	TDr21_111	Ahala-14	Bumba	66	TDr21_112	Bozongo-5	Bumba
5	TDr21_010	Ahala-2	Bumba	36	TDr21_128	Ahala-15	Bumba	67	TDr21_074	Ahala-30	Bumba
6	TDr21_015	Moenge-2	Bumba	37	TDr21_016	Ahala-16	Bumba	68	TDr21_116	Bozongo-6	Bumba
7	TDr21_046	Libanza-3	Bumba	38	TDr21_044	Libanza-11	Bumba	69	TDr21_157	Ahala-31	Bumba
8	TDr21 158	Ahala-3	Bumba	39	TDr21 166	Libanza-12	Bumba	70	TDr21 187	Ahala-32	Bumba
9	TDr21 131	Libanza-4	Bumba	40	TDr21_097	Moenge-5	Bumba	71	TDr21_017	Ahala-33	Bumba
10	TDr21_177	Ahala-4	Bumba	41	TDc21_172	Bwanzele-2	Buta	72	TDr21_110	Ahala-34	Bumba
11	TDr21_179	Moenge-3	Bumba	42	TDr21_127	Ahala-17	Bumba	73	TDr21_020	Libanza-15	Bumba
12	TDr21_085	Libanza-5	Bumba	43	TDr21_012	Ahala-18	Bumba	74	TDr21_004	Ahenge-1	Bumba
13	TDc21_070	Bwanzele-1	Buta	44	TDr21 165	Ahala-19	Bumba	75	TDr21 167	Ahala-35	Bumba
14	TDr21_021	Wasalaka	Bumba	45	TDr21_006	Libanza-13	Bumba	76	TDr21_164	Libanza-16	Bumba
15	TDr21_186	Ahala-5	Bumba	46	TDr21_175	Moenge-6	Bumba	77	TDr21_031	Moenge-12	Bumba
16	TDr21 033	Bozongo-1	Bumba	47	TDr21 109	Ahala-20	Bumba	78	TDr21_087	Ahala-36	Bumba
17	TDa21_084	Ekolo-1	Kisangani	48	TDr21 161	Ahala-21	Bumba	79	TDr21 013	Ahenge-2	Bumba
18	TDr21 047	Bozongo-2	Bumba	49	TDr21 093	Moindo-2	Bumba	80	TDr21_082	Libanza-17	Bumba
19	TDr21 181	Moenge-4	Bumba	50	TDr21 105	Ahala-22	Bumba	81	TDr21 089	Ahala-37	Bumba
20	TDr21 154	Ahala-6	Bumba	51	TDr21 101	Moenge-7	Bumba	82	TDr21 183	Libanza-18	Bumba
21	TDr21 108	Libanza-6	Bumba	52	TDr21 129	Moenge-8	Bumba	83	TDr21 191	Ahala-38	Bumba
22	TDc21 117	Libanza-7	Bumba	53	TDr21 106	Moenge-9	Bumba	84	TDr21 030	Moenge-13	Bumba
23	TDr21 045	Ahala-7	Bumba	54	TDc21 190	Neboneboto-1	Buta	85	TDr21 155	Ahala-39	Bumba
24	TDr21_066	Ahala-8	Bumba	55	TDr21 024	Ahala-23	Bumba	86	TDr21 118	Ahala-40	Bumba
25	TDc21_059	Libanza-8	Bumba	56	TDr21_039	Ahala-24	Bumba	87	TDc21_091	Libanza-19	Bumba
26	TDr21_092	Libanza-9	Bumba	57	TDr21 113	Ahala-25	Bumba	88	TDr21_037	Ahala-41	Bumba
27	TDr21 119	Ahala-9	Bumba	58	TDr21_139	Ahala-26	Bumba	89	TDr21_057	Libanza-20	Bumba
28	TDr21_060	Abala-10	Bumba	59	TDr21_140	Libanza-14	Bumba	90	TDr21_143	Libanza-21	Bumba
29	TDr21_007	Bozongo-3	Bumba	60	TDr21_110	Ahala-27	Bumba	91	TDr21_148	Ahala-42	Bumba
30	TDr21_007	Libanza-10	Bumba	61	TDr21_171	Moenge-10	Bumba	92	TDr21_110	Engho	Bumba
31	TDr21_000	Abala-11	Bumba	62	$TDr21_101$	Moenge-11	Bumba	93	TDd21_112	Biamaiaune-1	Kisangani
94	TDr21_102	Ahala-43	Bumba	126	TDd21_104	Bilenge-2	Kisangani	158	TDa21_174	Ekolo-2	Kisangani
95	TDr21_100	Moenge-14	Bumba	120	TDd21_075	Bilenge-7	Isangi	159	TDn21_063	Lihoma	Isanoi
96	$TDr21_071$	Ahala-44	Bumba	127	TDd21_124	Bilenge-3	Kisangani	160	TDa21_000	Ekolo-3	Kisangani
97	TDr21_001	Moonge-15	Bumba	120	TDd21_009	Bilonge-4	Kisangani	161	$TD_{a}21_{-10}$	Ekolo-4	Kisangani
98	TDr21_051	Moenge-16	Bumba	129	TDd21_094	Celenge	Kisangani	162	TDa21_073	Ekolo-5	Kisangani
99	TDr21_103	Moenge-17	Bumba	131	TDa21_103	IFA Kis-4	Kisangani	163	$TD_{a}21_{0}75$	Ekolo-6	Kisangani
100	TDr21_134	Libanza-22	Bumba	132	TDa21_1008	Masua	Isangi	164	TDa21_111	Ekolo-7	Kisangani
101	TDr21_101	Abenge-3	Bumba	133	TDp21_000	Bainabaina	Isangi	165	TDa21_005	Ekolo-8	Kisangani
101	$TDr21_0000$	Abala-45	Bumba	134	$TDp21_001$ TDp21_049	Bosondi-3	Isangi	166	TDa21_000	Ekolo-9	Kisangani
102	$TDr21_041$ TDr21_115	Abulungenge-1	Bumba	135	$TDr21_049$	Abala-48	Bumba	167	$TD_{a}21_{004}$	Ekolo-10	Kisangani
103	$TDr21_102$	Ahulungenge-?	Bumba	136	TDd21_000	Bilonge-5	Kisangani	168	$TD_{a}21_{0}68$	Ekolo-11	Kisangani
105	TDr21_102	Abongo-4	Bumba	137	TDd21_100	Bwanzolo-3	Buta	160	$TD_{2}21_{0}000$	Ekolo-12	Kisangani
105	$TD_{r}^{21} 1002$	Allenge-4	Bumba	137	TDd21_011	Biamaiauno 5	Vicenceni	109	$TD_{a}21_{0}52$	Ekolo 16	Kisangani
100	TDr21_100	Ahulungenge 4	Bumba	120	TDd21_040	Biamajaune 6	Kisangani	170	$TD_{a}21_{0}00$	Ekolo 17	Kisangani
107	TDr21_000	Ahala-46	Bumba	140	$TDn21_029$	Bipaluma	Icangi	171	$TDn21_014$ TDn21_159	Bosondi-4	Kisangani
100	TD:21_120	Allala-40	Bumba	140	TD ₂ 21_020	Biamaruali 2	Visangani	174	TDp21_139	Privammala 4	Pambasa
109	$TDr21_100$ TDr21_160	Andid-4/	Bumba	141	$TD_{a}21_{072}$	Biamaiauno 7	Kisangani	170	$TDp21_182$	A dia	Bute
110	$TD_{r}21_{077}$	Abulungonge 4	Bumba	142	$TDd21_0/2$	Biamajaune-7	Kisangani	178	$TDp21_123$	Ambaga	Buta
111	$TD_{21}_{0/7}$	Ahanga E	Bumba	145	TD421_000	Piamajaune-8	Kisangani	170	$TDp21_036$	Ambaga Burangala 5	Duta
112	$1DC21_1/6$ TD=21_107	Anenge-5	bumba	144	TDb21_090	Diamajaune-9	Kisangani	1/9	TDe21_040	bwanzele-5	Duta
113	$1D121_107$ TDr21_127	IFA_KIS-1	Kisangani	145	TDb21_022	Litenu-1	Kisangani	100	$1D(21_{14})$ TD:21_122	Ekpore	Bambasa
114 115	TDr21_137 TDr21_055	IFA_KIS-2 IFA_Kis-3	Kisangani	146 147	TDb21_002 TDb21_086	Litehu-2 Litehu-3	Kisangani Kisangani	181	TDc21_132 TDc21_035	Ekpego Ngbongboto-4	Bambesa Bambesa

Accession

s/n	Accession Identity	Landrace Name	Territory	s/n	Identity	Landrace Name	Territory	s/n	Identity	Name	Territory
116	TDc21_138	Ngbongboto-2	Buta	148	TDb21_130	Litehu-4	Kisangani	183	TDc21_135	Manzaka	Buta
117	TDc21_188	Ngbongboto-3	Buta	149	TDa21_120	Ilumbelumbe-1	Kisangani	184	TDp21_121	Bwanzele-7	Buta
118	TDa21_149	Biamawali-1	Kisangani	150	TDa21_125	Inene-1	Kisangani	185	TDc21_018	Bwanzele-8	Buta
119	TDa21_189	Biamawali-2	Kisangani	151	TDa21_160	Ilumbelumbe-2	Kisangani	186	TDd21_146	Avuadipudi	Lisala
120	TDa21_095	Biamajaune-2	Kisangani	152	TDa21_173	Inene-2	Kisangani	187	TDp21_058	Mboloko	Lisala
121	TDa21_180	Biamajaune-3	Kisangani	153	TDr21_054	TDr8902665	IITÃ	188	TDp21_122	Mokongo	Lisala
122	TDp21_065	Bosondi-1	Isangi	154	TDr21_151	TDr9519177	IITA	189	Tda21_150	Maswe_1	Lisala
123	TDp21_078	Bosondi-2	Isangi	155	TDa21_042	TDa1100316	IITA	190	Tda21_192	Maswe_2	Lisala
124	TDd21_114	Biamajaune-4	Kisangani	156	TDd21_156	Bilenge-6	Kisangani	191	TDb21_023	Lihote	Lisala
125	TDd21_145	Bilenge-1	Kisangani	157	TDb21_076	Liseleka	Isangi				

The experiment was conducted in a 12 by 16 lattice design with two replicates. The field layout was generated using "Agricolae" package in R [18]. Each replicate was comprised of 16 incomplete blocks with 12 experimental plots. In each replicate, the experimental unit was comprised of 5 m long ridges containing five plants at 1 m intra- and inter-row spacing. The planting was done with yam setts ranging between 150 to 200 g each, treated using a cocktail of fungicide (Mancozeb 7.5 g/liter of water) and insecticide (Cypermethrin 7.5 mL/liter of water). Following the sprouting of the planted setts, the plants were exposed to natural field infestation of yam mosaic virus and yam anthracnose disease, and no fertilizers were applied during the evaluation process. Weeding was done manually when necessary.

2.2. Data Collection

Data were collected on the traits of economic significance to farmers and consumers (Table 2).

S/N	Trait	Nature of the Trait	Collection Period	Collection Method
1	Plant vigor	Visual assessment of the vigor of the vine and leaves of the new plant in a plot	4 MAP	Using a 1–3 scale where 1 = weak (75% of the plants or all the plants in a plot are small and have few leaves and thin vines), 2 = medium (intermediate or normal), and 3 = vigorous (75% of the plants or all the plants in a plot are robust, with thick vines and leaves very well developed or with abundant foliage)
2	Plant leaf density	Observation of variation in leaf mass or abundance on a mature plant and rating of density	4 MAP	Using a 1–7 scale where 3 = low, 5 = intermediate, and 7 = high
3	Senescence class	Visual observation of the status of foliage senescence (leaf or vine yellowing) of plants in plot at 6 months and onward and rating of the maturity class (status)	8 MAP	Using a 1–9 scale where 1 = very late (all the plants in a plot still show green foliage), 3 = late (75% of the plants in a plot still show green foliage, but up to 25% plants in a plot show leaves senescence), 5 = medium (50% of the plants still show green leaves and 50% showing senescence), 7 = early (75% of the plants in a plot show senescent foliage), and 9 = very early (all the plants in a plot are completely senesced).
4	Number of tubers per plot	The total quantity of the harvested tubers in a plot	At harvest	By count
5	Tuber size class	The average length of five tubers measured from the corm to the base in centimeters	At harvest	Using a 1–3 scale where 1 = small (less than 15 cm in length), 2 = medium (between 15 and 25 cm in length), and 3 = big/large (more than 25 cm in length)
6	Intensity of tuber flesh texture	The texture of tuber flesh after being cut	Post-harvest	Using a 1–3 scale where 1 = smooth, 2 = grainy, and 3 = very grainy
7	Intensity of tuber flesh oxidation	Degree of flesh surface color change or browning of cut yam tubers scored at different time intervals (0, 30, 60, 180 min)	Post-harvest	Using a 1–3 scale where 1 = no oxidization, 2 = slightly oxidizing, and 3 = highly oxidizing

Table 2. List of some traits assessed during the trial evaluation.

S/N	Trait	Nature of the Trait	Collection Period	Collection Method
8	YMV severity	Visual assessment of the grade of reaction of the plant to the virus infection, varying from mottle, mosaics until total leaf deformation, recording of the severity as a proportion or percentage of plant surface affected	2-6 MAS	Using a visual five ordinal scale (1–5 scale) where 1 = no visible symptoms; 2 = mosaic on few leaves, symptom recovery over time; 3 = mild symptoms on many leaves but no leaf distortion; 4 = severe mosaic on most leaves, leaf distortion; and 5 = severe mosaic (bleaching), severe leaf distortion and stunting
9	YADS severity	Visual assessment of anthracnose severity by observing the relative or absolute area of plant tissue affected by yam anthracnose disease and recording of the severity as a proportion or percentage of plant surface affected	2-6 MAS	Using a visual 1–5 general scale where 1 = no visible symptoms of anthracnose disease, 2 = few anthracnose spots or symptoms on 1 to ~25% of the plant, 3 = anthracnose symptoms covering ~26 to ~50% of the plant, 4 = symptoms on >51% of the plant, and 5 = severe necrosis and death of the plant

Table 2. Cont.

MAP = Month after planting; MAS = Month after sprouting.

The area under the disease progression curve (AUDPC), a valuable quantitative summary of disease severity for YMV and YAD over time, was estimated using the trapezoidal method [19]. This method discretizes the time variable and calculates the average disease severity between each pair of adjacent time points:

$$AUDPC = \sum_{i=1}^{N} \frac{(Yi + Yi + 1)}{2} (ti + 1 - ti)$$
(1)

where N is the number of assessments made, Y_i is the anthracnose or virus severity score on date *i*, and *t* is the time in months between assessments Y_i and Y_{i+1} .

Pathological reactions to yam mosaic virus (YMV) and to yam anthracnose disease (YAD) (severity scores) were recorded monthly from two to six months after sprout (Figure 1).





Figure 1. Visual scale for yam anthracnose (A) and yam mosaic virus diseases (B) scoring (pictures from Asfaw, 2016 [20]).

The plant vigor and leaf density were assessed at two and three months after sprout emergence, respectively. Senescence class, a measure of maturity class was assessed at six months after sprout emergence. Parameters used for yield assessment at harvest included number of tubers harvested per plot, tuber size category, and fresh tuber and/or bulbil weight per plot. The intensity of tuber flesh oxidation, tuber flesh texture, and tuber dry matter content in percentage were collected post-harvest. All the traits were assessed using the recommendations of Asfaw, 2016 [20] and yam crop ontology: https: //yambase.org/tools/onto/ (access on 25 February 2022).

Genotype fresh weight per plot was converted to the total tuber yield adjusted (*TTYA*) in tons per hectare using the formula below:

$$TTYA = \frac{TTWPx0}{PLS}$$
(2)

where TTWP is the total tuber weight per plot, and *PLS* is the plot size. Sett multiplication ration (*SMR*) was estimated as

$$SMR = \frac{\text{Weight of fresh tuber harvested}}{\text{Weight of sett planted}}$$
(3)

The dry matter content (*DMC*) was determined by grating 200 g of fresh tuber flesh into a container and oven-drying it at 120 $^{\circ}$ C for 48 h, at which constant weight was observed. The percentage dry matter content was estimated as

$$\% DMC = \frac{\text{Dry tuber flesh weight}}{\text{Wet tuber flesh weight}} \times 100$$
(4)

2.3. Statistical Analyses

A mixed linear model was used to conduct analysis of variance (ANOVA) using the lmerTest package in R [21] following the alpha lattice model below:

$$Y_{ijkl} = \mu + Gen_i + Rep_i + Rep(Blk)_{j(k)} + Env_l + Gen \times Env_{(il)} + Error_{ijkl}$$
(5)

where Y_{ijk} is the phenotypic performance of accession for traits under consideration, μ is the average accession performance, Gen_i is the effect of accession *i*, Rep_j is the effect of replication *j*, $Rep(Blk)_{j(k)}$ is the block *k* effect nested in replication *j*, Env_l is the effect of environment *l*, $Gen^*Env_{(il)}$ is the effect of the accession *i* by environment *l* interaction, and *Error*_{ijkl} is the residual effect.

For the analysis, accession (landrace) and environment were considered to be random effects while species was considered to be a fixed effect. Error (δ^2 e), genotypic (δ^2 g), and phenotypes (δ^2 p) variances were calculated from expected mean squares (EMS) of ANOVA following Kresovich, 1990 [22].

Error variance;

$$\delta^2 \mathbf{e} = \mathbf{MS}\mathbf{e},\tag{6}$$

Genotypic variance;

$$\delta^2 g = \frac{Msg - Msgl}{rl} \tag{7}$$

Genotypic by environment interaction variance;

$$\delta^2 g l = \left(\frac{Msg - Msgl}{lr}\right) \tag{8}$$

Phenotypic variance;

$$\delta^2 p = \delta^2_g + \left(\frac{\delta^2 e}{rl}\right) + \left(\frac{\delta^2 g l}{l}\right) \tag{9}$$

where, MSg = mean square of genotype; MSgl = mean square due to accession by environmental interaction; MSe = error mean square (mean square of environment); l = number of locations/environment; r = number of replications.

Broad-sense heritability (*H*2), phenotypic coefficient of variance (*PCV*), and genotypic coefficient of variance (*GCV*) were calculated using the values derived from respective variance components. Broad-sense heritability (*H*2) was classified as low (<30%), medium (30–60%), and high (>60%), according to Johnson et al. [23]. Following Deshmukh et al. [24],

phenotypic and genotypic coefficients of variation greater than 20% were rated as high, between 10 and 20% were rated as medium, and lower than 10% were regarded as low.

$$H2 = \frac{\delta^2 g}{\delta^2 g + \frac{\delta^2 gl}{l} + \frac{\delta^2 e}{rl}} \times 100$$
(10)

$$PCV = \left(\frac{\sqrt{\delta^2 p}}{\mu}\right) \times 100 \tag{11}$$

$$GCV = \left(\frac{\sqrt{\delta^2 g}}{\mu}\right) \times 100 \tag{12}$$

where $\delta^2 p$ = phenotypic variance, $\delta^2 g$ = genotypic variance, $\delta^2 g l$ = genotype by environment interaction variance; $\delta^2 e$: residual variance, r = number of replication; l = number of environment; μ : grand mean of the trait.

The relationship matrix, among the assessed traits, was constructed using Pearson's correlation coefficient and visualized using the ggpairs function in the GGally package [25]. Principal component analysis (PCA) was done using the PRCOMP function implemented in R [26] to identify the most discriminant traits with high contribution to the observed genotypic variation. Hierarchical cluster analysis was done based on the Ward.D2 method using the Gower dissimilarity matrix. The final hierarchical cluster was built and viewed using the dendextend package [27] and the circlize package [28] in R. The optimum number of clusters was identified using the NbClust package [29]. Path coefficient analysis was estimated and viewed using the lavaan function in the lavaan package [30]. In this model, tuber yield and tuber dry matter content were considered response variables against key agronomic and tuber quality traits as predictor variables. The path diagram was then constructed using the semPlot package [31] to depict the direct effect of these traits on tuber yield and dry matter content for suitability for indirect selection. Performance of landrace accession against check genotypes was assessed using Shukla's stability variance implemented in the VitSel application Version 1.0 [32].

3. Results

3.1. Variability in Agronomic and Tuber Quality Traits of Yam Landraces and Species

The analysis of variance (ANOVA) that shows the statistical difference for accessions and environment is presented in Table 3. Combined ANOVA revealed significant interaction effects of accession by environment at p < 0.05 for all the estimated parameters except for seed multiplication ratio, indicating environmental influence on the observed phenotypic expression of the landrace accessions for these traits. The interaction effect of species by environment was not significant for any parameter, suggesting that species performance was not environment dependent. Accession effect was significant at p < 0.001 for all the traits evaluated, indicating significant differences in the observed phenotypic performance of the accessions. Significant variation at p < 0.05 was observed for species effect in all the estimated parameters, indicating that the species performance differs for all the traits evaluated. Environment effect was significant for tuber dry matter content and yam anthracnose disease at p < 0.01, tuber oxidative browning at p < 0.05, and YMV severity at p < 0.001, indicating the existence of environmental differences with respect to these traits. Environment-specific analysis of variance revealed that both landrace and species effects were significant at p < 0.001 for all the studied traits in both environments.

Combined Environment													
Source	DF	Yield (t/ha)	DMC	SMR	NTPP	TUBSZE	TUBOXI	FLSTXT	YMV	YAD	LFDEN	PLTVIG	SENSC
Species	5	235.74 *	215.50 ***	162.74 ***	20.86 **	1.93***	6.47 ***	3.33 ***	1.64 ***	25.37 ***	3.69 ***	0.52 **	29.90 ***
Env	1	69.04	131.51 **	134.55	22.15	0.12	1.46 *	0.15	11.97 ***	13.16 **	1.83	0.19	5.92
Accession	177	397.72 ***	46.95 ***	95.02 ***	18.75 ***	0.79 ***	1.44 ***	0.45 ***	2.15 ***	2.79 ***	4.01 ***	0.75 ***	5.88 ***
Species * Env	5	50.85	15.53	11.28	7.22	0.09	0.28	0.00	0.07	0.46	0.19	0.19	1.75
Env * Accession	177	137.51 **	13.86 *	34.80	11.24 ***	0.19 ***	0.24 ***	0.13 **	0.47 ***	1.13 ***	1.54 ***	0.27 ***	1.29 *
Residual	335	91.71	11.01	31.02	6.54	0.12	0.15	0.09	0.31	0.53	0.77	0.13	0.99
CV%		48.60	9.11	61.27	76.97	13.75	79.55	20.76	10.43	11.91	15.79	15.95	26.73
Mean		20.49	35.35	9.20	3.35	2.58	0.48	1.41	5.23	6.30	5.60	2.31	3.75
					A	kodali enviro	onment						
Species	5	1205.61 ***	891.25 ***	596.08 ***	52.03 ***	4.84 ***	16.16 ***	18.66 ***	13.58 ***	32.20 ***	20.82 ***	3.11 ***	86.32 ***
Landrace	177	289.85 ***	29.01 ***	78.26 ***	11.70 ***	0.46 ***	0.89 ***	0.25 ***	1.16 ***	1.967 ***	2.84 ***	0.46 ***	3.22 ***
Residuals	152	100.06	9.00	32.51	3.52	0.12	0.19	0.10	0.30	0.46	0.63	0.11	1.20
CV%		47.36	8.33	57.96	49.98	13.3	84.06	22.07	10.94	10.33	13.47	13.34	28.88
Mean		21.36	36.09	9.97	3.70	2.61	0.52	1.40	5.06	6.56	5.88	2.44	3.83
					Sir	ni-Simi envii	ronment						
Species	5	878.19 ***	655.61 ***	413.15 ***	114.25 ***	7.11 ***	14.43 ***	18.53 ***	9.68 ***	36.83 ***	20.15 ***	4.23 ***	99.18 ***
Landrace	177	250.81 ***	30.21 ***	54.65 ***	17.51 ***	0.51 ***	0.76 ***	0.28 ***	1.33 ***	2.07 ***	2.81 ***	0.60 ***	4.55 ***
Residuals	152	83.40	12.76	26.30	9.54	0.14	0.13	0.09	0.30	0.59	0.89	0.15	0.80
CV%		45.96	10.34	60.19	106.97	14.49	77.71	20.52	10.19	12.69	17.69	17.37	24.32
Mean		19.62	34.60	8.43	2.97	2.56	0.45	1.41	5.41	6.03	5.32	2.18	3.68

Table 3. Combined and environment specific mean squares for agronomic and tuber quality traits in yam accessions.

DMC = Dry matter content; SMR = Sett multiplication ratio; NTPP = Number of tubers per plot; TUBSZE = Tuber size; TUBOXI = Intensity of tuber oxidation; FLSTXT = Tuber flesh texture; YMV = Yam mosaic virus disease; YAD = Yam anthracnose disease; LFDEN = Leaf density; PLTVIG = Plant vigor; SENSC = Senescence class; DF = Degree of freedom; CV = Coefficient of variation *, **, *** = significant at p < 0.05, 0.01, and 0.001 respectively.

Variation in the landrace species' mean performance in the combined analysis (Table 4) showed that *D. cayenensis* had the highest yield performance (28.49 t/ha) but was statistically similar to *D. alata* (25.72 t/ha) and significantly different from other species. *D. rotundata* had the highest dry matter content (37%), statistically similar to *D. cayenensis* (36.89%) and different from other species. *D. alata* had the highest number of *D. cayenensis* (11.73) but significantly different form other species. *D. alata* had the highest number of tubers per plot (4.55), similar to *D. bulbifera* but significantly different from other species. *D. alata* had the highest number of tubers per plot (4.55), similar to *D. bulbifera* but significantly different from other species. *D. cayenensis*, *D. rotundata*, *D. alata*, and *D. praehensilis* had significantly larger tuber size than the two other species. *D. alata* and *D. dumetorum* had the significantly minimal tuber flesh oxidative browning (-0.01 and -0.12, respectively), while *D. bulbifera* and *D. dumetorum* had significantly smoother flesh textures (0.90 and 0.94, respectively).

	Combined Environment											
Species	Yield (t/ha)	DMC	SMR	NTPP	TUBSZE	TUBOXI	FLSTXT	YMV	YAD	LFDEN	PLTVIG	SENSC
TDa	25.72 ^a	35.17 ^{bc}	13.90 ^a	4.55 ^a	2.65 ^a	-0.01 ^d	2.41 ^a	4.66 ^c	7.72 ^a	5.93 ^a	2.22 ^b	5.79 ^b
TDb	16.53 ^b	28.56 ^d	1.62 ^c	3.53 ^{ab}	1.17 ^c	0.71 ^{bc}	0.90 ^d	3.88 ^d	6.54 ^c	6.40 ^a	2.76 ^a	6.96 ^a
TDc	28.49 ^a	36.89 ab	11.73 ^a	3.04 ^b	2.86 ^a	1.07 ^b	1.20 bc	5.35 ^b	5.95 ^{cd}	6.03 ^a	2.53 ^a	2.12 ^e
TDd	15.71 ^b	25.84 ^e	7.98 ^b	2.71 ^b	2.00 ^b	$-0.12^{\text{ d}}$	0.94 ^d	4.49 ^c	7.17 ^b	5.66 ^{ab}	2.59 ^a	4.45 ^c
TDp	20.83 ^b	34.94 ^c	8.23 ^b	2.65 ^b	2.65 ^a	1.48 ^a	1.37 ^b	4.78 ^c	5.59 ^d	5.15 ^b	2.26 ^b	2.90 ^d
TDr	18.77 ^b	37.00 ^a	8.15 ^b	3.17 ^b	2.69 ^a	0.48 ^c	1.20 ^c	5.66 ^a	5.84 ^d	5.46 ^b	2.26 ^b	3.15 ^d
					Ako	dali environr	nent					
TDa	29.09 ^a	35.65 ^b	15.77 ^a	4.49 ^a	2.65 ^a	0.06 ^c	2.37 ^a	4.58 ^b	7.79 ^a	6.28 ^a	2.41 ab	5.80 ^b
TDb	17.83 ^{bc}	28.61 ^c	1.48 ^c	3.59 ^{ab}	1.26 ^c	0.61 ^b	0.85 ^d	3.78 ^c	6.92 ^{bc}	6.45 ^a	2.78 ^a	7.07 ^a
TDc	29.34 ^a	38.28 ^a	12.16 ab	3.24 ab	2.82 ^a	1.32 ^a	1.22 ^c	5.20 ^a	6.30 ^{cd}	6.36 ^a	2.57 ^{ab}	2.67 ^d
TDd	16.74 ^c	25.91 ^c	8.47 ^b	2.67 ^b	1.99 ^b	-0.16 ^c	0.98cd	4.48 ^b	7.72 ^{ab}	5.82 ^{ab}	2.66 ^a	4.55 ^c
TDp	21.53 ^b	34.90 ^a	8.76 ^b	2.95 ^ь	2.63 ^a	1.51 ^a	1.47 ^b	4.65 ^b	5.92 ^d	5.21 ^b	2.33 ^b	3.32 ^d
TDr	18.90 bc	37.97 ^a	8.77 ^b	3.76 ab	2.74 ^a	0.51 ^b	1.18 ^c	5.41 ^a	6.07 ^{cd}	5.76 ^{ab}	2.40 ab	3.13 ^d
					Simi-	Simi environ	ment					
TDa	22.36 ab	34.69 ^a	12.03 ^a	4.61 ^a	2.66 ^a	-0.08 d	2.44 ^a	4.73 ^b	7.64 ^a	5.56 ^{ab}	2.03 c	5.77 ^b
TDb	15.24 bc	28.51 ^b	1.75 ^c	3.46 ab	1.07 ^c	0.80 bc	0.95 ^{bc}	3.98 ^c	6.15 ^{bc}	6.35 ^a	2.75 ^a	6.85 ^a
TDc	27.63 ^a	35.51 ^a	11.3 a	2.84 ab	2.89 a	0.81 ^b	1.17 ^b	5.50 ^a	5.59 °	5.70 ^{ab}	2.48 ab	1.57 ^e
TDd	14.68 ^c	25.76 ^b	7.49 ^b	2.75 ab	2.02 ^b	$-0.09^{\text{ d}}$	0.88 ^c	4.50 bc	6.62 ^b	5.49 ^{ab}	2.52 ab	4.35 ^c
TDp	20.12 abc	34.98 a	7.69 ^b	2.35 ^b	2.68 a	1.44 ^a	1.27 ^b	4.90 ^b	5.26 ^c	5.07 ^b	2.20 bc	2.48 ^d
TDr	18.64 ^{bc}	36.02 ^a	7.52 ^b	2.58 ^b	2.64 ^a	0.46 ^c	1.21 ^b	5.91 ^a	5.60 ^c	5.16 ^b	2.11 ^c	3.18 ^d

Table 4. Mean variations in agronomic and tuber quality traits of yam genotypes based on landrace species.

DMC = Dry matter content; SMR = Sett multiplication ratio; NTPP = Number of tubers per plot; TUBSZE = Tuber size; TUBOXI = Intensity of tuber oxidation; FLSTXT = Tuber flesh texture; YMV = Yam mosaic virus disease; YAD = Yam anthracnose disease; LFDEN = Leaf density; PLTVIG = Plant vigor; SENSC = Senescence class. TDa = Tropical *Dioscorea alata*; TDb = Tropical *Dioscorea bulbifera*; TDc = Tropical *Dioscorea cayenensis*; TDd = Tropical *Dioscorea fraehensilis*; TDr = Tropical *Dioscorea rotundata*. The letters a, b, c, d & e represent the LSD level of significance.

Response to pathological disease revealed that *D. bulbifera* had the highest tolerance to YMV severity (AUDPC = 3.88), while *D. rotundata* had the least tolerance (AUPDC = 5.66). However, *D. praehensilis* (AUDPC = 5.59) and *D. rotundata* (AUDPC = 5.84) had the highest tolerance to YAD severity, while *D. alata* had the least tolerance (AUDPC = 7.72). *D. bulbifera*, *D. cayenensis*, and *D. alata* had significantly higher leaf density (6.40, 6.30, and 5.93, respectively), while *D. bulbifera*, *D. dumetorum*, and *D. cayenensis* had significantly better plant vigor (2.76, 2.59, and 2.53, respectively). *D. bulbifera* had significantly higher senescence class (6.96 = early maturing), while *D. cayenensis* had the lowest rating (2.12 = very late maturing) (Table 4).

3.2. Genetic Variability and Broad-Sense Heritability of Agronomic and Tuber Quality Traits Yam Accessions

Genotypic and phenotypic variance components, genotypic and phenotypic coefficients of variation, and broad-sense heritability of agronomic and tuber quality traits in yam accessions are presented in Table 5. Genotypic coefficients of variation (GCV) ranged from a moderate classification of 12.96% for tuber dry matter content to high classification 52.16% for tuber flesh oxidative browning. A similar result was observed for phenotypic coefficients of variation (PCV) which ranged from a moderate classification of 14% for tuber dry matter content to a high classification of 68.49% for the number of tubers per plant. Broad-sense heritability (H²) varied between 46.97% (moderate) and 91.40% (high). High H² (>60%) was observed in all the estimated parameters except for number of tubers per plot, where moderate H² was observed.

	Genetic Parameters										
Traits	$\delta^2 g$	$\delta^2 p$	GCV (%)	PCV (%)	H ² (%)						
Yield (t/ha)	80.20	114.94	43.71	52.32	69.78						
DMC	21.00	24.48	12.96	14.00	85.77						
SMR	22.43	31.03	51.50	60.57	72.30						
NTPP	2.45	5.22	46.92	68.49	46.97						
TUBSZE	0.25	0.30	19.34	21.19	84.12						
TUBOXI	0.53	0.59	52.16	54.86	89.45						
FLSTXT	0.35	0.39	42.08	44.42	91.40						
YMV	0.60	0.72	14.80	16.22	83.93						
YAD	0.86	1.15	14.73	17.03	74.34						
LFDEN	0.81	1.19	16.07	19.48	67.85						
PLTVIG	0.16	0.23	17.31	20.75	69.00						
SENSC	2.76	3.10	44.25	46.90	88.97						

Table 5. Genetic variance, coefficient of variation, and broad-sense heritability in yam landrace accessions.

 $\delta^2 g$ = Genotypic variance; $\delta^2 p$ = Phenotypic variance; GCV = Genotypic coefficient of variation; PCV = Phenotypic coefficient of variation; H² = Broad-sense heritability; DMC = Dry matter content; SMR = Sett multiplication ratio; NTPP = Number of tubers per plot; TUBSZE = Tuber size; TUBOXI = Intensity of tuber oxidation; FLSTXT = Tuber flesh texture; YMV= Yam mosaic virus disease; YAD = Yam anthracnose disease; LFDEN = Leaf density; PLTVIG = Plant vigor; SENSC = Senescence class.

3.3. Principal Component Analysis of Agronomic and Tuber Quality Traits

The principal component analysis that was used to identify the most discriminant traits with high contributions to the observed genotypic variation is presented in Table 6. The first four principal components (PC), with Eigen values greater than one, accounted for 66.21% of the genetic variation in the study. The first PC accounted for 28.49% of variance, with major contributions from tuber yield, seed multiplication ratio, tuber size, leaf density, and plant vigor. The second PC accounted for 16.39%, with major contributions from number of tubers per plot, tuber size, tuber flesh texture, YAD severity, and senescence class. The third PC accounted for 12.68%, with major contributions from tuber dry matter content, tuber size, tuber flesh oxidative browning, YMV severity, and plant vigor. The fourth PC accounted for 8.65%, with major contributions from tuber dry matter content, tuber size, tuber flesh oxidative browning, tuber flesh texture, and YMV severity (Table 6).

Table 6. Principal component analysis and contributions of agronomic and tuber quality traits to the genetic variability.

Trait	PC1	PC2	PC3	PC4
Yield (t/ha)	0.457	-0.182	0.003	-0.120
DMC	0.011	-0.232	-0.437	0.252
SMR	0.455	-0.076	-0.123	-0.158
NTPP	0.166	0.287	-0.110	0.134
TUBSZE	0.316	-0.264	-0.322	0.227
TUBOXI	0.076	-0.162	0.327	0.762
FLSTXT	0.196	0.289	-0.391	0.345
YMV	-0.117	-0.221	-0.506	-0.284
YAD	0.220	0.472	-0.123	-0.016
LFDEN	0.452	0.019	0.142	-0.128
PLTVIG	0.382	-0.065	0.352	-0.177
SENSC	-0.003	0.608	-0.052	0.009
Eigen value	1.849	1.402	1.234	1.019
Variance (%)	28.490	16.390	12.680	8.654
Cumulative (%)	28.490	44.880	57.560	66.213

DMC = Dry matter content; SMR = Sett multiplication ratio; NTPP = Number of tubers per plot; TUBSZE = Tuber size; TUBOXI = Intensity of tuber oxidation; FLSTXT = Tuber flesh texture; YMV = Yam mosaic virus disease; YAD = Yam anthracnose disease; LFDEN = Leaf density; PLTVIG = Plant vigor; SENSC = Senescence class. PC1 to PC12 indicate Principal Components.

3.4. Relationships among Agronomic and Tuber Quality Traits in Yam Landraces

The relationship among evaluated yam parameters is presented in Figure 2. A significant positive relationship was observed between tuber yield and sett multiplication ratio (r = 0.79), tuber size (r = 0.49), plant leaf density (r = 0.63) at p < 0.001, tuber flesh texture (r = 0.22) at p < 0.01, and number of tubers per plot (r = 0.16) at p < 0.05. A significant negative relationship was not observed. Dry matter content showed a significant positive relationship with tuber size (r = 0.28) and YMV severity (r = 0.41) at p < 0.001, while a negative relationship was observed for number of tubers per plot (r = -0.15, p < 0.05), YAD severity (r = -0.20), and leaf density (r = -0.23) at p < 0.01. A significant positive relationship indicates similar direction in trait performance, while a significant negative relationship indicates opposite direction in traits expression.

		25 30 35 40		2 4 6 8 10		0.0 1.0 2.0		56789		4.0 5.0 6.0	
	Yieldt.ha.	-0.04	0.79***	0.16*	0.49***	0.07	0.22**	-0.13	0.14	0.63***	10 30 50
25 35		DMC	0.01	-0.15*	0.28***	0.00	-0.05	0.41***	-0.20**	-0.23**	
			SMR	0.27***	0.45***	-0.05	0.34***	-0.04	0.30***	0.59***	5 15
2468				NTPP	0.08	0.00	0.39***	-0.06	0.26***	0.31***	
					TUBSZE	0.11	0.32***	0.15*	0.06	0.31***	1.5 2.5
0.0 1.0 2.0	- 888 - 96 - 98 - 8 - 8 - 8 - 8 - 8 - 8 - 8 - 8 - 8 -	000 80 000 00 000 00 000 00 000 00 000 00	6000 0000 0000 0000 0000 0000 0000 000			TUBOXI	-0.01	-0.25***	-0.14	0.14	
			0 0000 0 0000 0 0000 0 0000 0 0000 0 0000 0				FLSTXT	-0.10	0.42***	0.25***	1.0 2.0
5 7 9							0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	YMV	-0.18*	-0.10	
				8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8					YAD	0.30***	5 7 9
4.0 5.5				600 C						LFDEN	
	10 30 50		5 10 15 20		1.5 2.0 2.5 3.0		1.0 1.5 2.0 2.5		56789		

Figure 2. Correlation coefficients among agronomic and tuber quality traits. DMC = Dry matter content; SMR = Sett multiplication ratio; NTPP = Number of tubers per plot; TUBSZE = Tuber size; TUBOXI = Intensity of tuber oxidation; FLSTXT = Tuber flesh texture; YMV = Yam mosaic virus disease; YAD = Yam anthracnose disease; LFDEN = Leaf density; PLTVIG = Plant vigor; SENSC = Senescence class. *, **, *** = significant at p < 0.05, 0.01, and 0.001 respectively.

3.5. Yam Clustering Based on Hierarchical Clustering

Hierarchical clustering employed for the grouping of yam accessions based on the evaluated agronomic and tuber quality characters produced four clusters (Figure 3). Cluster one consisted of accessions of *D. alata* (30), characterized by high tuber yield, a high number of tubers per plot, high leaf density, large tuber size, low tuber flesh oxidative browning, very grainy tuber flesh texture, high susceptibility to YAD severity, and medium senescence

class. Cluster two had the largest cluster membership, which consisted of accession of *D. rotundata* (42), *D. praehensilis* (13), *D. cayenensis* (11), *D. alata* (1), and *D. dumetorum* (1), characterized by high yield, high dry matter content, large tuber size, high leaf density, high plant vigor, smooth tuber flesh texture, and moderate tuber oxidative browning. Cluster three had the minimum cluster members and consisted of accessions of *D. dumetorum* (13), *D. bulbifera* (6), and *D. alata* (3), characterized by a high number of tubers per plot, high leaf density, high plant vigor, low tuber flesh oxidative browning, smooth tuber flesh texture, early senescence class, and moderate tolerance to YMV severity but susceptibility to YAD severity. Cluster four consisted of accessions of *D. rotundata* (60) and *D. cayenensis* (1), characterized by high dry matter content, low tuber flesh oxidative browning, smooth tuber flesh textures, and susceptibility to YMV severity with moderate tolerance of YAD severity (Table 7).



Figure 3. Hierarchical clustering showing grouping patterns of yam landrace accessions into four clusters using twelve key traits covering agronomic and tuber quality based on the Gower dissimilarity matrix. C1, Cluster one (blue); C2,; Cluster two (yellow); C3, Cluster three (green); C4, Cluster four (red). D. praehensilis (2).

Table 7. Description of clusters of yam landraces.

Trait	Cluster 1 (30)	Cluster 2 (68)	Cluster 3 (22)	Cluster 4 (63)	F-Value
Tuber yield (t/ha)	23.90 ^a	24.79 ^a	18.41 ^b	15.01 ^c	33.31 ***
Dry matter content (%)	35.57 ^b	36.32 ^{ab}	27.26 ^c	37.02 ^a	62.88 ***
Sett multiplication ratio	12.56 ^a	10.85 ^b	7.98 ^c	6.25 ^d	36.61 ***
Number of tuber per plot	3.99 ^a	3.35 ^b	3.55 ^{ab}	2.95 ^c	7.68 ***
Tuber size	2.87 ^a	2.82 ^a	1.91 ^c	2.43 ^b	50.11 ***
Tuber oxidative browning	0.26 ^b	0.88 ^a	0.31 ^b	0.22 ^b	14.72 ***
Tuber flesh texture	2.49 ^a	1.23 ^b	1.12 ^b	1.18 ^b	154.52 ***
Yam mosaic virus disease	5.01 ^b	5.27 ^b	4.50 ^c	5.56 ^a	16.55 ***
Yam anthracnose disease	7.12 ^a	6.06 ^b	6.83 ^a	5.98 ^b	28.63 ***
Leaf density	6.01 ^a	5.91 ^a	6.03 ^a	4.93 ^b	47.36 ***
Plant vigor	2.36 ^b	2.49 ^a	2.54 ^a	2.02 ^c	48.68 ***
Senescence class	5.08 ^a	2.85 ^c	5.72 ^a	3.42 ^b	48.84 ***

Significance level: "p < 0.001" = ***. Means followed by the same superscripts are not significantly different using the least significant difference (LSD) test at a 5% p-value threshold. The bold values indicate significant traits at each cluster. The letters a, b & c represent the LSD level of significance.

3.6. Path Analysis among Assessed Traits of Dioscorea Species

The path analysis done to depict the direct effect of agronomic traits on tuber yield and dry matter content for suitability for indirect selection is presented in Figure 4. The path analysis began with structural equation modelling where tuber yield and dry matter content were considered response variables against correlated agronomic and tuber quality parameters. The model resulted in excellent fit. The chi-square test of the model fit was not significant (χ^2 (4) = 2.455, *p* = 0.653). Overall, fit indices were in good range (RMSEA = 0.00 [0.00, 0.09], *p* = 0.81; CFI = 1.00; SRMR = 0.01). Most of the direct effects in the model were significant.



Figure 4. Path coefficient analysis between response and independent yam variables. Y.. = Tuber yield; DMC = Dry matter content; SMR = Sett multiplication ratio; NTP = Number of tubers per plot; TUB = Tuber size; TUBO = Tuber oxidation; FLS = Tuber flesh texture; YMV = Yam mosaic virus disease; YAD = Yam anthracnose disease; LFD = Leaf density; PLT = Plant vigor; SEN = Senescence class. Red indicates direct negative impact, and green indicates direct positive impact.

Setts multiplication ratio significantly predicted tuber yield (b = 1.12, SE = 0.10, p < 0.001) such that a unit increase in setts multiplication ratio was associated with a 1.12-unit increase in tuber yield. Tuber size significantly predicted tuber yield (b = 2.64, SE = 0.83, p < 0.001) such that a one-unit increase in tuber size was associated with a 2.64-unit increase in tuber yield. YMV severity significantly predicted tuber yield (b = -1.21, SE = 0.45, p < 0.01) such that a one-unit increase in YMV severity was associated with a 1.21-unit decrease in tuber yield. Leaf density significantly predicted tuber yield (b = 2.08, SE = 0.65, p < 0.001) such that a one-unit increase in leaf density was associated with a 2.08-unit increase in tuber yield (Figure 3).

Tuber size significantly predicted tuber dry matter content (b = 2.21, SE = 0.60, p < 0.001) such that a one-unit increase in tuber size was associated with a 2.21-unit increase in tuber dry matter content. YMV severity significantly predicted tuber dry matter content (b = 1.54, SE = 0.39, p < 0.001) such that a one-unit increase in YMV severity was associated with a 1.54-unit increase in tuber dry matter content. Plant vigor significantly predicted tuber dry matter dry matter content (b = -3.70, SE = 0.84, p < 0.001) such that a one-unit increase in plant vigor was associated with a 3.70-unit decrease in tuber dry matter. Senescence class significantly predicted tuber dry matter content (b = -0.60, SE = 0.17, p < 0.001) such that a one-unit increase in senescence class was associated with a 0.60-unit decrease in tuber dry matter content.

3.7. Performance of Landrace Accession against Standard Check Genotypes

Landraces performances for traits of interest were compared to that of the average performance of the three checks used in the study. Tuber yield and tuber dry matter were set as traits of higher values, and AUDPC estimates for YMV and YADS were set as traits of lower values, while tuber flesh oxidation was set as trait of values within range. The performance of landrace accession against the standard check genotypes revealed that 51 landrace accessions with Shukla's stability variances varying from 0 to 389 had better performance than the average performance of the three checks included in the study (Table S1). Of the 51 landraces accessions, only 20 accessions were observed to have stable performance (Shukla's stability variance of less than 5) with respect to the parameters under assessment. The 20 accessions were observed to be accessions of *D. alata* (TDa21_169; TDa21_080; TDa21_73; TDa21_152; TDa21_050; TDa21_005; TDa21_034), *D. cayenensis* (TDc21_059; TDc21_190), and *D. rotundata* (TDr21_162; TDr21_089; TDr21_142; TDr21_037; TDr21_153; TDr21_154; TDr21_134; TDr21_131; TDr21_163; TDr21_099) (Table 8).

Table 8. List of the stable twenty landrace accessions with better performance over checks mean for farmers' and consumers' preferred traits.

S/N	Genotype	Yield	DMC	YMV	YAD	TUBOXI	Stability	Rank
1	TDa21_169	25.88	38.43	5.68	6.89	0.80	0.00	0
2	TDr21_162	21.19	43.40	5.85	5.44	0.05	0.00	0
3	TDc21_059	19.30	38.00	6.32	6.11	0.06	0.15	7
4	TDr21_089	21.25	41.43	5.87	6.08	0.95	0.17	11
5	TDr21_142	17.74	38.54	5.02	6.12	0.05	0.18	13
6	TDr21_037	21.58	36.58	5.86	6.70	0.05	0.35	19
7	TDa21_080	21.35	36.10	5.89	6.89	0.05	0.46	22
8	TDr21_153	18.61	40.32	5.86	6.87	0.05	0.46	24
9	TDr21_167	24.39	37.23	5.84	6.12	0.95	0.48	25
10	TDr21_154	25.52	36.30	5.87	5.96	0.27	1.18	46
11	TDc21_190	33.08	36.54	6.71	5.40	0.05	1.59	52
12	TDr21_134	20.11	39.09	6.71	5.47	0.05	1.60	54
13	TDa21_073	23.35	38.21	5.47	6.56	0.05	1.72	57
14	TDa21_152	23.84	38.05	5.05	6.90	0.28	2.06	64
15	TDr21_131	25.25	39.39	5.03	5.38	0.05	2.18	67
16	TDa21_050	19.85	36.74	5.47	6.71	0.05	2.32	68
17	TDa21_005	21.75	38.87	5.26	7.12	0.13	2.98	75
18	TDr21_163	33.02	36.54	5.04	6.22	0.05	3.05	77
19	TDa21_034	24.09	37.78	5.26	6.90	0.20	3.16	78
20	TDr21_099	35.81	37.57	5.04	6.96	0.95	3.46	80
	Checks mean	17.40	35.42	6.72	7.21			

DMC = Dry matter content; TUBOXI = Intensity of tuber oxidation; YMV = Yam mosaic virus disease; YAD = Yam anthracnose disease; Stability = Shukla's stability variance.

4. Discussion

4.1. Variability in Agronomic and Tuber Quality Traits of Dioscorea Species as Identifiers of Gene Reservoirs for Yam Genetic Improvement in DR Congo

Yam production in DR Congo is challenged by numerous constraints, including, but not limited to, low yield, poor tuber quality characteristics, and pathological diseases, which have been the major focus of modern breeding programs in countries where they exist. The identification of the genetic potential and gene reservoir for genetic improvement from the existing genetic pool of landraces for high yield potential, good tuber quality attributes, and resistance/tolerance to pathological diseases could offer a potential hope for consideration for yam improvement. The study revealed varying degrees of potential of the landrace species for farmers' and consumers' preferred traits (high yield, high dry matter content, resistance to pathological diseases, and non to low tuber oxidative browning). Accessions of *D. cayenensis* and *D. alata* had the highest yield potential among all the species considered. These species are popular for their high plant vigor and leaf density, which enhance the yield potentials, hence the reason for the wide distribution of *D. alata* worldwide [1,33]. In addition, *D. cayenensis* requires a longer cycle, which is not a desired trait to many yam cultivators but allows the advantage of more assimilates production and translocation into the tubers compared to most cultivated species. Accessions of *D. rotundata* had the highest dry matter content and were probably the reason for the preference for consumption and industrial potential in many yam-producing communities [34,35]. Accessions of *D. alata* and *D. dumetorum* had very low tuber flesh oxidative browning properties compared to other species. This trait has been reported as a determinant in yam cultivar acceptability in many studies [15].

For resistance to pathological diseases, accessions of *D. bulbifera* had the best genetic tolerance to YMV disease; however, this species is not known for regular cultivation, as it is regarded as forest/wild species [36,37]. Of the cultivated species, accessions of *D. dumetorum*, *D. alata*, and *D. praehensilis* had better tolerance than the popular preference species for consumption (*D. rotundata*) and, as such, can be considered in breeding programs for the improvement of *D. rotundata*, particularly *D. praehensilis* due to their similar genome information [38]. This corroborates the findings of Adewumi et al. [39], who observed better tolerance of *D. praehensilis* genotypes over that of *D. rotundata* for YMV severity. Accessions of *D. praehensilis* and *D. rotundata* had the best genetic tolerance to YAD severity among the considered species while *D. alata* had the least tolerance. *D. alata* is very susceptible to YAD severity [16,40,41].

4.2. Genetic Parameters and Broad-Sense Heritability of Evaluated Traits

The high GCV and PCV (>20%) observed in some of the evaluated traits, such as tuber yield, seed multiplication ratio, number of tubers per plot, tuber flesh oxidative browning, tuber flesh texture, and senescence class indicates potentials for high selection intensity. This is essential, as it will facilitate the selection of accessions with superior performance in yam breeding programs. High GCV and PCV recorded for tuber yield in this study were in agreement with the findings of Padhan and Panda [42] conducted on advanced breeding populations of white yam. High H² (>60%) recorded in this study for all traits except for number of tubers per plot indicates a high correspondence between phenotypic and genotypic variance and, therefore, high response to selection. Many studies have also obtained similar findings for some of the observed parameters. Agre et al. [1] observed high H² estimates for tuber yield per plant and YMV in *D. rotundata*. Bhattacharjee et al. [16] also reported high broad-sense heritability for YAD in *D. alata*.

4.3. Correlation Coefficients, Principal Components, and Hierarchical Clusters among Assessed Traits of Landrace Accessions

Landraces accession with high leaf density, large tuber size, high seed multiplication ratio, grainy flesh texture, and high number of tubers per plot could be considered in breeding for improved yield following their observed relationship in the study. In consideration for improved tuber dry matter content, landrace accessions with a reduced number of tubers per plot, larger tuber size, tolerance to YAD severity, and reduced leaf density could be considered following the relationship observed in our study. Agre et al. [43] similarly observed a positive relationship between tuber yield and yield components, such as tuber size and number of tubers per plot in a panel of water yam.

The traits that best discriminated the landrace accession in this study were those which resolved on PC1 with major contribution. These traits, including tuber yield, tuber size, leaf density, plant vigor, YMV severity, YAD severity, and flesh texture could be utilized in evaluating genetic diversity among similar species of yam. Agre et al. [43,44] and Siadjeu, et al. [45] have previously reported the significant contribution of the majority of these traits in discriminating yam accessions.

The hierarchical clustering revealed genetic similarities among landraces accessions that were grouped in the same cluster. Clustering of *D. rotundata*, *D. praehensilis*, and *D. cayenensis* accessions in clusters two and four corroborates the findings of Scarcelli et al. [38], who

reported *D. praehensilis* as the progenitor of *D. rotundata*. This also suggests the existence of a possible genetic relationship between *D. rotundata* and *D. cayenensis*. Many studies have supported this theory [46,47] and, as such, called it the *D. cayenensis-rotundata* complex. From the clustering, *D. alata* showed characteristics for high tuber yield, high setts multiplication ratio, higher number of tubers per plot, larger tuber size, low tuber oxidative browning, high leaf density, moderate resistance to YMV severity, and early maturity. *D. rotundata*, *D. cayenensis*, and *D. praehensilis* showed characteristics for high yield, high dry matter content, large tuber size, better resistance to YAD severity, high leaf density, high plant vigor, and late maturity. *D. bulbifera* and *D. dumetorum* showed characteristics for a higher number of tubers per plot, leaf density, plant vigor, and early maturity.

4.4. Traits Prediction (Indirect Selection) for Yield and Tuber Quality Attributes

One of the challenges in yam improvement is long growing cycles of the genotypes. Thus, any means to select for improving yield and good tuber quality characteristics in yam accessions using agronomic characteristics (indirect selection) will be of advantage. Our study suggests that low YMV severity, leaf density, tuber size, and setts multiplication ratio predict tuber yield, while tuber size, plant vigor, YMV severity, and senescence class predict tuber dry matter content. Of the observed predictor traits, tuber size and YMV severity predict both tuber yield and tuber dry matter content.

5. Conclusions

This study explored a panel of 191 yam accessions within six *Dioscorea* species for the identification of superior genotypes for farmers' and consumers' preferred traits (tuber yield, tuber dry matter, YMV severity, YAD severity, and tuber flesh oxidative browning). We observed variations in the performance of the landrace species with respect to all the agronomic and tuber quality traits assessed in the study. All the assessed parameters have the moderate to high heritability necessary for response to selection. We observed significant relationships among the assessed traits and paths, and coefficient analysis revealed predictor traits for indirect selection. Four cluster groupings with contrasting characteristics were also identified. Our study identified 20 stable landrace accessions within three *Dioscorea* spp. with above-average check performance for farmers' and consumers' preferred traits. These accessions could be advised to farmers, as well as considered in future yam improvement programs in DR Congo. Further characterization of these landraces is required with high throughput molecular markers to ascertain their genetic uniqueness before incorporation into future breeding programs. This will provide more insight into the challenge of linguistic polymorphism and the genetic diversity of these species for effective use as source of genetic reservoirs for yam improvement in DR Congo.

Supplementary Materials: The following are available online at https://www.mdpi.com/article/ 10.3390/agriculture12050599/s1, Table S1: List of the landrace accessions with better performance over checks mean for most preferred farmers and consumer traits and their stability.

Author Contributions: Conceptualization, I.I.A., D.O.O., P.A.A., J.G.A., and J.-C.L.M.; Methodology, I.I.A. and P.A.A.; Data analysis, I.I.A. and P.A.A.; Supervision, D.O.O., P.A.A., and J.G.A.; Writing—original draft, I.I.A. and P.A.A.; Writing—review and editing, I.I.A., P.A.A., D.O.O., J.G.A., J.-C.L.M., I.M.C., and J.L.K. All authors have read and agreed to the published version of the manuscript.

Funding: The African trans-regional cooperation, through the Mobilité Université en Afrique (MOUNAF) project funded by the European Union Commission within the framework of the Intra-Africa Academic Mobility Scheme, granted a Ph.D. scholarship to the first author to study at the University of Kisangani, Congo. This study is also partially supported by the BMGF, and publication fees will be covered by the BMGF.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Data used are available within this manuscript, including the supplementary files. Data can be obtained upon request from the corresponding author.

Acknowledgments: The authors acknowledge the provision of research funds to the first author by the MOUNAF project. The Directorate of Research and Finance office of the University of Kisangani is also acknowledged for managing the MOUNAF project. We appreciate the guidance the inspector of the Inspection Provinciale de l'Agriculture and the cooperation of all yam local farmers, local aids, and authorities that facilitated germplasm collection at the six territories used for the study. We also thank all other Ph.D. colleagues within the project and the department for their motivational support.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Agre, P.; Norman, P.E.; Asiedu, R.; Asfaw, A. Identification of Quantitative Trait Nucleotides and Candidate Genes for Tuber Yield and Mosaic Virus Tolerance in an Elite Population of White Guinea Yam (*Dioscorea rotundata*) Using Genome-Wide Association Scan. *BMC Plant Biol.* 2021, 21, 552. [CrossRef] [PubMed]
- 2. FAOSTAT. FAO Food and Agriculture Organization of the United Nations Statistics Database. 2020. Available online: http://www.fao.org/faostat/en/#data/QC (accessed on 2 February 2022).
- 3. Asiedu, R.; Sartie, A. Crops that feed the World 1. Yams. Food Secur. 2010, 2, 305–315. [CrossRef]
- 4. Cormier, F.; Lawac, F.; Maledon, E.; Gravillon, M.-C.; Nudol, E.; Mournet, P.; Vignes, H.; Chaïr, H.; Arnau, G. A reference high-density genetic map of greater yam (*Dioscorea alata* L.). *Theor. Appl. Genet.* **2019**, *132*, 1733–1744. [CrossRef] [PubMed]
- 5. Coursey, D.G. Yams: An account of the nature, origins, cultivation and utilisation of the useful members of the Dioscoreaceae. *Science* **1967**, *163*, 244.
- 6. Egesi, C.N.; Odu, B.O.; Ogunyemi, S.; Asiedu, R.; Hughes, J. Evaluation of water yam (*Dioscorea alata* L.) germplasm for reaction to yam anthracnose and virus diseases and their effect on yield. *J. Phytopathol.* **2007**, *155*, 536–543. [CrossRef]
- Darkwa, K.; Olasanmi, B.; Asiedu, R.; Asfaw, A. Review of empirical and emerging breeding methods and tools for yam (*Dioscorea* spp.) improvement: Status and prospects. *Plant Breed.* 2019, 139, 474–497. [CrossRef]
- 8. Onda, Y.; Mochida, K. Exploring Genetic Diversity in Plants Using High-Throughput Sequencing Techniques. *Curr. Genom.* 2016, 17, 358–367. [CrossRef]
- 9. Lebot, V. Tropical Root and Tuber Crops: Cassava, Sweet Potato, Yams, Aroids; CABI: Wallingford, UK, 2008.
- 10. Adejumobi, I.I.; Agre, P.A.; Onautshu, D.O.; Adheka, J.G. Diversity, trait preferences, management and utilization of yams landraces (*Dioscorea species*): An orphan crop in DR Congo. *Sci. Rep.* **2022**, *12*, 2252. [CrossRef]
- Bukatuka, F.C.; Ngombe, K.N.; Mutwale, K.P.; Moni, B.M.; Makengo, K.G.; Pambu, L.A.; Bongo, N.G.; Mbombo, M.P.; Musuyu, M.D.; Maloueki, U.; et al. Bioactivity and Nutritional Values of Some *Dioscorea species* Traditionally Used as Medicinal Foods in Bandundu, DR Congo. *Eur. J. Med. Plants* 2016, 14, e25124. [CrossRef]
- Jeancy, N.L.; Paul, M.; Alasca, E.L.; Yves-Dady, B. Yam production on the sandy soil of Bateke Plateau (DR Congo). J. Appl. Biosci. 2021, 17, 16886–16896. [CrossRef]
- 13. Institute National de la Statistiqu (INS). *INS Annuaire Statistique 2014;* Institute National de la Statistiqu: Montrouge, France; Ministère du Plan et Révolution de la Modernité: Toulouse, France, 2015.
- 14. Jehani, M.D.; Patel, P.R.; Chaudhary, A.K. Evaluation of bioagents against Colletotrichum capsici caused anthracnose disease of yam (*Dioscorea alata* L.). *J. Pharmacogn. Phytochem.* **2019**, *8*, 4788–4790.
- Gatarira, C.; Agre, P.; Matsumoto, R.; Edemodu, A.; Adetimirin, V.; Bhattacharjee, R.; Asiedu, R.; Asfaw, A. Genome-Wide Association Analysis for Tuber Dry Matter and Oxidative Browning in Water Yam (*Dioscorea alata* L.). *Plants* 2020, *9*, 969. [CrossRef] [PubMed]
- Bhattacharjee, R.; Nwadili, C.O.; Saski, C.A.; Paterne, A.; Scheffler, B.E.; Augusto, J.; Lopez-Montes, A.; Onyeka, J.T.; Kumar, P.L.; Bandyopadhyay, R. An EST-SSR based genetic linkage map and identification of QTLs for anthracnose disease resistance in water yam (*Dioscorea alata* L.). *PLoS ONE* 2018, 13, e0197717. [CrossRef] [PubMed]
- Adjumati, G.T.; Pembele, A.I.; Ocan, D. Use of charcoal (biochar) to enhance tropical soil fertility: A case of Masako in Democratic Republic of Congo. J. Soil Sci. Environ. Manag. 2020, 11, 17–29. [CrossRef]
- de Mendiburu, F. Agricolae: Statistical Procedures for Agricultural Research. R Package Version 1.2-0. 2014. Available online: http://CRAN.R-project.org/package=agricolae (accessed on 8 January 2022).
- Campbell, C.L.; Madden, L.V. Book Review: Introduction to Plant Disease Epidemiology; John Wiley and Sons: New York, NY, USA, 1990; Volume 19.
- 20. Asfaw, A. Standard Operating Protocol for Yam Variety Performance Evaluation Trial; International Institute of Tropical Agriculture (IITA): Ibadan, Nigeria, 2016.
- Kuznetsova, A.; Brockhoff, P.B.; Christensen, R.H.B. lmerTest Package: Tests in Linear Mixed Effects Models. J. Stat. Softw. 2017, 82, 1–26. [CrossRef]
- 22. Kresovich, S. Quantitative genetics in maize breeding. Field Crops Res. 1990, 23, 78–79. [CrossRef]
- Johnson, H.W.; Robinson, H.F.; Comstock, R.E. Genotypic and Phenotypic Correlations in Soybeans and Their Implications in Selection 1. Agron. J. 1955, 47, 477–483. [CrossRef]

- 24. Deshmukh, S.; Basu, M.; Reddy, P. Genetic variability, character association and path coefficients of quantitative traits in Virginia bunch varieties of groundnut. *Indian J. Agric. Sci.* **1986**, *56*, 816–821.
- Schloerke, B.; Cook, D.; Larmarange, J.; Briatte, F.; Marbach, M.; Thoen, E.; Elberg, A.; Crowley, J. Ggally: Extension to ggplot2, R Package Version 0.5.0.; R Foundation for Statistical Computing: Vienna, Austria, 2020; Available online: https://cran.r-project.org/ web/packages/GGally/ (accessed on 10 January 2022).
- 26. Team, R.C. A Language and Environment for Statistical Computing; R Foundation for Statistical Computing: Vienna, Austria, 2017.
- 27. Galili, T. dendextend: An R package for visualizing, adjusting and comparing trees of hierarchical clustering. *Bioinformatics* **2015**, *31*, 3718–3720. [CrossRef]
- 28. Gu, Z. Package 'circlize', R Packag. 2015.
- Charrad, M.; Ghazzali, N.; Boiteau, V.; Maintainer, A.N. Determining the Best Number of Clusters in a Data Set. Cran. 2015. Available online: http://cran.rediris.es/web/packages/NbClust/NbClust.pdf (accessed on 10 January 2022).
- 30. Rosseel, Y. Lavaan: An R package for structural equation modeling. J. Stat. Softw. 2012, 48, 1–36. [CrossRef]
- 31. Jacobucci, R.; Grimm, K.; McArdle, J. Package 'semPlot'. *Struct. Equ. Model.* 2016, 23, 4.
- Dieng, I.; Rodriguez, F.; Pacheco, A.; Alvarado, G.; Burgueno, J. ViTSel, Version 1.0; Africa Rice Centre (AfricaRice): Ibadan, Nigeria; Centro Internacional de Mejoramiento de Maiz y Trigo (CIMMYT): Texcoco, Mexico; International Institute of Tropical Agriculture (IITA): Ibadan, Nigeria, 2020.
- Garedew, B.; Haile, B.; Ayiza, A. Distribution, Diversity and Potential Production of Yams (*Dioscorea* spp.) in Sheko District, Southwest Ethiopia. Am. J. Life Sci. 2017, 5, 86. [CrossRef]
- 34. Villar, A.E.V.; Sarmiento-Zea, N.; León-Pulido, J.; Rojas-Pérez, L.C. Bioethanol Production from Yam (*Dioscorea rotundata*) Using Simultaneous Saccharification and Fermentation (SSF). *TecnoLógicas* 2021, 24, e1724. [CrossRef]
- Gabriel, A.; Agre, P.; Edemodu, A.; Okon, A.E.; Bhattacharjee, R.; de Koyer, D. Genetic diversity of white yam (*Dioscorea rotundata* poir) accessions maintained in tissue culture and cultivated in field using simple sequence repeat (ssr) markers and morphological characterization. *J. Agric. Crops* 2021, 7, 60–68. [CrossRef]
- 36. Kundu, B.B.; Vanni, K.; Farheen, A.; Jha, P.; Pandey, D.K.; Kumar, V. *Dioscorea bulbifera* L. (Dioscoreaceae): A review of its ethnobotany, pharmacology and conservation needs. *South Afr. J. Bot.* **2021**, *140*, 365–374. [CrossRef]
- 37. Ikiriza, H.; Ogwang, P.E.; Peter, E.L.; Hedmon, O.; Tolo, C.U.; Abubaker, M.; Abdalla, A.A.M. *Dioscorea bulbifera*, a highly threatened African medicinal plant, a review. *Cogent Biol.* **2019**, *5*, 1631561. [CrossRef]
- Scarcelli, N.; Cubry, P.; Akakpo, R.; Thuillet, A.-C.; Obidiegwu, J.; Baco, M.N.; Otoo, E.; Sonké, B.; Dansi, A.; Djedatin, G.; et al. Yam genomics supports West Africa as a major cradle of crop domestication. *Sci. Adv.* 2019, *5*, eaaw1947. [CrossRef]
- 39. Adewumi, A.S.; Agre, P.A.; Asare, P.A.; Adu, M.O.; Taah, K.J.; Mondo, J.M.; Akaba, S. Exploring the Bush yam (*Dioscorea praehensilis* Benth) as a Source of Agronomic and Quality Trait Genes in White Guinea. *Agronomy* **2022**, *12*, 55. [CrossRef]
- Egesi, C.N.; Asiedu, R.; Ude, G.; Ogunyemi, S.; Egunjobi, J.K. AFLP marker diversity in water yam (*Dioscorea alata* L.). *Plant Genet. Resour.* 2006, 4, 181–187. [CrossRef]
- 41. Egesi, C.N.; Onyeka, T.J.; Asiedu, R. Severity of anthracnose and virus diseases of water yam (*Dioscorea alata* L.) in Nigeria I: Effects of yam genotype and date of planting. *Crop Prot.* **2007**, *26*, 1259–1265. [CrossRef]
- 42. Padhan, B.; Panda, D. Potential of Neglected and Underutilized Yams (Dioscorea spp.) for Improving Nutritional Security and Health Benefits. *Photosynthetica* **2018**, *56*, 496. [CrossRef] [PubMed]
- Agre, P.; Asibe, F.; Darkwa, K.; Edemodu, A.; Bauchet, G.; Asiedu, R.; Adebola, P.; Asfaw, A. Phenotypic and molecular assessment of genetic structure and diversity in a panel of winged yam (*Dioscorea alata*) clones and cultivars. *Sci. Rep.* 2019, *9*, 18221. [CrossRef] [PubMed]
- 44. Agre, P.A.; Dassou, A.G.; Loko, L.E.Y.; Idossou, R.; Dadonougbo, E.; Gbaguidi, A.; Mondo, J.M.; Muyideen, Y.; Adebola, P.O.; Asiedu, R.; et al. Diversity of white Guinea yam (*Dioscorea rotundata* Poir.) cultivars from Benin as revealed by agro-morphological traits and SNP markers. *Plant Genet. Resour.* **2021**, *19*, 437–446. [CrossRef]
- Siadjeu, C.; Toukam, G.M.S.; Bell, J.M.; Nkwate, S. Genetic diversity of sweet yam *Dioscorea dumetorum* (Kunth) Pax revealed by morphological traits in two agro-ecological zones of Cameroon. *Afr. J. Biotechnol.* 2015, 14, 781–793. [CrossRef]
- Onyilagha, J.C.; Lowe, J. Studies on the Relationship of *Dioscorea cayenensis* and *Dioscorea rotundata* Cultivars. *Euphytica* 1986, 35, 733–739. [CrossRef]
- Loko, Y.L.; Adjatin, A.; Dansi, A.; Vodouhè, R.; Sanni, A. Participatory evaluation of Guinea yam (*Dioscorea cayenensis* Lam.–*D. rotundata* Poir. complex) landraces from Benin and agro-morphological characterization of cultivars tolerant to drought, high soil moisture and chips storage insects. *Genet. Resour. Crop Evol.* 2015, *62*, 1181–1192. [CrossRef]