

Correction

Correction: Zhao et al. Behavioral Responses, Fumigant Toxicity, and Contact Toxicity of Eugenol and Cinnamaldehyde Against *Sitobion avenae* Under Controlled Conditions. *Agriculture* 2026, 16, 1538

Zilan Zhao ^{1,2}, Xiao Ran ² , Mingzhe Zhang ¹, Yunhui Zhang ², Xiangrui Li ², Jianchun Qin ^{1,*}  and Xun Zhu ^{2,*} 

¹ College of Plant Science, Jilin University, Changchun 130062, China; 19932276905@163.com (Z.Z.); mzzhang@jlu.edu.cn (M.Z.)

² State Key Laboratory for Biology of Plant Diseases and Insect Pests, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing 100193, China; ranxiao99@126.com (X.R.); yhzhang@ippcaas.cn (Y.Z.); xrli@ippcaas.cn (X.L.)

* Correspondence: qinjc@jlu.edu.cn (J.Q.); zhuxun@caas.cn (X.Z.)

Missing Funding

In the original publication [1], the funders the National Key Research and Development Program of China (2024YFD1400400), the Earmarked Fund for China Agriculture Research System (CARS-03), and the National Natural Science Foundation of China (No. 32272604) were not included. The authors state that the scientific conclusions are unaffected. This correction was approved by the Academic Editor. The original publication has also been updated.

Table Legend

In the original publication, there was a mistake in the legend for Table 1. “Cinnamaldehyde” should be revised to “Repellent Effect”. The correct Table 1 legend appears below. The authors state that the scientific conclusions are unaffected. This correction was approved by the Academic Editor. The original publication has also been updated.

Table 1. The repellent effects of cinnamaldehyde on *S. avenae* (mean $\pm$ SD).

Concentration (μ L/mL)	Nc	Nt	Repellent Effect	95% CI
0.375	40	50	$-10.00\% \pm 0.14$ c	$-36.41-14.19$
0.750	56	34	$24.44\% \pm 0.10$ b	$14.88-34.01$
1.500	79	11	$75.56\% \pm 0.10$ a	$50.26-100.85$
3.000	83	7	$84.44\% \pm 0.04$ a	$74.88-94.01$
6.000	86	4	$91.11\% \pm 0.04$ a	$81.55-100.01$

Note: Nc = total number of insects in the negative control (solvent only) and Nt = total number of insects in the treatment arm, both pooled from three independent replicates. Levene’s test confirmed homogeneity of variances across all concentration groups ($F_{(4,10)} = 1.862$, $p = 0.194$). One-way ANOVA revealed a highly significant concentration effect ($F_{(4,10)} = 117.265$, $p < 0.001$). Different lowercase letters denote significant differences among concentrations (Tukey’s HSD, $p < 0.05$).



Received: 17 August 2026

Accepted: 18 August 2026

Published: 28 August 2026

Copyright: © 2026 by the authors.

Licensee MDPI, Basel, Switzerland.

This article is an open access article distributed under the terms and

conditions of the [Creative Commons](https://creativecommons.org/licenses/by/4.0/)

Attribution (CC BY) license.

Reference

1. Zhao, Z.; Ran, X.; Zhang, M.; Zhang, Y.; Li, X.; Qin, J.; Zhu, X. Behavioral Responses, Fumigant Toxicity, and Contact Toxicity of Eugenol and Cinnamaldehyde Against *Sitobion avenae* Under Controlled Conditions. *Agriculture* **2026**, *16*, 1538. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.