





Anti-Metastatic Effects of Plant Sap-Derived Extracellular Vesicles in a 3D Microfluidic Cancer Metastasis Model

Kimin Kim 1 , Jik-Han Jung 2 , Hye Ju Yoo 1 , Jae-Kyung Hyun 3 , Ji-Ho Park 2 , Dokyun Na 4 and Ju Hun Yeon 1,*

- ¹ Department of Integrative Biosciences, University of Brain Education, Cheonan 31228, Korea; abc-632@hanmail.net (K.K.); hyeju_yoo@naver.com (H.J.Y.)
- ² Department of Bio and Brain engineering, Korea Advanced Institute of Science and Technology, Daejeon 34141, Korea; jjhan@kaist.ac.kr (J.-H.J.); jihopark@kaist.ac.kr (J.-H.P.)
- ³ Electron Microscopy Research Center, Korea Basic Science Institute, Cheongju 28119, Korea; jaekyung.hyun@oist.jp
- ⁴ School of Integrative Engineering, Chung-Ang University, Seoul 06911, Korea; blisszen@lile.cau.ac.kr
- * Correspondence: jhyeon@ube.ac.kr; Tel.: +82-41-529-2621; Fax: +82-41-529-2674




Figure S1. Uptake of DM-EVs into HUVECs. (**a**) Representative fluorescence microscopic images of cellular uptake of fluorescently labeled DM-EVs. (**b**) A comparison of fluorescence intensity with different time of 0, 1, 3, 6, 9, 12 h, respectively. Scale bar: 50 μm.



© 2020 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).