

Platinum and Iridium Oxide Co-modified TiO₂ Nanotubes Array Based Photoelectrochemical Sensors for Glutathione

Jing Tian¹, Peng Zhao¹, Shasha Zhang¹, Guona Huo¹, Zhaochen Suo¹, Zhao Yue^{2,*}, Shoumin Zhang¹, Weiping Huang¹ and Baolin Zhu^{1,3,*}

¹ College of Chemistry, The Key Laboratory of Advanced Energy Materials Chemistry (Ministry of Education), Tianjin Key Lab of Metal and Molecule-based Material Chemistry, Nankai University, Tianjin 300071, China; tianjingnk@nankai.edu.cn (J.T.); zhaopeng@nankai.edu.cn (P.Z.); 2120170933@mail.nankai.edu.cn (S.Z.); 15230153709@163.com (G.H.); zc.suo@mail.nankai.edu.cn (Z.S.); zhangsm@nankai.edu.cn (S.Z.); hwp914@nankai.edu.cn (W.H.)

² Department of Microelectronics, Nankai University, Tianjin 300350, China

³ National Demonstration Center for Experimental Chemistry Education (Nankai University), Tianjin 300071, China

* Correspondence: yuezhaoy@nankai.edu.cn (Z.Y.); zhubaolin@nankai.edu.cn (B.Z.)




Figure S1. The curve of Pt-IrO₂/TiO₂NTs/Ti (after 30 days) for the detection of different concentrations for GSH.