

A Review: Antimicrobial Therapy for Human Pythiosis

Sadeep Medhasi ¹, Ariya Chindamporn ² and Navaporn Worasilchai ^{3,*}

¹ Department of Transfusion Medicine and Clinical Microbiology, Faculty of Allied Health Sciences, Chulalongkorn University, Bangkok 10330, Thailand; sadeep.m@chula.ac.th

² Department of Microbiology, Faculty of Medicine, Chulalongkorn University, Bangkok 10330, Thailand; ariya.c@chula.ac.th

³ Department of Transfusion Medicine and Clinical Microbiology, Faculty of Allied Health Sciences, and Immunomodulation of Natural Products Research Group, Chulalongkorn University, Bangkok 10330, Thailand

* Correspondence: navaporn.w@chula.ac.th; Tel.: +66-2218-1065

Abstract: Human pythiosis is associated with poor prognosis with significant mortality caused by *Pythium insidiosum*. Antimicrobials' in vitro and in vivo results against *P. insidiosum* are inconsistent. Although antimicrobials are clinically useful, they are not likely to achieve therapeutic success alone without surgery and immunotherapy. New therapeutic options are therefore needed. This non-exhaustive review discusses the rationale antimicrobial therapy, minimum inhibitory concentrations, and efficacy of antibacterial and antifungal agents against *P. insidiosum*. This review further provides insight into the immunomodulating effects of antimicrobials that can enhance the immune response to infections. Current data support using antimicrobial combination therapy for the pharmacotherapeutic management of human pythiosis. Also, the success or failure of antimicrobial treatment in human pythiosis might depend on the immunomodulatory effects of drugs. The repurposing of existing drugs is a safe strategy for anti-*P. insidiosum* drug discovery. To improve patient outcomes in pythiosis, we suggest further research and a deeper understanding of *P. insidiosum* virulence factors, host immune response, and host immune system modification by antimicrobials.

Keywords: human pythiosis; antibacterial; antifungal; immunomodulatory; drug repurposing

Citation: Medhasi, S.; Chindamporn, A.; Worasilchai, N. A Review: Antimicrobial Therapy for Human Pythiosis. *Antibiotics* **2022**, *11*, 450. <https://doi.org/10.3390/antibiotics11040450>

Academic Editors: George Notas and Federico Pea

Received: 14 January 2022

Accepted: 22 March 2022

Published: 26 March 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Human pythiosis is an infectious disease with high morbidity and mortality [1]. *Pythium insidiosum*, a fungus-like aquatic oomycete microorganism, is a causative agent of pythiosis. The motile flagellate zoospore plays a significant role in initiating an infection. The zoospores of *P. insidiosum* adhere to the skin cut or wound sites and encyst on the surface of the injured tissue(s). The encysted spore develops a germination tube (hypha) that uses chemotaxis to find the host and infiltrate human blood vessels [1,2]. Pythiosis risk is higher in tropical and subtropical regions, including Southeast Asia, eastern coastal Australia, and South America [3].

Human pythiosis is associated with a poor prognosis due to the difficulties in diagnosing the infection and the lack of effective therapeutic agents against this disease [4]. The clinical features of human pythiosis are classified into four forms: (i) vascular pythiosis characterized by arteritis, thrombosis, gangrene, aneurysm, or limb claudication; (ii) ocular pythiosis characterized by corneal ulcers, decreased visual acuity, conjunctival redness, eyelid swelling, or multiple, linear, tentacle-like infiltrates and dot-like or pinhead-shaped infiltrates in the surrounding cornea; (iii) cutaneous and subcutaneous pythiosis characterized by a granulomatous and ulcerating lesion in the face or limbs, cellulitis, soft tissue abscess, or lymphadenopathy; and (iv) disseminated pythiosis characterized by the

infection of internal organs [1,4,5]. The risk factors for vascular pythiosis include thalassemia, hemoglobinopathy, paroxysmal nocturnal hemoglobinuria, aplastic anemia, and leukemia because *P. insidiosum* has a higher affinity for iron [6,7].

When an infection is diagnosed as *P. insidiosum*, the therapeutic options include surgery, pharmacotherapy, and immunotherapy (Figure 1) [8]. Surgical intervention is the mainstay treatment for managing human pythiosis, but such treatment substantially increases the financial burden on patients, postsurgical complications, and uncontrolled infection [9]. Immunotherapy is a promising approach for human pythiosis treatment where antigens of *P. insidiosum* from *in vitro* cultures are injected into the patient [10,11]. The mechanism behind *P. insidiosum* antigen (PIA) immunotherapy in human pythiosis includes a switch from the host's T helper-2 (Th2) to T helper-1 (Th1) mediated immune response in the host; the Th1 response producing higher levels of interferon- γ (IFN- γ) and interleukin 2 (IL-2) [10,12,13]. Even though a good prognosis in PIA-treated patients can be implied by Th2 to Th1 switching, the efficacy of *P. insidiosum* antigen is inconclusive when used as immunotherapy in human pythiosis [7,12,14]. *In vitro* studies have demonstrated the anti-*P. insidiosum* effect of antifungals even though the *P. insidiosum* lacks the antifungal drug-target: ergosterol biosynthetic pathway [15]. However, a significant concern with antifungals is the contradictory results in susceptibility to *P. insidiosum* in *in vitro* and clinical use [15,16].



Figure 1. Photograph of human pythiosis. A 46-year-old Thai male with thalassemia was diagnosed with vascular pythiosis. CTA showed the occlusion of the right aorta, and ELISA showed the positive IgG against *P. insidiosum* (with permission). Abbreviations: CTA, computed tomography angiography; ELISA, enzyme-linked immunosorbent assay.

This review focuses on evidence supporting and disputing the effectiveness of antimicrobials to expand the pharmacotherapeutic role of antimicrobials in the management of human pythiosis. We do not explicitly discuss the biology of *P. insidiosum*, the pathogenesis of the *P. insidiosum* infection in humans, or the management of human pythiosis with immunotherapy and surgical intervention. Finally, we conclude with general remarks on future strategic options for managing human pythiosis.

2. Principles of Antimicrobial Therapy

Antimicrobial therapy should achieve a clinical response by eliminating the invading microorganism(s) while minimizing cost, adverse effects, and antimicrobial resistance [17,18]. When selecting appropriate antimicrobial therapy, both pharmacokinetic and pharmacodynamic properties of the drug(s) must be considered to ensure that effective agents are administered in sufficient doses for therapeutic success [19]. For species such as *P. insidiosum*, identifying potential targets for antimicrobials is necessary for managing pythiosis. The microbial cell wall is a critical target for antimicrobials, and the cell wall of *P. insidiosum* is primarily composed of β -glucan and cellulose [20]. However, the cell wall of *P. insidiosum* lowers the penetration of drug molecules and prevents drug access to targets inside the cell wall [21]. The gene expression of cytochrome oxidase 2 (COX2) in Thai *P. insidiosum* strains was 2.5-fold higher at 37 °C compared to the expression at 27 °C [22]. In addition, the elicitor protein, *ELI025*, was highly up-regulated in *P. insidiosum* hyphae at 37 °C compared to hyphae grown at 28 °C and facilitated the evasion of the host antibody response [23]. COX2 and *ELI025* can be candidate targets for controlling *P. insidiosum* infection.

Several antifungal and antibacterial drugs have been examined for their susceptibility profile against *P. insidiosum* in an *in vitro* study (Figure 2). They have been tried to manage human pythiosis but have been successful only in a few cases [4]. *P. insidiosum* keratitis was successfully managed in a 20-year-old Japanese man following triple antibiotic therapy (minocycline ointment four times a day, chloramphenicol eye drops hourly, and linezolid 1200 mg orally twice a day) [24]. Recently, a *P. insidiosum* keratitis patient was successfully managed with topical 0.2% linezolid and topical 1% azithromycin, administered hourly [25]. Antimicrobial susceptibility testing (AST) is a procedure to determine the concentration of an antimicrobial that inhibits microbial growth *in vitro* by establishing minimum inhibitory concentration (MIC), which is the lowest concentration of an antimicrobial that inhibits visible growth of a microorganism [26,27]. Table 1 summarizes the methods used to determine the MIC of antimicrobial drugs against *P. insidiosum* discussed in our review.




Figure 2. Mode of MIC value of each antibacterial/antifungal class against *P. insidiosum* isolates reviewed in previous publications. Mode of MIC value of *P. insidiosum* isolates against antimicrobial

drugs in class different antimicrobial classes: tetracyclines (4 µg/ml) [28,29], macrolides (6 µg/ml) [29], oxazolidinones (8 µg/ml) [29], lincosamides (4 µg/ml) [30], streptogramins (2 µg/ml) [30], phenicols (16 µg/ml) [30], aminoglycosides (64 µg/ml) [31], nitrofurantoin (no data) [30], mupirocin (4 µg/ml) [29], polyenes (64 µg/ml) [29], allylamines and azoles (4 µg/ml) [9], and echinocandins (4 µg/ml) [32].

Table 1. Summary of methods for determining MICs of antimicrobial drugs against *P. insidiosum*.

Antimicrobial Class	Drug	MIC Determination Method(s)	Reference(s)
Tetracyclines	Tetracycline	Broth microdilution	[28]
	Tigecycline	Broth microdilution, disk diffusion, and Etest	[28,29,31]
	Minocycline	Broth microdilution, disk diffusion, and Etest	[28,29]
Macrolides	Azithromycin	Broth microdilution, disk diffusion, and Etest	[28,29]
	Clarithromycin	Broth microdilution, disk diffusion, and Etest	[28,29]
Oxazolidinones	Linezolid	Broth microdilution, disk diffusion, and Etest	[29]
Lincosamides	Clindamycin	Broth dilution	[30]
Streptogramins	Quinupristin and dalfopristin	Broth dilution	[30]
Phenicols	Chloramphenicol	Broth dilution	[30]
Aminoglycosides	Gentamicin	Broth microdilution	[31]
	Neomycin	Broth microdilution	[31]
	Paromomycin	Broth microdilution	[31]
	Streptomycin	Broth microdilution	[31]
	Nitrofurantoin	Broth dilution	[30]
	Mupirocin	Broth microdilution, disk diffusion, and Etest	[29]
Polyenes	Amphotericin B	Etest	[29]
Allylamines	Terbinafine	Broth dilution and radial growth	[9]
Azoles	Miconazole	Broth microdilution	[9]
	Ketoconazole	Broth microdilution	[9]
	Fluconazole	Broth microdilution and agar diffusion	[9]
	Itraconazole	Broth microdilution, radial growth, and agar diffusion	[9]
	Posaconazole	Broth microdilution and agar diffusion	[9]
	Voriconazole	Broth microdilution, radial growth, and agar diffusion	[9]
	Caspofungin	Broth dilution	[32]
Echinocandins	Anidulafungin	Broth dilution	[32]

Abbreviations: MIC, minimal inhibitory concentration.

3. Why Do Antimicrobial Treatments Fail?

Factors contributing to the antimicrobial treatment failure include antimicrobial agent's pharmacokinetic and pharmacodynamic issues related to the antimicrobial agent, lack of pathogen control, development of infection complications, drug-resistant pathogens, conflicting AST results, disparities between *in vitro* and *in vivo* efficacy, host immune response, and wrong choice of antimicrobial drug (Figure 3) [33,34]. Pharmacokinetics variability can be defined as differences in plasma antimicrobial exposure, impacting treatment success [35]. Antimicrobials, like beta-lactams and aminoglycosides, achieve suboptimal plasma concentrations in critically ill patients due to increased volume of distribution and increased renal and hepatic clearance [36,37]. As another example, linezolid's pharmacokinetic variability results in adverse effects and ineffective therapy because of the narrow therapeutic window of linezolid [38].




Figure 3. Antimicrobial treatment in the management of *P. insidiosum* infection. Antibacterial and antifungal drugs exhibit immunomodulation activity and can improve treatment strategies for human pythiosis. Several mechanisms contribute to antimicrobial failure during the treatment of diseases.

P. insidiosum produces six enzymes (ERG3, ERG5, ERG11, ERG20, ERG24, and ERG26) included in the sterol biosynthetic pathways [16]. However, more than 40 enzymes are involved in the sterol biosynthetic pathways; thus, drugs targeting sterol pathways exhibit limited efficacy against *P. insidiosum*. These drugs cannot be exploited for rationalized and successful management of pythiosis [39]. Different strategies could be considered to prevent the antimicrobial treatment failure in pythiosis, namely: delivering adequate concentration of antimicrobial drug at the site of infection [40], increased periods of exposure of *P. insidiosum* to the antimicrobial drug [41], redesigning drug to penetrate the outer membrane of *P. insidiosum* and avoid being pumped out of the membrane [42], and modulate host immunity [43].

4. Immune Response and Antimicrobial Therapy

The innate immune system protects the host from various toxins and infectious agents, including bacteria, fungi, viruses, and parasites via phagocytosis and intracellular killing, recruitment of other inflammatory cells, and presentation of antigens [44]. The innate immune system is highly complex and comprises physical and anatomical barriers, effector cells, antimicrobial peptides, soluble mediators, and cell receptors [45]. However, pathogens can breach the early innate immune mechanisms. In these circumstances, a strategy to modify the function of immune cells can lead to the elimination of the pathogenic intruder [46]. Interestingly, host immunity is often overlooked in the process of pathogen clearance. A favorable innate immune response can considerably reduce the need for more prolonged antimicrobial therapy in infections [47].

Once *P. insidiosum* enters and adheres to the host tissues, the soluble exoantigens from *P. insidiosum* trigger the Th2 response and lock the host immune system into a Th2 subset. Further, *P. insidiosum* protects itself from the host immune system by concealing inside the eosinophilic material formed by the eosinophil degranulation, which helps protect the *P. insidiosum* from being fully presented to the host's immune system [10]. Toll-like receptors (TLRs) play a central role in the innate immune system by recognizing pathogen-associated molecular patterns and triggering downstream signaling pathways that activate the innate immune response [48]. Wongprompitak et al. demonstrated that both zoospores and hyphae of *P. insidiosum* induced a TLR2-mediated innate immune response with a subsequent increase in the levels of the pro-inflammatory cytokines IL-6 and IL-8 [49].

To combat the pathogen and prevent its spread, it is rational to administer antimicrobial drugs that interact with the host's innate immune system to provide profound indirect effects and enhance pathogen clearance. Antimicrobial drugs have been shown to modify the immune responses to infection, guiding improved treatment strategies in human pythiosis (Table 2 and Figure 3).

Table 2. Immunomodulatory effects of antimicrobials.

Antimicrobial Class	Drug	Immunopharmacological Effect	Reference(s)
Tetracyclines	Tigecycline, minocycline	Potentiate the innate immune response and augment resolution of inflammation	[50]
Macrolides	Azithromycin	Reduce the production of IL-12, resulting in enhanced Th2 response	[51]
Oxazolidinones	Linezolid	Suppress synthesis of proinflammatory cytokines, such as interleukin-1 β (IL-1 β), IL-6, IL-8, interferon- γ (IFN- γ), and tumor necrosis factor- α (TNF- α)	[52–54]
Lincosamides	Clindamycin	Suppress the release of inflammatory cytokines such as TNF- α and IL-1 β and enhance the phagocytosis of microorganisms by host cells	[55,56]
Streptogramins	Quinupristin-dalfopristin	Decrease the concentration of pro-inflammatory cell wall components (lipoteichoic acid and teichoic acid) and the activity of TNF	[57]
Phenicol	Chloramphenicol	Elevate the anti-inflammatory IL-10 levels	[58]
Polyenes	Amphotericin B	Activate the host's innate immunity and augment the IL-1 β -induced inducible nitric-oxide synthase (iNOS) expression and the production of nitric oxide (NO)	[59]
Allylamines	Terbinafine	Stimulate proinflammatory cytokines	[60]
Azoles	Fluconazole, voriconazole	Enhance microbicidal activity of monocytes, macrophages, and neutrophils	[61,62]

5. Antibacterial Drugs against *P. insidiosum*

Previous *in vitro* screening of antibacterial drugs has identified tetracycline, minocycline, tigecycline, azithromycin, clarithromycin, erythromycin, gentamicin, streptomycin, paromomycin, neomycin, linezolid, nitrofurantoin, quinupristin-dalfopristin, chloramphenicol, clindamycin, and mupirocin, which demonstrated inhibitory activity against *P. insidiosum* [29–31,63–65]. Among the arsenal of antibiotics, the best-studied antibiotics in human pythiosis are tetracyclines, macrolides, and oxazolidinones. This section discusses different classes of antibacterial drugs to manage human pythiosis.

5.1. Tetracyclines

Tetracycline antibiotics such as tetracycline, tigecycline, and minocycline inhibit bacterial protein synthesis by binding with the bacterial 30S ribosomal subunit [66]. Tetracyclines can inhibit mammalian collagenase activity and assist wound healing [67]. Further, tetracyclines potentiate the innate immune response and augment the resolution of inflammation [50].

Based on the *in vivo* studies in rabbits, minocycline in combination with immunotherapy may be an effective therapeutic medical treatment of pythiosis to heal injuries [68]. Worasilchai et al. evaluated the *in vitro* susceptibility of human, environmental, and animal *P. insidiosum* isolates to eight antibiotic classes and demonstrated that tetracyclines and macrolides inhibited the *in vitro* growth of *P. insidiosum* isolates at concentrations 10 to 100 times lower than those observed for previously studied antifungal drugs [28]. Also, the combination of tetracyclines and macrolides resulted in a synergistic effect that reduced MICs against *P. insidiosum* isolates. Loreto et al. also reported a similar *in vitro* susceptibility of *P. insidiosum* isolates to tetracyclines and their superior potency compared to amphotericin B, echinocandins, and triazole antifungals [29].

5.2. Macrolides

Macrolides are the group of antibiotics that inhibit bacterial protein synthesis by binding with the bacterial 50S ribosomal subunit. Common macrolides include erythromycin, clarithromycin, and azithromycin [69]. Among the macrolides, azithromycin, in particular, is highly accumulated in phagocytes and is targeted to the sites of infection [70]. Azithromycin reduces the production of IL-12, resulting in enhanced Th2 response [51]. Th2 cells are involved in wound healing and tissue repair [71,72]. The immunomodulatory activities of macrolides are evident with both pro-inflammatory and anti-inflammatory effects. For example, erythromycin can suppress pro-inflammatory cytokine production, such as IL-6, IL-8, and tumor necrosis factor- α (TNF- α) [73].

Jesus et al. investigated the antimicrobial activity of azithromycin alone and in combination with minocycline against *P. insidiosum* in a rabbit model [74]. The results revealed a strong *in vivo* activity of azithromycin (20 mg/kg/day twice daily) alone and combination with minocycline (10 mg/kg/day twice daily) against subcutaneous lesions. In an *in vitro* susceptibility study, the MICs of azithromycin and clarithromycin were less than 4 μ g/ml for *P. insidiosum* isolates [29].

5.3. Oxazolidinones

Oxazolidinones such as linezolid inhibit bacterial protein synthesis by binding with the 50S subunit of the ribosome [75]. The suppression of the synthesis of pro-inflammatory cytokines, such as interleukin-1 β (IL-1 β), IL-6, IL-8, IFN- γ , and TNF- α by linezolid has highlighted an exciting role of linezolid in immunomodulatory effects [52–54]. Linezolid may significantly reduce the inflammatory damage induced by the excessive release of pro-inflammatory cytokines during critical infections [76].

In a rabbit model of *P. insidiosum* keratitis, topical linezolid demonstrated superior efficacy and safety compared to azithromycin and tigecycline after prolonged treatment for more than 3–4 weeks [77].

5.4. Lincosamides, Streptogramins, and Phenicol

Lincosamides, streptogramins, and phenicols inhibit bacterial protein synthesis by interacting with the 50S subunit of bacterial ribosomes [78]. Among lincosamides, clindamycin possesses immunomodulatory activity by suppressing the release of inflammatory cytokines such as TNF- α and IL-1 β and enhancing the phagocytosis of microorganisms by host cells [55,56]. Quinupristin and dalbapristin, used in a fixed combination, belong to a class of streptogramins [78]. Quinupristin-dalbapristin decreased the concentration of pro-inflammatory cell wall components (lipoteichoic acid and teichoic acid) and TNF activity in cerebrospinal fluid compared to the ceftriaxone-treated rabbits [57]. A previous report showed that chloramphenicol, a member of the phenicols group, elevated the IL-10 levels, a potent anti-inflammatory cytokine [58].

Lincosamides, streptogramins, and phenicols have shown the ability to inhibit the growth of *P. insidiosum* isolates. The microdilution-based MIC ranges (with geometric means) of lincosamides, streptogramins, and phenicols against *P. insidiosum* were reported to be 2 to >4 $\mu\text{g/ml}$, 1 to >2 $\mu\text{g/ml}$, and 8 to >16 $\mu\text{g/ml}$, respectively [30].

5.5. Aminoglycosides

Aminoglycosides such as gentamicin, streptomycin, paromomycin, and neomycin bind to the bacterial ribosome and inhibit protein synthesis [79]. Streptomycin stimulated the *in vitro* growth of one of the Thai *P. insidiosum* isolates [80]. Aminoglycoside antibiotics inhibited the *in vitro* growth of *P. insidiosum*; however, they may not be clinically relevant due to the high MIC values [31]. Therefore, aminoglycosides for clinical use in managing human pythiosis are questionable.

5.6. Miscellaneous Antibacterial Drugs

Nitrofurantoin is used to treat urinary tract infections and works by attacking bacterial ribosomal proteins non-specifically, causing complete inhibition of protein synthesis [81]. *P. insidiosum* mycelial growth was inhibited with nitrofurantoin (MIC range of 64 to >64 $\mu\text{g/ml}$) in an *in vitro* susceptibility test [30].

Mupirocin inhibits bacterial protein and RNA synthesis by reversibly inhibiting isoleucyl-transfer RNA [82]. A study evaluating the *in vitro* susceptibility of Brazilian *P. insidiosum* strains showed that mupirocin could inhibit the growth of *P. insidiosum* isolates at MIC lower than 4 $\mu\text{g/ml}$ [29].

6. Antifungal Drugs against *P. insidiosum*

Studies have focused on several antifungal medications, such as polyenes, azoles, allylamines, and echinocandins, for the adjunctive therapy in managing human pythiosis [4]. Despite the evidence of anti-*P. insidiosum* effects, it has been highly challenging to achieve consistently effective antifungal treatment in human pythiosis.

6.1. Polyenes

Amphotericin B is a polyene antifungal that binds to ergosterol in the fungal cell membrane, which alters cell membrane permeability leading to the loss of intracellular components [83]. Two Australian cases with subcutaneous pythiosis responded well to amphotericin B treatment [84]. However, the evidence of the effectiveness of amphotericin B against other forms of human pythiosis and substantial activity against *P. insidiosum* is lacking [14,85–87].

Studies have shown amphotericin B's immunomodulatory properties, which activate the host's innate immunity [88]. Nitric oxide (NO) is an endogenous regulator of inflammation and an antibacterial agent, and it plays a crucial role in wound repair [89,90]. Amphotericin B can augment the IL-1 β -induced inducible nitric-oxide synthase (iNOS) expression and NO production [59]. In addition, amphotericin B is reported to induce oxidative stress and improve antifungal efficacy [91,92].

6.2. Allylamines and Azoles

The primary mode of action of allylamines, such as terbinafine, is the inhibition of the enzyme squalene monooxygenase. Therefore, these drugs inhibit the fungal synthesis of ergosterol [93]. Azoles, such as miconazole, ketoconazole, fluconazole, itraconazole, posaconazole, and voriconazole, exhibit antifungal activity by inhibiting the 14 α -lanosterol demethylase, a key enzyme in ergosterol biosynthesis, in fungi [94,95]. Studies have suggested that the enhanced microbiocidal activity of monocytes, macrophages, and neutrophils against intracellular *Candida albicans* is enhanced when combined with azoles [61,62]. However, terbinafine has been reported to stimulate pro-inflammatory cytokines [60].

Susaengrat et al. reported favorable responses to voriconazole and itraconazole in Thai vascular pythiosis patients [96]. Synergistic effects have been demonstrated for terbinafine and fluconazole against *P. insidiosum* isolates *in vitro* [97]. A synergistic combination of itraconazole and terbinafine was effective during the *in vitro* susceptibility testing of a *P. insidiosum* isolate from the 2-year-old patient with a deeply invasive facial infection [98]. The growth of *P. insidiosum* isolates was inhibited by terbinafine, and the efficacy of terbinafine increased against *P. insidiosum* isolates when combined with cetrimide, an antiseptic [63]. Pediatricians used a combination of itraconazole and terbinafine to manage a child with vascular pythiosis [99]. *In vitro* susceptibility testing of *P. insidiosum* showed MICs from 0.5 to 128 μ g/ml for terbinafine, 2 to 32 μ g/ml for miconazole, 4 to 64 μ g/ml for ketoconazole, 1 to >128 μ g/ml for itraconazole, 2 to >16 μ g/ml for voriconazole, greater than 1 to >32 μ g/ml for fluconazole, and >8 μ g/ml for posaconazole based on the strains of *P. insidiosum* [9,14].

6.3. Echinocandins

Echinocandins, such as caspofungin, anidulafungin, and micafungin, act by inhibiting beta-(1,3)-D-glucan synthase, an enzyme that is necessary for the synthesis of beta-(1,3)-D-glucan, which is an essential component of the fungal cell wall [99]. Studies have documented the immunomodulatory effects of echinocandins with increased fungal beta-(1,3)-D-glucan exposure and caspofungin-induced neutrophil-mediated fungal damage and anidulafungin- and micafungin-induced phagocyte-mediated fungal damage [100,101].

Synergistic anti-*P. insidiosum* effects were observed with caspofungin and terbinafine *in vitro* [97]. The MICs of caspofungin and anidulafungin against human *P. insidiosum* isolates ranged from 2 to 8 μ g/ml [32]. However, when used alone, echinocandins showed poor *in vitro* and *in vivo* activity against *P. insidiosum* [29,102]. Caspofungin demonstrated less fungistatic activity against *P. insidiosum* [103].

6.4. Miscellaneous Antifungal Drugs

Amorolfine, a morpholine derivative, inhibits fungal ergosterol biosynthesis and leads to changes in the membrane permeability, which in turn causes fungal growth inhibition and cell death [104]. Only recently, amorolfine hydrochloride exhibited *in vitro* inhibitory activity against *P. insidiosum* [15]. The MICs of amorolfine hydrochloride tested against *P. insidiosum* isolates were 16 to 64 mg/L. Further, amorolfine hydrochloride produced alterations in *P. insidiosum* hyphae, with changes in the surface of hyphae, intracellular organelles, the cell wall, and plasma membrane of *P. insidiosum*.

7. Repurposing Antimicrobials against *P. insidiosum*

Due to the limited success of pharmacological interventions against *P. insidiosum* in humans, identifying novel therapeutic strategies is required to treat *P. insidiosum* infection in humans. Drug repurposing is a process for identifying new therapeutic indications different from the scope of the initial pharmacological indication [105]. For example, antibiotics such as macrolides, tetracyclines, and fluoroquinolones have been used in the clinical

management of coronavirus disease 2019 (COVID-19) [106]. Using the drug repurposing strategy, existing FDA-approved antimicrobials can forgo early phases of drug development in managing human pythiosis [107]. Disulfiram irreversibly inhibits aldehyde dehydrogenase (ALDH1A1) and is an alcohol-deterrent medication that causes a severe adverse reaction when patients use alcohol. Disulfiram effectively treats individuals dependent on alcohol but highly motivated to discontinue alcohol use [108]. Krajaejun et al. evaluated disulfiram for its anti-*P. insidiosum* activity using agar- and broth-based methods and revealed that *P. insidiosum* strains were susceptible to disulfiram with MICs ranging from 8 to 32 mg/liter [109]. Further, disulfiram was found to bind and inactivate aldehyde dehydrogenase and urease of *P. insidiosum*.

Researchers utilize computational and experimental approaches to identify the promising candidates in the drug repurposing process [110]. The computational system uses various databases and computational tools, such as Gene Signature Database (GeneSigDB), Gene Set Enrichment Analysis (GSEA), The Pharmacogenetics and Pharmacogenomics Knowledge Base (PharmGKB), DrugBank, ChemBank, Genecard, Online Mendelian Inheritance in Man (OMIM), PubMed, e-Drug3D, DrugPredict, Promiscuous, Mantra2.0, Protein Data Bank (PDB), DRAR-CPI, repoDB, Repurpose DB, DeSigN, Cmap, and DPDR-CPI, etc. [105,111]. Computational techniques employed for drug repurposing include (i) profile-based drug repositioning, (ii) network-based drug repositioning, and (iii) data-based drug repositioning [112]. Experimental-based approaches validate the computer-generated hits for preclinical drug evaluation [111]. An experimental technique for drug repurposing involves protein target-based and cell/organism-based screens in *in vitro* and *in vivo* assays [113].

Using combination regimens of antibacterial plus antifungal or antibacterial plus antibacterial to achieve synergistic activity is one of the drug repurposing strategies against *P. insidiosum* [114]. Synergism between antibacterial and antifungal against *P. insidiosum* was observed for *in vitro* minocycline with amphotericin B, itraconazole, and micafungin and clarithromycin with micafungin [65]. Susaengrat et al. reported two cases of relapsed vascular pythiosis patients who were successfully clinically managed with a combination of antibacterial plus antifungal [96]. However, isolate-specific combinations for treatment must be implemented because of the varying effectiveness of any given drug combination for different isolates of *P. insidiosum* [115]. Studies have found the enhanced killing effects of multiple classes of antibiotics when combined with NO [116,117]. We expect that NO-containing antibiotics might improve the therapeutic outcomes in patients with pythiosis.

8. Conclusions and Future Perspectives

Evidence supports using the antimicrobials reviewed in our article as a new therapeutic option in treating human pythiosis. *In vitro* studies have demonstrated the tetracyclines, macrolides, oxazolidinones, lincosamides, streptogramins, phenicols, aminoglycosides, polyenes, allylamines, azoles, and echinocandins reviewed in our papers inhibit the growth of *P. insidiosum* and have the potential implications for further research on their use in the management of human pythiosis. However, prolonged use of antimicrobials and prolonged treatment with antimicrobials is not warranted due to the side effects and threat of antimicrobial resistance. A practical pharmacological intervention guideline for human pythiosis remains to be discovered and is necessary to assist practitioner and patient decisions, lower treatment costs, and optimize patient outcomes. Despite the disease affecting the most vulnerable populations with higher mortality rates, pythiosis is not included in the Sanford Guide, which provides evidence-based recommendations for treating infectious diseases [118].

In the future, human pythiosis could be managed with antimicrobials owing to their anti-inflammatory and immunomodulatory activities. Clinicians can optimize drug combinations based on the anti-*P. insidiosum* susceptibility testing for the management of pythiosis. Studies have shown the growth inhibitory effects of antimicrobials against *P. insidiosum*; nevertheless, studies regarding the mechanism of action of the antimicrobials

against *P. insidiosum* are vital for clinical approval. Researchers must consider the pharmacodynamics principle involved in selecting the antimicrobials to assess the anti-*P. insidiosum* activity.

Microbial virulence factors are molecules produced by microorganisms and may cause disease in the host (e.g., toxins, enzymes, exopolysaccharides, lipopolysaccharides, lipoproteins, etc.) [22]. The potential virulence factors of *P. insidiosum* include glucan 1,3-beta-glucosidase, heat shock protein (Hsp) 70, and enolase [23]. Keeratijarut et al. reported genetic, immunological, and biochemical characteristics of Exo-1,3- β -glucanase (Exo1) in *P. insidiosum* and found up-regulated *exo1* expression at 37 °C compared to 28 °C, thus suggesting a drug target against *P. insidiosum* [119]. A new therapeutic approach with anti-virulence therapy combined with antimicrobials might prevent the pathogenesis of *P. insidiosum* and limit host damage. Metabolites have been isolated from *Pseudomonas stutzeri* and *Klebsiella pneumoniae*, and these organisms have shown anti-*P. insidiosum* activity [15,120]. Therefore, the role of potential microbial metabolites in the treatment of pythiosis must be subjected to intense research in the future.

With the evidence of the effectiveness of some antimicrobials in the management of human pythiosis, we suggest using new drug delivery systems to release the drug to the target site in the body and minimize the off-target accumulation of the drug. Antibiotics can be reformulated using nanotechnology-derived delivery systems to improve the targeting and specificity at the infected areas [121]. Due to the genetic variability among individuals, not all individuals with pythiosis exhibit similar therapeutics responses to antimicrobials [122]. Therefore, it is essential to incorporate the pharmacogenomics assay into the clinics to personalize antimicrobial treatment in pythiosis.

Author Contributions: S.M. and N.W.; conceptualization, N.W. and A.C.; funding acquisition, S.M.; writing—original draft preparation, N.W. and A.C.; writing—review and editing. All authors have read and agreed to the published version of the manuscript.

Funding: This work was partially funded by the National Research Council of Thailand and Health Systems Research Institute (No. 64-087). The funder had no role in the study design, collection, and interpretation of the data or the decision to submit the work for publication.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: The authors would like to thank the members of the Mycology, Epidemiology, and Education Research Group (MERG), Sureerat Watcharasuwanseree from Udon Thani hospital, and the pythiosis patient for supporting the clinical part and clinical photograph of human pythiosis.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Permpalung, N.; Worasilchai, N.; Chindamporn, A. Human Pythiosis: Emergence of Fungal-Like Organism. *Mycopathologia* **2020**, *185*, 801–812, doi:10.1007/s11046-019-00412-0.
2. Gastra, W.; Lipman, L.J.; De Cock, A.W.; Exel, T.K.; Pegge, R.B.; Scheurwater, J.; Vilela, R.; Mendoza, L. *Pythium insidiosum*: an overview. *Vet Microbiol* **2010**, *146*, 1–16, doi:10.1016/j.vetmic.2010.07.019.
3. Mar Htun, Z.; Laikul, A.; Pathomsakulwong, W.; Yurayart, C.; Lohnoo, T.; Yingyong, W.; Kumsang, Y.; Payattikul, P.; Sae-Chew, P.; Rujirawat, T.; et al. Identification and Biotyping of *Pythium insidiosum* Isolated from Urban and Rural Areas of Thailand by Multiplex PCR, DNA Barcode, and Proteomic Analyses. *J Fungi (Basel)* **2021**, *7*, doi:10.3390/jof7040242.
4. Chitasombat, M.N.; Jongkhajornpong, P.; Lekhanont, K.; Krajaejun, T. Recent update in diagnosis and treatment of human pythiosis. *PeerJ* **2020**, *8*, e8555, doi:10.7717/peerj.8555.
5. Krajaejun, T.; Imkhieo, S.; Intaramat, A.; Ratanabanangkoon, K. Development of an immunochromatographic test for rapid serodiagnosis of human pythiosis. *Clin Vaccine Immunol* **2009**, *16*, 506–509, doi:10.1128/CVI.00276-08.

6. Zanette, R.A.; Bitencourt, P.E.R.; Alves, S.H.; Figuera, R.A.; Flores, M.M.; Wolkmer, P.; Hecktheuer, P.A.; Thomas, L.R.; Pereira, P.L.; Loreto, E.S.; et al. Insights into the pathophysiology of iron metabolism in *Pythium insidiosum* infections. *Vet Microbiol* **2013**, *162*, 826–830, doi:10.1016/j.vetmic.2012.10.036.
7. Chitasombat, M.N.; Larbcharoensub, N.; Chindamporn, A.; Krajaejun, T. Clinicopathological features and outcomes of pythiosis. *Int J Infect Dis* **2018**, *71*, 33–41, doi:10.1016/j.ijid.2018.03.021.
8. Mendoza, L.; Ajello, L.; McGinnis, M.R. Infections Caused by the Oomycetous Pathogen *Pythium insidiosum*. *Journal of Medical Mycology* **1996**, *6*, 151–164.
9. Yolanda, H.; Krajaejun, T. Review of methods and antimicrobial agents for susceptibility testing against *Pythium insidiosum*. *Heliyon* **2020**, *6*, e03737, doi:10.1016/j.heliyon.2020.e03737.
10. Mendoza, L.; Newton, J.C. Immunology and immunotherapy of the infections caused by *Pythium insidiosum*. *Med Mycol* **2005**, *43*, 477–486, doi:10.1080/13693780500279882.
11. Wanachiwanawin, W.; Mendoza, L.; Visuthisakchai, S.; Mutsikapan, P.; Sathapatayavongs, B.; Chaiprasert, A.; Suwanagool, P.; Manuskiatti, W.; Ruangsetakit, C.; Ajello, L. Efficacy of immunotherapy using antigens of *Pythium insidiosum* in the treatment of vascular pythiosis in humans. *Vaccine* **2004**, *22*, 3613–3621, doi:10.1016/j.vaccine.2004.03.031.
12. Yolanda, H.; Krajaejun, T. History and Perspective of Immunotherapy for Pythiosis. *Vaccines (Basel)* **2021**, *9*, doi:10.3390/vaccines9101080.
13. Tondolo, J.S.M.; Ledur, P.C.; Loreto, E.S.; Verdi, C.M.; Bitencourt, P.E.R.; de Jesus, F.P.K.; Rocha, J.P.; Alves, S.H.; Sasaki, G.L.; Santurio, J.M. Extraction, characterization and biological activity of a (1,3)(1,6)-beta-d-glucan from the pathogenic oomycete *Pythium insidiosum*. *Carbohydr Polym* **2017**, *157*, 719–727, doi:10.1016/j.carbpol.2016.10.053.
14. Permpalung, N.; Worasilchai, N.; Plongla, R.; Upala, S.; Sanguankeo, A.; Paitoonpong, L.; Mendoza, L.; Chindamporn, A. Treatment outcomes of surgery, antifungal therapy and immunotherapy in ocular and vascular human pythiosis: a retrospective study of 18 patients. *J Antimicrob Chemother* **2015**, *70*, 1885–1892, doi:10.1093/jac/dkv008.
15. Wittayapipath, K.; Yenjai, C.; Prariyachatigul, C.; Hamal, P. Evaluation of antifungal effect and toxicity of xanthyletin and two bacterial metabolites against Thai isolates of *Pythium insidiosum*. *Sci Rep* **2020**, *10*, 4495, doi:10.1038/s41598-020-61271-0.
16. Lerksuthirat, T.; Sangcakul, A.; Lohnoo, T.; Yingyong, W.; Rujirawat, T.; Krajaejun, T. Evolution of the Sterol Biosynthetic Pathway of *Pythium insidiosum* and Related Oomycetes Contributes to Antifungal Drug Resistance. *Antimicrob Agents Chemother* **2017**, *61*, doi:10.1128/AAC.02352-16.
17. Thom, K.A.; Schweizer, M.L.; Osih, R.B.; McGregor, J.C.; Furuno, J.P.; Perencevich, E.N.; Harris, A.D. Impact of empiric antimicrobial therapy on outcomes in patients with *Escherichia coli* and *Klebsiella pneumoniae* bacteremia: a cohort study. *BMC Infect Dis* **2008**, *8*, 116, doi:10.1186/1471-2334-8-116.
18. Rawson, T.M.; Wilson, R.C.; O'Hare, D.; Herrero, P.; Kambugu, A.; Lamorde, M.; Ellington, M.; Georgiou, P.; Cass, A.; Hope, W.W.; et al. Optimizing antimicrobial use: challenges, advances and opportunities. *Nat Rev Microbiol* **2021**, *19*, 747–758, doi:10.1038/s41579-021-00578-9.
19. Jarrell, A.S.; Kruer, R.M.; Johnson, D.; Lipsett, P.A. Antimicrobial Pharmacokinetics and Pharmacodynamics. *Surg Infect (Larchmt)* **2015**, *16*, 375–379, doi:10.1089/sur.2014.180.
20. Melida, H.; Sandoval-Sierra, J.V.; Dieguez-Uribeondo, J.; Bulone, V. Analyses of extracellular carbohydrates in oomycetes unveil the existence of three different cell wall types. *Eukaryot Cell* **2013**, *12*, 194–203, doi:10.1128/EC.00288-12.
21. Pires, L.; Bosco Sde, M.; Baptista, M.S.; Kurachi, C. Photodynamic therapy in *Pythium insidiosum* - an in vitro study of the correlation of sensitizer localization and cell death. *PLoS One* **2014**, *9*, e85431, doi:10.1371/journal.pone.0085431.
22. Kammarnjassadakul, P.; Rangsiapanurath, W.; Sriprapun, M.; Ngamsakulrungruj, P.; Palaga, T.; Sritunyalucksana, K.; Chindamporn, A. Cytochrome Oxidase 2 (COX2), β -Tubulin (TUB) and Chitin Synthase Subunit 2 (CHS2) Expression in *Pythium insidiosum* Thai Strains. *Walailak Journal of Science and Technology* **2021**, *18*, doi:10.48048/wjst.2021.9433.
23. Lerksuthirat, T.; Lohnoo, T.; Inkomlue, R.; Rujirawat, T.; Yingyong, W.; Khositnithikul, R.; Phaonakrop, N.; Roytrakul, S.; Sullivan, T.D.; Krajaejun, T. The elicitor-like glycoprotein, ELI025, is secreted by the pathogenic oomycete *Pythium insidiosum* and evades host antibody responses. *PLoS One* **2015**, *10*, e0118547, doi:10.1371/journal.pone.0118547.

24. Maeno, S.; Oie, Y.; Sunada, A.; Tanibuchi, H.; Hagiwara, S.; Makimura, K.; Nishida, K. Successful medical management of *Pythium insidiosum* keratitis using a combination of minocycline, linezolid, and chloramphenicol. *Am J Ophthalmol Case Rep* **2019**, *15*, 100498, doi:10.1016/j.ajoc.2019.100498.
25. Gurnani, B.; Narayana, S.; Christy, J.; Rajkumar, P.; Kaur, K.; Gubert, J. Successful management of pediatric *pythium insidiosum* keratitis with cyanoacrylate glue, linezolid, and azithromycin: Rare case report. *Eur J Ophthalmol* **2021**, 11206721211006564, doi:10.1177/11206721211006564.
26. Mercer, D.K.; Torres, M.D.T.; Duay, S.S.; Lovie, E.; Simpson, L.; von Kockritz-Blickwede, M.; de la Fuente-Nunez, C.; O'Neil, D.A.; Angeles-Boza, A.M. Antimicrobial Susceptibility Testing of Antimicrobial Peptides to Better Predict Efficacy. *Front Cell Infect Microbiol* **2020**, *10*, 326, doi:10.3389/fcimb.2020.00326.
27. Leekha, S.; Terrell, C.L.; Edson, R.S. General principles of antimicrobial therapy. *Mayo Clin Proc* **2011**, *86*, 156-167, doi:10.4065/mcp.2010.0639.
28. Worasilchai, N.; Chindamporn, A.; Plongla, R.; Torvorapanit, P.; Manothummetha, K.; Chuleerarux, N.; Permpalung, N. In Vitro Susceptibility of Thai *Pythium insidiosum* Isolates to Antibacterial Agents. *Antimicrob Agents Chemother* **2020**, *64*, doi:10.1128/AAC.02099-19.
29. Loreto, E.S.; Tondolo, J.S.; Pilotto, M.B.; Alves, S.H.; Santurio, J.M. New insights into the in vitro susceptibility of *Pythium insidiosum*. *Antimicrob Agents Chemother* **2014**, *58*, 7534-7537, doi:10.1128/AAC.02680-13.
30. Loreto, E.S.; Tondolo, J.S.M.; Santurio, J.M.; Alves, S.H. Screening of antibacterial drugs for antimicrobial activity against *Pythium insidiosum*. *Med Mycol* **2019**, *57*, 523-525, doi:10.1093/mmy/myy056.
31. Mahl, D.L.; de Jesus, F.P.; Loreto, E.; Zanette, R.A.; Ferreira, L.; Pilotto, M.B.; Alves, S.H.; Santurio, J.M. In vitro susceptibility of *Pythium insidiosum* isolates to aminoglycoside antibiotics and tigecycline. *Antimicrob Agents Chemother* **2012**, *56*, 4021-4023, doi:10.1128/AAC.00073-12.
32. Worasilchai, N.; Permpalung, N.; Chongsathidkiet, P.; Leelahavanichkul, A.; Mendoza, A.L.; Palaga, T.; Reantragoon, R.; Finkelman, M.; Sutcharitchan, P.; Chindamporn, A. Monitoring Anti-*Pythium insidiosum* IgG Antibodies and (1 \rightarrow 3)-beta-d-Glucan in Vascular Pythiosis. *J Clin Microbiol* **2018**, *56*, doi:10.1128/JCM.00610-18.
33. Richter, A.; Fessler, A.T.; Bottner, A.; Koper, L.M.; Wallmann, J.; Schwarz, S. Reasons for antimicrobial treatment failures and predictive value of in-vitro susceptibility testing in veterinary practice: An overview. *Vet Microbiol* **2020**, *245*, 108694, doi:10.1016/j.vetmic.2020.108694.
34. Bassetti, M.; Montero, J.G.; Paiva, J.A. When antibiotic treatment fails. *Intens Care Med* **2018**, *44*, 73-75, doi:10.1007/s00134-017-4962-2.
35. Cicchese, J.M.; Dartois, V.; Kirschner, D.E.; Linderman, J.J. Both Pharmacokinetic Variability and Granuloma Heterogeneity Impact the Ability of the First-Line Antibiotics to Sterilize Tuberculosis Granulomas. *Front Pharmacol* **2020**, *11*, 333, doi:10.3389/fphar.2020.00333.
36. Shah, S.; Barton, G.; Fischer, A. Pharmacokinetic considerations and dosing strategies of antibiotics in the critically ill patient. *J Intensive Care Soc* **2015**, *16*, 147-153, doi:10.1177/1751143714564816.
37. Veiga, R.P.; Paiva, J.A. Pharmacokinetics-pharmacodynamics issues relevant for the clinical use of beta-lactam antibiotics in critically ill patients. *Crit Care* **2018**, *22*, 233, doi:10.1186/s13054-018-2155-1.
38. Alghamdi, W.A.; Al-Shaer, M.H.; Klinker, K.P.; Peloquin, C.A. Variable linezolid exposure and response and the role of therapeutic drug monitoring: Case series. *Clin Case Rep* **2020**, *8*, 1126-1129, doi:10.1002/ccr3.2835.
39. Fugi, M.A.; Gunasekera, K.; Ochsenreiter, T.; Guan, X.; Wenk, M.R.; Maser, P. Genome profiling of sterol synthesis shows convergent evolution in parasites and guides chemotherapeutic attack. *J Lipid Res* **2014**, *55*, 929-938, doi:10.1194/jlr.M048017.
40. Levison, M.E.; Levison, J.H. Pharmacokinetics and pharmacodynamics of antibacterial agents. *Infect Dis Clin North Am* **2009**, *23*, 791-815, vii, doi:10.1016/j.idc.2009.06.008.
41. Jacobs, M.R. Optimisation of antimicrobial therapy using pharmacokinetic and pharmacodynamic parameters. *Clin Microbiol Infect* **2001**, *7*, 589-596, doi:10.1046/j.1198-743x.2001.00295.x.
42. Zhang, R.; Qin, X.; Kong, F.; Chen, P.; Pan, G. Improving cellular uptake of therapeutic entities through interaction with components of cell membrane. *Drug Deliv* **2019**, *26*, 328-342, doi:10.1080/10717544.2019.1582730.
43. Lee, C.R.; Cho, I.H.; Jeong, B.C.; Lee, S.H. Strategies to minimize antibiotic resistance. *Int J Environ Res Public Health* **2013**, *10*, 4274-4305, doi:10.3390/ijerph10094274.
44. Actor, J.K. Innate Immunity. In *Elsevier's Integrated Review Immunology and Microbiology*, Second ed.; Elsevier/Mosby: Philadelphia, USA, 2012; pp. 43-51.
45. Aristizábal, B.; González, A. Innate immune system. In *Autoimmunity: From Bench to Bedside [Internet]*, Anaya JM, S.Y., Rojas-Villarraga A, et al., Ed.; El Rosario University Press: Bogota (Colombia), 2013.

46. Bergman, P.; Raqib, R.; Rekha, R.S.; Agerberth, B.; Gudmundsson, G.H. Host Directed Therapy Against Infection by Boosting Innate Immunity. *Front Immunol* **2020**, *11*, 1209, doi:10.3389/fimmu.2020.01209.
47. Gjini, E.; Brito, P.H. Integrating Antimicrobial Therapy with Host Immunity to Fight Drug-Resistant Infections: Classical vs. Adaptive Treatment. *PLoS Comput Biol* **2016**, *12*, e1004857, doi:10.1371/journal.pcbi.1004857.
48. Zheng, W.J.; Xu, Q.; Zhang, Y.Y.; E, X.F.; Gao, W.; Zhang, M.G.; Zhai, W.J.; Rajkumar, R.S.; Liu, Z.J. Toll-like receptor-mediated innate immunity against herpesviridae infection: a current perspective on viral infection signaling pathways. *Virol J* **2020**, *17*, doi:10.1186/s12985-020-01463-2.
49. Wongprompitak, P.; Pleewan, N.; Tantibhedhyangkul, W.; Chaiprasert, A.; Prabhasawat, P.; Inthasin, N.; Ekpo, P. Involvement of Toll-like receptor 2 on human corneal epithelium during an infection of *Pythium insidiosum*. *Asian Pac J Allergy* **2020**, *38*, 129-138, doi:10.12932/Ap-110518-0311.
50. Garrido-Mesa, J.; Rodriguez-Nogales, A.; Algieri, F.; Vezza, T.; Hidalgo-Garcia, L.; Garrido-Barros, M.; Utrilla, M.P.; Garcia, F.; Chueca, N.; Rodriguez-Cabezas, M.E.; et al. Immunomodulatory tetracyclines shape the intestinal inflammatory response inducing mucosal healing and resolution. *Br J Pharmacol* **2018**, *175*, 4353-4370, doi:10.1111/bph.14494.
51. Yamauchi, K.; Shibata, Y.; Kimura, T.; Abe, S.; Inoue, S.; Osaka, D.; Sato, M.; Igarashi, A.; Kubota, I. Azithromycin suppresses interleukin-12p40 expression in lipopolysaccharide and interferon-gamma stimulated macrophages. *Int J Biol Sci* **2009**, *5*, 667-678, doi:10.7150/ijbs.5.667.
52. Takahashi, G.; Sato, N.; Yaegashi, Y.; Kojika, M.; Matsumoto, N.; Kikkawa, T.; Shozushima, T.; Akitomi, S.; Aoki, K.; Ito, N.; et al. Effect of linezolid on cytokine production capacity and plasma endotoxin levels in response to lipopolysaccharide stimulation of whole blood. *J Infect Chemother* **2010**, *16*, 94-99, doi:10.1007/s10156-009-0012-5.
53. Pichereau, S.; Moran, J.J.; Hayney, M.S.; Shukla, S.K.; Sakoulas, G.; Rose, W.E. Concentration-dependent effects of antimicrobials on *Staphylococcus aureus* toxin-mediated cytokine production from peripheral blood mononuclear cells. *J Antimicrob Chemother* **2012**, *67*, 123-129, doi:10.1093/jac/dkr417.
54. Lambers, C.; Burian, B.; Binder, P.; Ankersmit, H.J.; Wagner, C.; Muller, M.; Zeitlinger, M. Early immunomodulatory effects of linezolid in a human whole blood endotoxin model. *Int J Clin Pharmacol Ther* **2010**, *48*, 419-424, doi:10.5414/cpp48419.
55. Hirata, N.; Hiramatsu, K.; Kishi, K.; Yamasaki, T.; Ichimiya, T.; Nasu, M. Pretreatment of mice with clindamycin improves survival of endotoxic shock by modulating the release of inflammatory cytokines. *Antimicrob Agents Chemother* **2001**, *45*, 2638-2642, doi:10.1128/AAC.45.9.2638-2642.2001.
56. Veringa, E.M.; Lambe, D.W., Jr.; Ferguson, D.A., Jr.; Verhoef, J. Enhancement of opsonophagocytosis of *Bacteroides* spp. by clindamycin in subinhibitory concentrations. *J Antimicrob Chemother* **1989**, *23*, 577-587, doi:10.1093/jac/23.4.577.
57. Trostorf, F.; Reinert, R.R.; Schmidt, H.; Nichterlein, T.; Stuert, K.; Schmitz-Salue, M.; Sadowski, I.; Bruck, W.; Nau, R. Quinupristin/dalfopristin attenuates the inflammatory response and reduces the concentration of neuron-specific enolase in the cerebrospinal fluid of rabbits with experimental *Streptococcus pneumoniae* meningitis. *J Antimicrob Chemother* **1999**, *43*, 87-94, doi:10.1093/jac/43.1.87.
58. Majumdar, S.; Dutta, K.; Manna, S.K.; Basu, A.; Bishayi, B. Possible protective role of chloramphenicol in TSST-1 and coagulase-positive *Staphylococcus aureus*-induced septic arthritis with altered levels of inflammatory mediators. *Inflammation* **2011**, *34*, 269-282, doi:10.1007/s10753-010-9233-0.
59. Suschek, C.V.; Bonmann, E.; Kapsokefalou, A.; Hemmrich, K.; Kleinert, H.; Forstermann, U.; Kroncke, K.D.; Mahotka, C.; Kolb-Bachofen, V. Revisiting an old antimicrobial drug: amphotericin B induces interleukin-1-converting enzyme as the main factor for inducible nitric-oxide synthase expression in activated endothelia. *Mol Pharmacol* **2002**, *62*, 936-946, doi:10.1124/mol.62.4.936.
60. Mizuno, K.; Fukami, T.; Toyoda, Y.; Nakajima, M.; Yokoi, T. Terbinafine stimulates the pro-inflammatory responses in human monocytic THP-1 cells through an ERK signaling pathway. *Life Sci* **2010**, *87*, 537-544, doi:10.1016/j.lfs.2010.08.010.
61. Vora, S.; Purimetla, N.; Brummer, E.; Stevens, D.A. Activity of voriconazole, a new triazole, combined with neutrophils or monocytes against *Candida albicans*: effect of granulocyte colony-stimulating factor and granulocyte-macrophage colony-stimulating factor. *Antimicrob Agents Chemother* **1998**, *42*, 907-910, doi:10.1128/AAC.42.4.907.
62. Baltch, A.L.; Smith, R.P.; Franke, M.A.; Ritz, W.J.; Michelsen, P.B.; Bopp, L.H. Effects of cytokines and fluconazole on the activity of human monocytes against *Candida albicans*. *Antimicrob Agents Chemother* **2001**, *45*, 96-104, doi:10.1128/AAC.45.1.96-104.2001.

63. Itaqui, S.R.; Verdi, C.M.; Tondolo, J.S.; da Luz, T.S.; Alves, S.H.; Santurio, J.M.; Loreto, E.S. In Vitro Synergism between Azithromycin or Terbinafine and Topical Antimicrobial Agents against *Pythium insidiosum*. *Antimicrob Agents Chemother* **2016**, *60*, 5023–5025, doi:10.1128/AAC.00154-16.
64. Jesus, F.P.; Ferreira, L.; Bizzi, K.S.; Loreto, E.S.; Pilotto, M.B.; Ludwig, A.; Alves, S.H.; Zanette, R.A.; Santurio, J.M. In vitro activity of carvacrol and thymol combined with antifungals or antibacterials against *Pythium insidiosum*. *J Mycol Med* **2015**, *25*, e89–93, doi:10.1016/j.mycmed.2014.10.023.
65. Jesus, F.P.; Ferreira, L.; Loreto, E.S.; Pilotto, M.B.; Ludwig, A.; Bizzi, K.; Tondolo, J.S.; Zanette, R.A.; Alves, S.H.; Santurio, J.M. In vitro synergism observed with azithromycin, clarithromycin, minocycline, or tigecycline in association with antifungal agents against *Pythium insidiosum*. *Antimicrob Agents Chemother* **2014**, *58*, 5621–5625, doi:10.1128/AAC.02349-14.
66. Chopra, I.; Roberts, M. Tetracycline antibiotics: mode of action, applications, molecular biology, and epidemiology of bacterial resistance. *Microbiol Mol Biol Rev* **2001**, *65*, 232–260 ; second page, table of contents, doi:10.1128/MMBR.65.2.232-260.2001.
67. Nakao, C.; Angel, M.; Mateo, S.D.; Komesu, M.C. Effects of Topical Tetracycline in Wound Healing on Experimental Diabetes in Rats. *The Open Diabetes Journal* **2009**, *2*, 53–59.
68. Zimmermann, C.E.P.; Jesus, F.P.K.; Schlemmer, K.B.; Loreto, E.S.; Tondolo, J.S.M.; Driemeier, D.; Alves, S.H.; Ferreira, L.; Santurio, J.M. In vivo effect of minocycline alone and in combination with immunotherapy against *pythium insidiosum*. *Vet Microbiol* **2020**, *243*, 108616, doi:10.1016/j.vetmic.2020.108616.
69. Patel, P.H.; Hashmi, M.F. Macrolides. In *StatPearls*; Treasure Island (FL), 2021.
70. Cramer, C.L.; Patterson, A.; Alchakaki, A.; Soubani, A.O. Immunomodulatory indications of azithromycin in respiratory disease: a concise review for the clinician. *Postgrad Med* **2017**, *129*, 493–499, doi:10.1080/00325481.2017.1285677.
71. Allen, J.E.; Wynn, T.A. Evolution of Th2 immunity: a rapid repair response to tissue destructive pathogens. *PLoS Pathog* **2011**, *7*, e1002003, doi:10.1371/journal.ppat.1002003.
72. Chen, F.; Liu, Z.; Wu, W.; Roza, C.; Bowdridge, S.; Millman, A.; Van Rooijen, N.; Urban, J.F., Jr.; Wynn, T.A.; Gause, W.C. An essential role for TH2-type responses in limiting acute tissue damage during experimental helminth infection. *Nat Med* **2012**, *18*, 260–266, doi:10.1038/nm.2628.
73. Kanoh, S.; Rubin, B.K. Mechanisms of action and clinical application of macrolides as immunomodulatory medications. *Clin Microbiol Rev* **2010**, *23*, 590–615, doi:10.1128/CMR.00078-09.
74. Jesus, F.P.; Loreto, E.S.; Ferreira, L.; Alves, S.H.; Driemeier, D.; Souza, S.O.; Franca, R.T.; Lopes, S.T.; Pilotto, M.B.; Ludwig, A.; et al. In Vitro and In Vivo Antimicrobial Activities of Minocycline in Combination with Azithromycin, Clarithromycin, or Tigecycline against *Pythium insidiosum*. *Antimicrob Agents Chemother* **2016**, *60*, 87–91, doi:10.1128/AAC.01480-15.
75. Bozdogan, B.; Appelbaum, P.C. Oxazolidinones: activity, mode of action, and mechanism of resistance. *Int J Antimicrob Agents* **2004**, *23*, 113–119, doi:10.1016/j.ijantimicag.2003.11.003.
76. Wang, J.; Xia, L.; Wang, R.; Cai, Y. Linezolid and Its Immunomodulatory Effect: In Vitro and In Vivo Evidence. *Front Pharmacol* **2019**, *10*, 1389, doi:10.3389/fphar.2019.01389.
77. Ahirwar, L.K.; Kalra, P.; Sharma, S.; Mohamed, A.; Mittal, R.; Das, S.; Bagga, B. Linezolid shows high safety and efficacy in the treatment of *Pythium insidiosum* keratitis in a rabbit model. *Exp Eye Res* **2021**, *202*, 108345, doi:10.1016/j.exer.2020.108345.
78. Schwarz, S.; Shen, J.; Kadlec, K.; Wang, Y.; Brenner Michael, G.; Fessler, A.T.; Vester, B. Lincosamides, Streptogramins, Phenicol, and Pleuromutilins: Mode of Action and Mechanisms of Resistance. *Cold Spring Harb Perspect Med* **2016**, *6*, doi:10.1101/cshperspect.a027037.
79. Krause, K.M.; Serio, A.W.; Kane, T.R.; Connolly, L.E. Aminoglycosides: An Overview. *Cold Spring Harb Perspect Med* **2016**, *6*, doi:10.1101/cshperspect.a027029.
80. McMeekin, D.; Mendoza, L. In vitro effect of streptomycin on clinical isolates of *Pythium insidiosum*. *Mycologia* **2000**, *92*, 371–373.
81. McOsker, C.C.; Fitzpatrick, P.M. Nitrofurantoin: mechanism of action and implications for resistance development in common uropathogens. *J Antimicrob Chemother* **1994**, *33 Suppl A*, 23–30, doi:10.1093/jac/33.suppl_a.23.
82. Parenti, M.A.; Hatfield, S.M.; Leyden, J.J. Mupirocin: a topical antibiotic with a unique structure and mechanism of action. *Clin Pharm* **1987**, *6*, 761–770.
83. Noor, A.; Preuss, C.V. Amphotericin B. In *StatPearls*; Treasure Island (FL), 2021.
84. Triscott, J.A.; Weedon, D.; Cabana, E. Human subcutaneous pythiosis. *J Cutan Pathol* **1993**, *20*, 267–271, doi:10.1111/j.1600-0560.1993.tb00654.x.

85. Krajaejun, T.; Sathapatayavongs, B.; Prachartam, R.; Nitiyanant, P.; Leelachaikul, P.; Wanachiwanawin, W.; Chaiprasert, A.; Assanasen, P.; Saipetch, M.; Mootsikapun, P.; et al. Clinical and epidemiological analyses of human pythiosis in Thailand. *Clin Infect Dis* **2006**, *43*, 569–576, doi:10.1086/506353.
86. Sekhon, A.S.; Padhye, A.A.; Garg, A.K. In vitro sensitivity of *Penicillium marneffei* and *Pythium insidiosum* to various antifungal agents. *Eur J Epidemiol* **1992**, *8*, 427–432, doi:10.1007/BF00158578.
87. Permpalung, N.; Worasilchai, N.; Manothummetha, K.; Torvorapanit, P.; Ratanawongphaibul, K.; Chuleerarux, N.; Plongla, R.; Chindamporn, A. Clinical outcomes in ocular pythiosis patients treated with a combination therapy protocol in Thailand: A prospective study. *Med Mycol* **2019**, *57*, 923–928, doi:10.1093/mmy/myz013.
88. Mesa-Arango, A.C.; Scorzoni, L.; Zaragoza, O. It only takes one to do many jobs: Amphotericin B as antifungal and immunomodulatory drug. *Front Microbiol* **2012**, *3*, 286, doi:10.3389/fmicb.2012.00286.
89. Malone-Povolny, M.J.; Maloney, S.E.; Schoenfisch, M.H. Nitric Oxide Therapy for Diabetic Wound Healing. *Adv Healthc Mater* **2019**, *8*, e1801210, doi:10.1002/adhm.201801210.
90. Luo, J.D.; Chen, A.F. Nitric oxide: a newly discovered function on wound healing. *Acta Pharmacol Sin* **2005**, *26*, 259–264, doi:10.1111/j.1745-7254.2005.00058.x.
91. Vriens, K.; Kumar, P.T.; Struyfs, C.; Cools, T.L.; Spincemaille, P.; Kokalj, T.; Sampaio-Marques, B.; Ludovico, P.; Lammertyn, J.; Cammue, B.P.A.; et al. Increasing the Fungicidal Action of Amphotericin B by Inhibiting the Nitric Oxide-Dependent Tolerance Pathway. *Oxid Med Cell Longev* **2017**, *2017*, 4064628, doi:10.1155/2017/4064628.
92. Mesa-Arango, A.C.; Trevijano-Contador, N.; Roman, E.; Sanchez-Fresneda, R.; Casas, C.; Herrero, E.; Arguelles, J.C.; Pla, J.; Cuenca-Estrella, M.; Zaragoza, O. The production of reactive oxygen species is a universal action mechanism of Amphotericin B against pathogenic yeasts and contributes to the fungicidal effect of this drug. *Antimicrob Agents Chemother* **2014**, *58*, 6627–6638, doi:10.1128/AAC.03570-14.
93. Ryder, N.S. Terbinafine: mode of action and properties of the squalene epoxidase inhibition. *Br J Dermatol* **1992**, *126 Suppl 39*, 2–7, doi:10.1111/j.1365-2133.1992.tb00001.x.
94. Kanafani, Z.A.; Perfect, J.R. Antimicrobial resistance: resistance to antifungal agents: mechanisms and clinical impact. *Clin Infect Dis* **2008**, *46*, 120–128, doi:10.1086/524071.
95. Ghannoum, M.A.; Rice, L.B. Antifungal agents: mode of action, mechanisms of resistance, and correlation of these mechanisms with bacterial resistance. *Clin Microbiol Rev* **1999**, *12*, 501–517, doi:10.1128/CMR.12.4.501.
96. Sussaengrat, N.; Torvorapanit, P.; Plongla, R.; Chuleerarux, N.; Manothummetha, K.; Tuangsirisup, J.; Worasilchai, N.; Chindamporn, A.; Permpalung, N. Adjunctive antibacterial agents as a salvage therapy in relapsed vascular pythiosis patients. *Int J Infect Dis* **2019**, *88*, 27–30, doi:10.1016/j.ijid.2019.08.032.
97. Cavalheiro, A.S.; Maboni, G.; de Azevedo, M.I.; Argenta, J.S.; Pereira, D.I.; Spader, T.B.; Alves, S.H.; Santurio, J.M. In Vitro activity of terbinafine combined with caspofungin and azoles against *Pythium insidiosum*. *Antimicrob Agents Chemother* **2009**, *53*, 2136–2138, doi:10.1128/AAC.01506-08.
98. Shenep, J.L.; English, B.K.; Kaufman, L.; Pearson, T.A.; Thompson, J.W.; Kaufman, R.A.; Frisch, G.; Rinaldi, M.G. Successful medical therapy for deeply invasive facial infection due to *Pythium insidiosum* in a child. *Clin Infect Dis* **1998**, *27*, 1388–1393, doi:10.1086/515042.
99. Sucher, A.J.; Chahine, E.B.; Balcer, H.E. Echinocandins: the newest class of antifungals. *Ann Pharmacother* **2009**, *43*, 1647–1657, doi:10.1345/aph.1M237.
100. Lamaris, G.A.; Lewis, R.E.; Chamilos, G.; May, G.S.; Safdar, A.; Walsh, T.J.; Raad, II; Kontoyiannis, D.P. Caspofungin-mediated beta-glucan unmasking and enhancement of human polymorphonuclear neutrophil activity against *Aspergillus* and non-*Aspergillus* hyphae. *J Infect Dis* **2008**, *198*, 186–192, doi:10.1086/589305.
101. Katragkou, A.; Roilides, E.; Walsh, T.J. Role of Echinocandins in Fungal Biofilm-Related Disease: Vascular Catheter-Related Infections, Immunomodulation, and Mucosal Surfaces. *Clin Infect Dis* **2015**, *61 Suppl 6*, S622–629, doi:10.1093/cid/civ746.
102. Zanette, R.A.; Jesus, F.P.; Pilotto, M.B.; Weiblen, C.; Potter, L.; Ferreira, L.; Alves, S.H.; Santurio, J.M. Micafungin alone and in combination therapy with deferasirox against *Pythium insidiosum*. *J Mycol Med* **2015**, *25*, 91–94, doi:10.1016/j.mycmed.2014.09.002.
103. Pereira, D.I.; Santurio, J.M.; Alves, S.H.; Argenta, J.S.; Potter, L.; Spanamberg, A.; Ferreira, L. Caspofungin in vitro and in vivo activity against Brazilian *Pythium insidiosum* strains isolated from animals. *J Antimicrob Chemother* **2007**, *60*, 1168–1171, doi:10.1093/jac/dkm332.
104. Polak, A.M. Preclinical data and mode of action of amorolfine. *Clin Exp Dermatol* **1992**, *17 Suppl 1*, 8–12, doi:10.1111/j.1365-2230.1992.tb00270.x.

105. Sahoo, B.M.; Ravi Kumar, B.V.V.; Sruti, J.; Mahapatra, M.K.; Banik, B.K.; Borah, P. Drug Repurposing Strategy (DRS): Emerging Approach to Identify Potential Therapeutics for Treatment of Novel Coronavirus Infection. *Front Mol Biosci* **2021**, *8*, 628144, doi:10.3389/fmolb.2021.628144.
106. Yacouba, A.; Olowo-Okere, A.; Yunusa, I. Repurposing of antibiotics for clinical management of COVID-19: a narrative review. *Ann Clin Microbiol Antimicrob* **2021**, *20*, 37, doi:10.1186/s12941-021-00444-9.
107. Kaul, G.; Shukla, M.; Dasgupta, A.; Chopra, S. Update on drug-repurposing: is it useful for tackling antimicrobial resistance? *Future Microbiol* **2019**, *14*, 829–831, doi:10.2217/fmb-2019-0122.
108. Stokes, M.; Abdijadid, S. Disulfiram. In *StatPearls [Internet]*; Treasure Island (FL), 2021.
109. Krajaejun, T.; Lohnoo, T.; Yingyong, W.; Rujirawat, T.; Kumsang, Y.; Jongkhajornpong, P.; Theerawatanasirikul, S.; Kittichotirat, W.; Reamtong, O.; Yolanda, H. The Repurposed Drug Disulfiram Inhibits Urease and Aldehyde Dehydrogenase and Prevents In Vitro Growth of the Oomycete *Pythium insidiosum*. *Antimicrob Agents Chemother* **2019**, *63*, doi:10.1128/AAC.00609-19.
110. Pushpakom, S.; Iorio, F.; Eyers, P.A.; Escott, K.J.; Hopper, S.; Wells, A.; Doig, A.; Williams, T.; Latimer, J.; McNamee, C.; et al. Drug repurposing: progress, challenges and recommendations. *Nat Rev Drug Discov* **2019**, *18*, 41–58, doi:10.1038/nrd.2018.168.
111. Li, J.; Zheng, S.; Chen, B.; Butte, A.J.; Swamidass, S.J.; Lu, Z. A survey of current trends in computational drug repositioning. *Brief Bioinform* **2016**, *17*, 2–12, doi:10.1093/bib/bbv020.
112. Ko, Y. Computational Drug Repositioning: Current Progress and Challenges. *Appl. Sci.* **2020**, *10*, 5076, doi:10.3390/app10155076.
113. Rudrapal, M.; Khairnar, S.; Jadhav, A. Drug Repurposing (DR): An Emerging Approach in Drug Discovery. In *Drug Repurposing - Hypothesis, Molecular Aspects and Therapeutic Applications*; London, UK, 2020.
114. Liu, Y.; Tong, Z.; Shi, J.; Li, R.; Upton, M.; Wang, Z. Drug repurposing for next-generation combination therapies against multidrug-resistant bacteria. *Theranostics* **2021**, *11*, 4910–4928, doi:10.7150/thno.56205.
115. Cheng, Y.S.; Williamson, P.R.; Zheng, W. Improving therapy of severe infections through drug repurposing of synergistic combinations. *Curr Opin Pharmacol* **2019**, *48*, 92–98, doi:10.1016/j.coph.2019.07.006.
116. Nguyen, T.K.; Selvanayagam, R.; Ho, K.K.K.; Chen, R.; Kutty, S.K.; Rice, S.A.; Kumar, N.; Barraud, N.; Duong, H.T.T.; Boyer, C. Co-delivery of nitric oxide and antibiotic using polymeric nanoparticles. *Chem Sci* **2016**, *7*, 1016–1027, doi:10.1039/c5sc02769a.
117. Rouillard, K.R.; Novak, O.P.; Pistiolis, A.M.; Yang, L.; Ahonen, M.J.R.; McDonald, R.A.; Schoenfisch, M.H. Exogenous Nitric Oxide Improves Antibiotic Susceptibility in Resistant Bacteria. *ACS Infect Dis* **2021**, *7*, 23–33, doi:10.1021/acsinfecdis.0c00337.
118. Available online: <https://www.sanfordguide.com/> (accessed on 3/4/2022).
119. Keeratijarut, A.; Lohnoo, T.; Rujirawat, T.; Yingyong, W.; Kalambaheti, T.; Miller, S.; Phuntumart, V.; Krajaejun, T. The Immunoreactive Exo-1,3-beta-Glucanase from the Pathogenic Oomycete *Pythium insidiosum* Is Temperature Regulated and Exhibits Glycoside Hydrolase Activity. *PLoS One* **2015**, *10*, e0135239, doi:10.1371/journal.pone.0135239.
120. Wittayapipath, K.; Laolit, S.; Yenjai, C.; Chio-Srichan, S.; Pakarasang, M.; Tavichakorntrakool, R.; Prariyachatigul, C. Analysis of xanthyletin and secondary metabolites from *Pseudomonas stutzeri* ST1302 and *Klebsiella pneumoniae* ST2501 against *Pythium insidiosum*. *BMC Microbiol* **2019**, *19*, 78, doi:10.1186/s12866-019-1452-4.
121. Subramaniam, S.; Joyce, P.; Thomas, N.; Prestidge, C.A. Bioinspired drug delivery strategies for repurposing conventional antibiotics against intracellular infections. *Adv Drug Deliv Rev* **2021**, *177*, 113948, doi:10.1016/j.addr.2021.113948.
122. Stocco, G.; Lucafo, M.; Decorti, G. Pharmacogenomics of Antibiotics. *Int J Mol Sci* **2020**, *21*, doi:10.3390/ijms21175975.