

**Table S2.** Distribution of minimum inhibitory concentration (MIC) of eight antimicrobial agents; MIC<sub>50</sub> and MIC<sub>90</sub> values for the studied *Rhodococcus equi* isolates from pigs (n=17).

Antimicrobial agents <sup>a</sup>	Number of Isolates with MIC (µg/mL) <sup>b</sup>												MIC <sub>50</sub>	MIC <sub>90</sub>
	0.75	1	1.5	2	3	4	6	8	12	16	24	>32		
PEN				2	1	3	2		1			8	12	>32
AMC	3	3	6	3	2								1.5	2
CTX						3	2	1				11	>32	>32
	<b>0.25</b>	<b>0.38</b>	<b>0.5</b>	<b>0.75</b>	<b>1</b>	<b>1.5</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>8</b>	<b>&gt;32</b>		
CIP	1	4	1	4	4	2						1	0.75	1.5
GEN	1	6	4	5							1		0.5	0.75
	<b>0.125</b>	<b>0.19</b>	<b>0.25</b>	<b>0.38</b>	<b>0.5</b>	<b>0.75</b>	<b>1</b>	<b>1.5</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>6</b>		
ERY	2	5	4	4	2								0.25	0.38
	<b>0.5</b>	<b>0.75</b>	<b>1.5</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>8</b>	<b>12</b>	<b>16</b>	<b>24</b>	<b>&gt;32</b>		
SXT	1		2	6			1	1	1			5	2	>32
	<b>1.5</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>6</b>	<b>8</b>	<b>12</b>	<b>16</b>	<b>24</b>	<b>32</b>	<b>48</b>	<b>&gt;256</b>		
DOX	1			3	4	6	3						8	12

<sup>a</sup> PEN – penicillin, AMC – amoxicillin/clavulanic acid, CTX – cefotaxime, CIP – ciprofloxacin, ERY – erythromycin, DOX – doxycycline, GEN – gentamicin, SXT – trimethoprim-sulfamethoxazole; <sup>b</sup> MIC range for tested antimicrobial agents.