

Supplementary Material

Solution-Mediated Modulation of *P. aeruginosa* Biofilm Formation by a Cationic Synthetic Polymer

Leanna Foster¹, Shin-ichi Yusa² and Kenichi Kuroda^{1,3,*}

¹ Macromolecular Science and Engineering, University of Michigan, Ann Arbor, MI, 48109; llfoster@umich.edu

² Department of Materials Science and Chemistry, University of Hyogo, 2167 Shosha, Himeji, Hyogo 671-2280, Japan; yusa@eng.u-hyogo.ac.jp

³ Department of Biologic and Materials Sciences & Prosthodontics, School of Dentistry, University of Michigan, Ann Arbor, MI 48109; kkuroda@umich.edu

* Correspondence: kkuroda@umich.edu




Figure S1. ¹H NMR spectra for (a) P-1 and (b) P-2 in D₂O. The ¹H NMR spectra were obtained with Bruker DRX-500.




Figure S2. GPC elution curves for (a) P-1 using two Shodex Ohpak SB-804 HQ columns and a 0.3 M Na_2SO_4 aqueous solution containing 0.5 M acetic acid as an eluent and (b) P-2 using a Shodex Asahipak GF-7M HQ column and a phosphate buffer (pH 9) containing 10 vol% acetonitrile as an eluent.