

Communication

Comparative Evaluation of the Antimicrobial and Mucus Induction Properties of Selected *Bacillus* Strains Against Enterotoxigenic *Escherichia coli*

Natalia Bravo Santano ¹, Erik Juncker Boll ², Lena Catrine Capern ², Tomasz Maciej Cieplak ², Enver Keleszade ¹, Michal Letek ^{3,†} and Adele Costabile ^{1,*}

¹ Department of Life Sciences, University of Roehampton, London, UK; nbravasantano@gmail.com (N.B.S.); keleszae@roehampton.ac.uk (E.K.)

² Animal Health Innovation, Chr. Hansen A/S, Hørsholm, Denmark; dkerbo@chr-hansen.com (E.J.B.); dkleca@chr-hansen.com (L.C.C.); dktoci@chr-hansen.com (T.M.C.)

³ Departamento de Biología Molecular, Facultad de Ciencias Biológicas y Ambientales, Universidad de León, León, Spain; michal.letek@unileon.es

* Correspondence: adele.costabile@roehampton.ac.uk, Tel.: +44-(0)-20-8392-3571

† These authors share co-senior authorship.

Received: 8 November 2020; Accepted: 25 November 2020; Published: date

Supplementary materials




Figure S1. Quantification of MUC2 and MUC3 in HT29-16E cells after treatment with *Bacillus subtilis* strain CHCC 15541. HT29-16E cells were incubated with CHCC 15541 for 6, 8, and 10 hours, and concentrations of both mucin 2 (MUC2) and mucin 3 (MUC3) were quantified by ELISA at these time points. Data are expressed as Mean \pm SEM of one experiment performed in duplicates.