

Table S1 Primers used in this study

Gene	Forward primer (5'-3')	Reverse primer (5'-3')
Primer used for plasmid construction		
RpSP13.8	ACGAGCTGTACAAGGGTACCATGAAATATCCCAAGACAGCTTGC	GCGGACTCTAGTTCATCTAGATTATCTTCCGTAGATTGTAATA
RpSP30.2	ACGAGCTGTACAAGGGTACCATGGAACCTCTGCGCCAGAAAAGG	GCGGACTCTAGTTCATCTAGATTATGCGGCAGCTTGAGGCCT
RpSP13.4	ACGAGCTGTACAAGGGTACCATGTTACGAGCCTCATCAAAGCG	GCGGACTCTAGTTCATCTAGATTAAATTCCCAGACTCAGCCTG
RpSP10.3	ACGAGCTGTACAAGGGTACCATGGAGCCTTTGTTTCTGAAGCGTG	GCGGACTCTAGTTCATCTAGATCAGCAGTCATCTTCGCATTC
RpSP17.8	ACGAGCTGTACAAGGGTACCATGCAGAGAAATACCATTGACAATGC	GCGGACTCTAGTTCATCTAGATTAAACCTTGTGGTGATGGTCCC
Primer used for qPCR		
NPR1	TGCTCTCCATTATGCTGTAG	GCACTGTGTATCCTCTTGAA
PDF1.2	TGGCAAAATCTATGCGCTTT	ATCCTTCGGTCAAACAGACG
PR1	TGAGATGTGGGTCGATGAGA	CGAGTTACGCCAAACCACTT
PR4	GGCCAAGATTCCTGTGGTAGAT	CACTGTTGTTTGAGTTCCTGTTCT
actin	GGAGAAGTTGGCTTACATTG	TCATTGATGGTTGGAACAGA