

Article

Detection of Carbapenem Resistance of *Proteus mirabilis* Strains Isolated from Foxes, Raccoons and Minks in China

Penghao Lv ^{1,†}, Guijuan Hao ^{1,*}, Yanli Cao ¹, Lulu Cui ¹, Guisheng Wang ^{2,*} and Shuhong Sun ^{1,*}

¹ Department of Preventive Veterinary Medicine, College of Veterinary Medicine, Shandong Agricultural University, Taian 271018, China; 2019110426@sda.edu.cn (P.L.); sdaucyl@163.com (Y.C.); 2021010090@sda.edu.cn (L.C.)

² Shandong Animal Disease Prevention and Control Center, Taian 261500, China

* Correspondence: haogj2020@sda.edu.cn (G.H.); wangguisheng@shandong.cn (G.W.); sunshuhong@sda.edu.cn (S.S.); Tel.: +86-18252026546 (G.H.); +86-18560113839 (G.W.); +86-13705389710 (S.S.)

† These authors contributed equally to this work.

Simple Summary: *Proteus* is found abundantly in soil and water, and it has been known to cause human urinary tract infections and food poisoning. Currently, the opportunistic pathogen *Proteus mirabilis* (*P. mirabilis*) is also found to be an emerging threat to animals, such as birds, fish, dogs, etc. In this study, we examined the antibiotic resistance genes and virulence genes of *P. mirabilis* isolates from raccoons, foxes and minks. Among a total of 53 *P. mirabilis* isolates, the proportion of bacteria resistant to three or more antibiotic classes was up to 73.58%, and the detection rate of carbapenem-resistant *P. mirabilis* isolates was up to 71.7%, putting human health at risk. The close evolutionary relationship between *P. mirabilis* isolates from animals and the farm environment suggested that multidrug-resistant *P. mirabilis* from animals could pose a great environmental threat. In addition, the carriage rate of virulence-associated genes was not positively correlated with *P. mirabilis* pathogenicity in a *Galleria mellonella* model, highlighting the importance of further understanding the virulence of *P. mirabilis* in future studies.

Abstract: *Proteus mirabilis*, an opportunistic pathogen, is found to be an emerging threat to both animals and humans for a variety of infections. However, the characteristics of *P. mirabilis* infections from foxes, raccoons and minks remain unclear. In this context, we identified the antibiotic resistance genes and virulence genes of *P. mirabilis* isolates from foxes, raccoons and minks in China. Most isolates showed resistance to florfenicol (90.57%), trimethoprim-sulfamethoxazole (73.58%), and imipenem (71.70%). A total of 73.58% of isolates were resistant to antibiotics from at least three or more classes, and were categorized as multi-drug resistant. A total of 33.33% of the isolates were resistant to antibiotics from seven classes. The most prevalent resistant were *sul1* (94.34%), followed by *floR*, *bla_{TEM}*, *aac(6')Ib-cr* and *bla_{OXA-1}* with the detection rate of 88.68%, 83.02%, 71.70% and 60.38%, respectively. Among the 51 *P. mirabilis* isolates that were resistant to beta-lactam antibiotics, all isolates carried at least one beta-lactam gene. In addition, *bla_{NDM}* and *bla_{OXA-24}* genes were firstly reported in carbapenem-resistant *P. mirabilis* isolates from foxes, raccoons and minks. All isolates exhibited the virulence genes *ureC*, *zapA*, *pmfA*, *atfA* and *mrpA*. *P. mirabilis* isolates carrying all detected 10 virulence genes from different animal species showed different lethal abilities in a *G. mellonella* larvae model. More importantly, the profiles of antibiotic resistance genes of isolates from fur animals and the environment were generally similar, and phylogenetic analysis showed that the *P. mirabilis* isolates from farm environment samples may have close relatedness with that from animals.

Keywords: *Proteus mirabilis*; antimicrobial resistance; carbapenem resistance; virulence genes

Citation: Lv, P.; Hao, G.; Cao, Y.; Cui, L.; Wang, G.; Sun, S. Detection of Carbapenem Resistance of *Proteus mirabilis* Strains Isolated from Foxes, Raccoons and Minks in China. *Biology* **2022**, *11*, 292. <https://doi.org/10.3390/biology11020292>

Academic Editor: Chrissoula Voudarou

Received: 20 December 2021

Accepted: 8 February 2022

Published: 11 February 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Proteus mirabilis (*P. mirabilis*), a Gram-negative bacterium with flagella and fimbriae, is widely found in water, soil, and the intestinal microbiota of animals and humans. As an opportunistic pathogen, *P. mirabilis* is usually harmless to human health [1]. However, when the body's immunity is weakened, *P. mirabilis* may move to non-intestinal sites and cause serious diseases, such as cystitis, sepsis, peritonitis and meningitis [2,3]. After *Escherichia coli* (*E. coli*) and *Klebsiella pneumoniae* (*K. pneumoniae*), *P. mirabilis* is the third most common cause of human urinary tract infections [4]. *P. mirabilis* is also involved in nosocomial outbreaks in neonates in India [3]. In addition, although *Proteus* does not belong to the foodborne pathogens, some food poisoning cases associated with *P. mirabilis* have been reported [5,6]. Gong et al. [6] demonstrated that *P. mirabilis* from humans could induce the symptoms of both vomiting and diarrhea in a mouse model and show high gastrointestinal pathogenicity. Comparative genomic analysis showed that *P. mirabilis* may acquire horizontally from other microbial genomes toxin genes to exert digestion tract infection and toxicity [7]. Currently, the opportunistic pathogen *P. mirabilis* is found to be an emerging threat to animals. It was reported that *P. mirabilis* could cause large-scale mortality of fish without association with any other pathogens [8]. In addition, *P. mirabilis* infections have been reported in birds with reproductive failure [9], cattle and fowl with diarrhea [10,11] and dogs with chronic otitis externa [12].

Carbapenems, such as imipenem and meropenem, are bactericidal beta-lactam antimicrobials with proven efficacy in severe infections caused by extended spectrum beta-lactamases (ESBLs, e.g., TEM, SHV, CTX-M enzymes) producing bacteria [13]. Carbapenem resistance occurs mainly among Gram-negative pathogens such as *Enterobacteriaceae*, *Pseudomonas* spp. and *Acinetobacter* spp., which is mainly due to the production of carbapenemase including some class A beta-lactamases (e.g., KPC, GES), class B metallo-beta-lactamases (e.g., NDM, IMP) and class D carbapenemases (e.g., OXA-23, -24, -48, -51, and -58) [14]. KPC-type producing *P. mirabilis* was identified for the first time in the blood culture of a diabetic patient in the United States in 2008; since then, KPC-producing *P. mirabilis* isolates have been found in humans in many countries [15–17]. NDM-producing *P. mirabilis* isolates have been isolated from commercial broilers in slaughterhouses, ducks, wildlife, and hospitalized patients in China [18–21]. Among the carbapenem-hydrolyzing class D beta-lactamases, OXA-48, the most frequently identified in *Enterobacteriales*, has only been very rarely reported in *Proteus* spp., whereas OXA-23 and OXA-58, exclusively identified in *Acinetobacter* species, have recently been increasingly identified in *P. mirabilis* from hospital settings, the community, and cattle feces [17,22,23]. In addition, the *bla*_{OXA-24} has also been identified in *P. mirabilis* isolates from hospital clinical specimens in Algeria [22]. Since *P. mirabilis* is intrinsically resistant to tetracyclines and polymyxins, the emergence of multidrug-resistant, or even extensively drug-resistant *P. mirabilis* isolates carrying carbapenemase genes complicated the clinical treatment of bacterial infections, causing a significant public health concern. Most previous studies were focused on the prevalence of antimicrobial resistance of *P. mirabilis* in human urinary tract infections, and chicken, beef, and pork meat [19,21,23,24]. However, there are few reports on the characteristics of *P. mirabilis* from foxes, raccoons and minks.

The present study aimed to characterize the antibiotic resistance profiles of 53 *P. mirabilis* isolates obtained from fox, raccoon and mink farms in China, and to compare the genotypic and phenotypic characteristics of antimicrobial resistance and virulence factors in these isolates.

2. Materials and Methods

2.1. Bacterial Isolates and Growth Conditions

A total of 53 *P. mirabilis* isolates were used in this study. They were isolated from fox, raccoon, or mink farms in different cities of Shandong Province, China between October

2019 and November 2020 (Table 1). Detailed information on these *P. mirabilis* isolates is provided in Tables S1 and S2 in the Supplementary Materials. The standard *Salmonella enterica* serovar Enteritidis (*S. Enteritidis*) strain CVCC3377 was purchased from the China Veterinary Culture Collection Center (Beijing, China). All isolates were streaked on xylose lysine deoxycholate (XLD) agar plates at 37 °C for 16–24 h, and then colonies were inoculated in Luria–Bertani (LB) medium. Cultures were grown at 37 °C and 180 rpm to the stationary phase for subsequent experiments.

Table 1. Fifty-three isolates of *P. mirabilis* from fox, mink and raccoon farms used in this study.

Isolates	Year of Isolation	Source	
		Animal	Farm Environment ¹
F1-F22	2019	12 isolates from fox feces	9 isolates from soil samples 1 isolate from feed sample
R1-R18	2019	15 isolates from raccoon feces 6 isolates from mink feces,	3 isolates from soil samples
M1-M13	2020	2 isolates from carcass samples, 3 isolates from throat samples, 1 isolate from annal sample	1 isolate from feed sample

¹ The soil samples were collected from the soil within 1 m around the animal cage, and the feed samples were collected from leftover animal feed in the feed trough. F1–F22 are from one fox farm, R1–R18 are from one raccoon farm and M1–M13 are from two mink farms in China.

2.2. Antimicrobial Susceptibility Testing

The CLSI disk diffusion method was used in this study to examine antibiotic resistance of *P. mirabilis* isolates to 11 antibiotics from 7 classes, including cefepime (FEP, 30 µg), cefotaxime (CTX, 30 µg), ceftazidime (CAZ, 30 µg), gentamicin (GM, 10 µg), imipenem (IPM, 10 µg), ampicillin (AMP, 10 µg), streptomycin (STR, 10 µg), ofloxacin (OFX, 5 µg), enrofloxacin (ENR, 15 µg), and trimethoprim-sulfamethoxazole (SXT, 25 µg). In brief, 0.5 McFarland *P. mirabilis* inoculum from liquid culture was spread onto Mueller–Hinton agar plates (Hopebiol), and the antibiotic discs were dispensed on the agar. Plates were incubated at 35 °C air incubator for 16 to 18 h. *E. coli* (ATCC 25922) was used as a quality control strain. The results were interpreted based on the Clinical and Laboratory Standards Institute (CLSI) M100-S27 guideline and CLSI Supplement VET01S [25,26]. The broth microdilution method was used for the determination of the minimum inhibitory concentrations (MICs) of *P. mirabilis* isolates to florfenicol (FFC). *P. mirabilis* isolates resistant to one or more antibiotics in three or more antibiotic classes were defined as multidrug-resistant (MDR).

2.3. Detection of Antimicrobial Resistance Genes

A total of 31 antimicrobial resistance genes in 5 categories, beta-lactam resistance genes (*bla*SHV, *bla*OXA-1, *bla*OXA-23, *bla*OXA-24, *bla*OXA-58, *bla*OXA-48, *bla*CTX-M, *bla*TEM, *bla*PSE, *bla*KPC, *bla*NDM, *bla*IMP, *bla*VIM) [27–31], aminoglycoside resistance genes (*aac*C1, *aac*C2, *aac*C3, *aad*A, *aad*B, *aph*A6, *aac* (6′)-Ib- cr) [32–35], quinolone resistance genes (*qnr*A, *qnr*B, *qnr*C, *qnr*S, *oqx*A, *aac* (6′)-Ib- cr) [27,35], folate pathway antagonist resistance genes (*sul*1, *sul*2, *sul*3) and phenicol resistance genes (*flo*R, *cml*A) were identified by PCR with the primers in Table A1 as previously described [27,36–38]. Bacterial DNA samples were prepared by the boiling method with a modification [39]. Briefly, approximately 10⁸ CFU/mL cells in ddH₂O were boiled at 100 °C for 10 min in a water bath, then cells were centrifuged for five minutes at 1000 rpm and the supernatants were used for the PCR directly. The PCR was carried out in a 25 µL reaction mixture containing 12.5 µL of 2 × Taq Master Mix (Vazyme, China), 1 µL of each primer (10 pmol), 9.5 µL of ddH₂O and 1 µL of DNA template using a T100TM Thermal Cycler (BIO-RAD, USA). ddH₂O was used as a

negative control. PCR products were analyzed by 1% agarose gel electrophoresis at 100 V for 20 min and imaged.

2.4. Detection of Virulence Genes

The virulence genes *mrpA*, *pmfA*, *atfA*, *atfC* (fimbriae), *ureC* (urease), *zapA* (protease), *ucaA* (adhesin), *rsbA* (migration), *rsmA* (repressor of secondary metabolites) and *hmpA* (hemolysin) of *P. mirabilis* were identified with the previous primers and described PCR reaction system (Table A2) [11,40], the bacterial DNA and PCR reaction mixture were prepared as described above.

2.5. Infection of *Galleria Mellonella* Larvae

In this study, the larvae of the *Galleria mellonella* (*G. mellonella*) wax moth was used to assess the pathogenicity of *P. mirabilis* as described previously with modifications [41]. Stationary bacterial cells were washed with PBS and then diluted in PBS to an optical density at OD₆₀₀ of 1, which corresponds to approximately 1×10^9 CFU/mL. For 50% lethal dose (LD₅₀) experiments, a series of 10-fold serial dilutions containing 10^4 to 10^8 CFU in PBS were injected into *G. mellonella* larvae. After surface disinfection using 70% ethanol, a Microliter Syringe (Shanghai Gaoge, China) was used to inject 10 µL aliquots of the inoculum into the hemocoel of each larvae via the last left proleg [42]. A group of 10 larvae were injected with 10 µL of PBS in parallel to ensure that death was not due to injection trauma. Larvae were placed in 9.2 cm Petri dishes and kept at 37 °C in the dark. Insects were considered dead when they displayed no movement in response to touch. The number of dead larvae was scored at 6, 12, and 24 h post-infection. For each strain, at least three independent experiments were performed, and LD_{50s} were calculated according to the formula of Käber [43].

2.6. Determination of In Vivo Bacterial Loads

Each group containing 15 larvae was infected with approximately 1×10^3 CFU per larva of *P. mirabilis* isolates (F13, R8 and M1) or *S. Enteritidis* CVCC3377, respectively. Three insects in each group were collected at different post-infection time points (6 h and death time within 12 h) to determine bacterial loads. After surface disinfection with 70% ethanol, larvae were homogenized in 2 mL of PBS by use of a High-Speed KZ-II Basic Homogenizer (Servicebio, China). Serial dilutions of the homogenates in PBS were plated on XLD agar, and colonies were counted after incubation at 37 °C for 24 h. Three independent experiments were performed.

2.7. Phylogenetic Tree Analysis

The 16S ribosomal RNA (16S rRNA) gene sequences of 53 *P. mirabilis* isolates were amplified with primers 27F and 1492R [44]. The PCR products were sequenced and identified using BLAST algorithms against the NCBI databases. A total of 8 representative *P. mirabilis* isolates from different isolation sources (chicken, snake, milk, etc.) were selected to construct a phylogenetic tree. Multiple alignments of the 16S rRNA gene sequences were conducted with the ClustalW algorithm, and phylogenetic trees were constructed by MEGA 7.0 software using the Neighbor-Joining method [45].

2.8. Data Availability

The partial 16S rRNA gene sequences of 53 *P. mirabilis* isolates were all deposited in the NCBI database under the GenBank accession numbers OL629182–OL629203 (F1–F22), OL629209–OL629226 (R1–R18) and OL629229–OL629241 (M1–M13).

3. Results

3.1. Antimicrobial Susceptibility of *P. mirabilis* Isolates

As shown in Table 2, all isolates were sensitive to cefepime (53/53, 100%), and most isolates were sensitive to ceftazidime (48/53, 90.57%) and ofloxacin (44/53, 83.02%). Most isolates were resistant to florfenicol (48/53, 90.57%), trimethoprim-sulfamethoxazole (39/53, 73.58%), imipenem (38/53, 71.70%), enrofloxacin (34/53, 64.15%), ampicillin (33/53, 63.46%), streptomycin (32/53, 60.38%), gentamicin (27/53, 50.94%), and cefotaxime (26/53, 49.06%). We found that the proportion of MDR isolates was 73.58% (39/53). Among the 39 MDR *P. mirabilis* isolates, isolates resistant to seven classes of antibiotics accounted for the highest proportion (33.33%, 13/39) (Figure 1), of which nine were from raccoons and four were from foxes.

Table 2. The sensitivity of 53 *P. mirabilis* isolates isolated from fox, raccoon and mink farms to 11 antibiotics.

Antibiotic Classes	Antibiotics	Number of Isolates		
		Resistant	Intermediate	Susceptible
Cephems	Cefepime (FEP)	0	1 (1.89%)	52 (98.11%)
	Cefotaxime (CTX)	26 (49.06%)	8 (15.10%)	19 (35.85%)
	Ceftazidime (CAZ)	5 (9.43%)	0	48 (90.57%)
Penicillins	Ampicillin (AMP)	33 (63.46%)	11 (20.75%)	9 (16.98%)
Carbapenems	Imipenem (IPM)	38 (71.70%)	0	15 (28.30%)
Aminoglycosides	Gentamicin (GM)	27 (50.94%)	1 (1.89%)	25 (47.17%)
	Streptomycin (STR)	32 (60.38%)	13 (24.53%)	8 (15.10%)
Quinolones	Ofloxacin (OFX)	9 (16.98%)	8 (15.09%)	36 (67.92%)
	Enrofloxacin (ENR)	34 (64.15%)	14 (26.42%)	5 (9.43%)
Folate pathway antagonists	Trimethoprim-sulfamethoxazole (SXT)	39 (73.58%)	1 (1.89%)	13 (24.53%)
Phenicol	Florfenicol (FFC)	48 (90.57%)	4 (7.55%)	1 (1.89%)

The isolates from the fox farm had generally higher resistance rate than the isolates from raccoon and mink farms (Figure 2). Among the *P. mirabilis* isolates from raccoons and foxes, the significantly different resistance rates to antibiotics were as follows, respectively: to CTX, 40.9% and 94.4% ($p < 0.001$); to AMP, 54.5% and 88.9% ($p < 0.05$); to GM, 40.9% and 83.3% ($p < 0.01$). Meanwhile, by comparing the antibiotic resistance rates between the isolates from minks and foxes, the difference was mainly in AMP ($p < 0.01$), IMP ($p < 0.01$), GM ($p < 0.001$) and STR ($p < 0.01$). The resistant rates of *P. mirabilis* isolates from raccoons and foxes to CTX, CAZ and OFX were 9.1%~94%, but no isolate from mink showed resistance to the above three antibiotics. No difference was found in ENR, SXT and FFC resistance among the isolates from the above three kinds of animals.




Figure 1. The proportion of MDR *P. mirabilis* isolates to different classes of antibiotics used in this study.




Figure 2. Comparison of the antibiotic resistance rates of *P. mirabilis* isolates between raccoons, foxes and minks. The difference was analyzed by chi-squared test. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

3.2. Antimicrobial Resistance Genes of *P. mirabilis*

The most frequently detected antimicrobial resistance gene was *sul1* (94.34%, 50/53), followed by *floR*, *bla_{TEM}*, *aac(6')-Ib-cr* and *bla_{OXA-1}*, with the detection rate of 88.68% (47/53), 83.02% (44/53), 71.70% (38/53) and 60.38% (32/53), respectively (Table 3). Among the 14 beta-lactamase genes detected, *bla_{TEM}* was the most prevalent beta-lactamase gene (83.02%), followed by *bla_{OXA-1}* (60.38%), *bla_{CTX-M}* (32.08%), *bla_{PSE}* (24.53%), *bla_{OXA-24}* (15.09%), and *bla_{NDM}* (13.21%). Among the 51 *P. mirabilis* isolates that were resistant to beta-lactam antibiotics, all isolates carried at least one related drug resistance gene, of which most isolates (23.5%) carried only the *bla_{TEM}* gene, followed by isolates harboring *bla_{TEM}*, *bla_{CTX-M}* and *bla_{OXA-1}* (19.6%) and isolates harboring both *bla_{TEM}* and *bla_{OXA-1}* (9.8%) (Figure 3A). For quinolone resistance genes, the detection rate of *aac(6')-Ib-cr*, *qnrA*, *qnrB* and *qnrC* was 71.70%, 5.66%, 22.64% and 15.09%, respectively, but *qnrS* and *oqxA* were not detected. Most isolates carry the folate pathway antagonist resistance gene *sul1* (94.34%), followed by *sul3* (54.72%) and *sul2* (11.32%), respectively. Among phenicol resistance genes, the detection rate of *floR* was up to 88.68%, followed by *cmlA* (5.66%). In addition, four aminoglycoside resistance genes (*aacC2*, *aadA*, *aadB* and *aphA6*) were detected, accounting for 9.43%, 83.02%, 13.23% and 11.32%, respectively.

By comparing the antimicrobial resistance genes from fox, raccoon and mink farms, we found that gene *bla_{TEM}*, *bla_{OXA-1}*, *aac(6')-Ib-cr*, *sul1*, and *floR* were common. However, *bla_{CTX-M}*, *qnrA* and *qnrC* were only detected in *P. mirabilis* isolates from raccoons and foxes. In addition, the genes *bla_{PSE}*, *qnrB*, *aacC2*, *sul2* and *cmlA* were specific for the isolates from minks. These results show the diversity of antibiotic resistance genes among different farms.

We also compared the detection rate of antibiotic resistance genes between samples from animals (feces, throat, anal and carcass) and the farm environment (feed and soil), and found that the detection rates of antibiotic resistance genes were generally similar (Figure 3B). The proportions of *bla_{TEM}*-positive, *sul1*-positive and *floR*-positive isolates from animal samples and farm environment samples were all above 85%. Although *bla_{TEM}*, *bla_{OXA-1}*, *bla_{CTX-M}*, *bla_{OXA-24}*, *bla_{NDM}*, *aac(6')-Ib-cr*, *qnrA*, *qnrC*, *sul1*, *sul3*, *aadA*, *aadB*, *aphA6* and *floR* genes were prevalent in both animal samples and farm environment samples, no significant difference ($p > 0.05$) was found between the positive rates of the two groups of samples.

Table 3. Detection rates of drug resistance genes of 53 *P. mirabilis* isolates used in this study.

Name	Gene	Detection Rate			
		Raccoon	Fox	Mink	Total
Beta-lactams	<i>bla</i> _{TEM}	33.96%	41.51%	7.55%	83.02%
	<i>bla</i> _{SHV}	0	0	0	0
	<i>bla</i> _{PSE}	0	0	24.53%	24.53%
	<i>bla</i> _{OXA-1}	32.08%	9.43%	18.87%	60.38%
	<i>bla</i> _{OXA-23}	0	0	0	0
	<i>bla</i> _{OXA-24}	0	3.78%	11.32%	15.09%
	<i>bla</i> _{OXA-58}	0	0	0	0
	<i>bla</i> _{OXA-48}	0	0	0	0
	<i>bla</i> _{CTX-M}	18.87%	13.21%	0	32.08%
	<i>bla</i> _{KPC}	0	0	0	0
	<i>bla</i> _{NDM}	11.32%	1.89%	0	13.21%
	<i>bla</i> _{IMP}	0	0	0	0
Aminoglycosides	<i>bla</i> _{VIM}	0	0	0	0
	<i>aac</i> _{C1}	0	0	0	0
	<i>aac</i> _{C2}	0	0	9.43%	9.43%
	<i>aac</i> _{C3}	0	0	0	0
	<i>aadA</i>	28.30%	32.08%	22.64%	83.02%
	<i>aadB</i>	7.56%	1.89%	3.78%	13.23%
	<i>aphA6</i>	0	5.66%	5.66%	11.32%
Quinolones	<i>aac</i> (6')-Ib-cr	32.08%	15.09%	24.53%	71.70%
	<i>qnrA</i>	3.78%	1.89%	0	5.66%
	<i>qnrB</i>	0	0	22.64%	22.64%
	<i>qnrC</i>	9.43%	5.66%	0	15.09%
	<i>qnrS</i>	0	0	0	0
	<i>oqxA</i>	0	0	0	0
	<i>sul1</i>	33.96%	41.51%	18.87%	94.34%
Folate pathway antagonists	<i>sul2</i>	0	0	11.32%	11.32%
	<i>sul3</i>	32.08%	20.75%	1.89%	54.72%
Phenicol	<i>cmlA</i>	0	0	5.66%	5.66%
	<i>floR</i>	33.96%	41.51%	13.21%	88.68%




Figure 3. (A) Beta-lactam antibiotics resistance genes among the 53 *P. mirabilis* isolates. (B) Comparison of the detection rate of resistance genes in *P. mirabilis* isolates from animal samples and farm environment samples.

3.3. Virulence Genes of *P. mirabilis* Isolates

Among the 10 virulence genes, *ureC*, *zapA*, *pmfA*, *atfA* and *mrpA* genes were all detected among 53 *P. mirabilis* isolates, followed by both *atfC* and *hmpA* genes, both with a detection rate of 98.11%, *rsmA* at 94.34%, *rsbA* at 71.70% and *ucaA* at 45.28% (Table 4).

Table 4. Detection rate of 10 virulence genes of *P. mirabilis* isolates from different farms.

Gene	Fox (n = 22)	Raccoon (n = 18)	Mink (n = 13)	Total (n = 53)
<i>ureC</i>	100%	100%	100%	100%
<i>zapA</i>	100%	100%	100%	100%
<i>pmfA</i>	100%	100%	100%	100%
<i>mrpA</i>	100%	100%	100%	100%
<i>atfC</i>	100%	94.44%	100%	98.11%
<i>atfA</i>	100%	100%	100%	100%
<i>ucaA</i>	45.45%	33.33%	61.54%	45.28%

<i>rsbA</i>	68.18%	61.11%	92.31%	71.70%
<i>rsmA</i>	95.45%	100%	84.62%	94.34%
<i>hmpA</i>	95.45%	100%	100%	98.11%

3.4. Pathogenicity of *P. mirabilis* Isolates

3.4.1. Pathogenicity Comparison of *P. mirabilis* Isolates from Different Animal Sources

To determine the pathogenicity of *P. mirabilis* isolates from different animal sources, three *P. mirabilis* isolates (F13, R8 and M1) that carry all detected 10 virulence genes were selected. The LD₅₀ of *P. mirabilis* F13, R8, and M1 in *G. mellonella* larvae model were 2.5×10^4 CFU/larvae, 2.0×10^4 CFU/larvae and 3.9×10^5 CFU/larvae at 6 h post infection, respectively (Table 5). The results indicate that the virulence of *P. mirabilis* M1 from minks seemed to be weaker than F13 from foxes and R8 from raccoons.

To further determine whether *G. mellonella* mortality is associated with the growth of bacteria in infected larvae, larvae were injected with 10^3 CFU of *P. mirabilis*. As shown in Figure 4A, infection of *G. mellonella* with the F13, R8, and M1 strain resulted in a 10^4 – 10^5 -fold increase in CFU at 6 h post infection, which was followed by further increases in the bacterial numbers of dead larvae within 12 h post infection. These results are similar to that of larvae infected with pathogenic *S. Enteritidis* CVCC3377, which caused a remarkable increase in bacterial numbers in *G. mellonella* over time. At 12 h post infection, larvae in *P. mirabilis* infection group were all dead, whereas 60% larvae survived in the *S. Enteritidis* infection group (Figure 4B). The results demonstrate that *P. mirabilis* isolates from fox, mink and raccoon farms could replicate rapidly in vivo and displayed high virulence in *G. mellonella* larvae model.

Table 5. The LD₅₀s of three *P. mirabilis* isolates from different animal species in *G. mellonella* larvae model.

Isolates	Bacteria Dose (CFU/Larvae)	Dead			LD ₅₀ (6 h) (CFU/Larvae)
		6 h	12 h	24 h	
F13	1×10^6	9	10	10	2.5×10^4
	1×10^5	6	10	10	
	1×10^4	6	10	10	
	1×10^3	0	10	10	
	1×10^2	0	10	10	
R8	1×10^6	9	10	10	2.0×10^4
	1×10^5	10	10	10	
	1×10^4	3	10	10	
	1×10^3	0	10	10	
	1×10^2	0	10	10	
M1	1×10^6	9	10	10	3.9×10^5
	1×10^5	0	10	10	
	1×10^4	0	10	10	
	1×10^3	0	9	10	
	1×10^2	0	6	10	




Figure 4. Comparison of pathogenicity between three *P. mirabilis* isolates from different animal sources and *S. Enteritidis*. **(A)** The bacterial loads in larvae infected with *P. mirabilis* and *S. Enteritidis*. **(B)** Survival rate in larvae infected with *P. mirabilis* and *S. Enteritidis* at 6 and 12 h post infection. Each group was infected with approximately 1×10^3 CFU per larva of *P. mirabilis* isolates (F13, R8 and M1) or *S. Enteritidis* CVCC3377, respectively. Three infected larvae from each group were pooled and homogenized, and the numbers of CFU were determined by plating. The survival rates in each group were calculated at 6 and 12 h post-infection. $n = 15$.

3.4.2. Pathogenicity of *P. mirabilis* Isolates with Different Virulence Genes

To evaluate the effect of virulence genes on the pathogenicity of *P. mirabilis* isolates, three isolates, which were all isolated from fox feces but with different virulence genes, were selected to compare their virulence in a *G. mellonella* model. As shown in Table 6, the LD₅₀ of F2, F3, and F6 isolates at 6 h post infection was 2.0×10^4 CFU/larvae, 7.9×10^3 CFU/larvae and 4.0×10^3 CFU/larvae, respectively. Although strain F2 carried all detected virulence genes, its LD₅₀ was 2.53- and 5-fold higher than that of strain F3 lacking *ucaA*, *rsbA* and *rsmA*, and strain F6 lacking *ucaA*. The results indicate that the carriage rate of virulence-associated genes was not positively correlated with *P. mirabilis* pathogenicity in *G. mellonella* larvae.

Table 6. Pathogenicity of *P. mirabilis* isolates with different virulence genes from foxes in *G. mellonella* larvae.

Tested Strain	Virulence Gene			Bacteria Dose (CFU/Larvae)	Dead			LD ₅₀ (6 h) (CFU/Larvae)
	<i>ucaA</i>	<i>rsbA</i>	<i>rsmA</i>		6 h	12 h	24 h	
F2	+	+	+	1×10^6	10	10	10	2.0×10^4
				1×10^5	7	10	10	
				1×10^4	5	10	10	
				1×10^3	0	1	3	
				1×10^2	0	0	2	
F3	-	-	-	1×10^6	10	10	10	7.9×10^3
				1×10^5	9	10	10	
				1×10^4	7	10	10	
				1×10^3	0	10	10	
				1×10^2	0	8	10	
F6	-	+	+	1×10^6	10	10	10	4.0×10^3
				1×10^5	10	10	10	
				1×10^4	9	10	10	
				1×10^3	0	10	10	
				1×10^2	0	10	10	



Figure 5. Phylogenetic tree constructed based on 16S rRNA sequences of 61 *P. mirabilis* isolates using the neighbor-joining method. Eight representative isolates from different sources are labeled with big parentheses, 14 isolates from farms environment in the present study are marked in turquoise, and isolates harboring *bla*_{NDM} or *bla*_{OXA-24} are marked in green or yellow, respectively. All sequences were aligned using ClustalW, then the aligned data were evaluated by the neighbor-joining approach using MEGA7 software with 2000 replications of bootstrap. Branches corresponding to partitions reproduced in less than 50% bootstrap replicates are collapsed. There were a total of 1337 positions in the final dataset.

3.5. Phylogenetic Analysis of *P. mirabilis* Isolates

The 53 *P. mirabilis* isolates and the selected eight representative isolates of *P. mirabilis* 16S rRNA gene sequences obtained on NCBI were used to perform phylogenetic analysis to understand the evolutionary relationship among the *P. mirabilis* isolates (Figure 5). The results show that the sequence similarity between the 53 isolates and eight representative bacteria was 99.5–99.9%, but the *P. mirabilis* isolates from fur farms have far lower relatedness with the representative *P. mirabilis* isolates from shrimp, chick, mastitic milk, etc. The 22 *P. mirabilis* isolates from foxes and 18 isolates from raccoons were co-clustered into three branches, but the 13 *P. mirabilis* isolates from minks formed many clades in the evolutionary phylogeny, indicating there may be multiple origins of *P. mirabilis* isolates from minks. Moreover, the isolates from animal samples and farm environment samples were distributed across different branches. This pattern suggested that the isolates from farm environment samples had close relatedness with those from animal samples.

4. Discussion

Bacterial antibiotic resistance is a global public health threat, and one important cause of it is the multidrug bacterial isolates of animal origins acting as an important source for human infections [46–48]. We investigated the characteristics of 53 *P. mirabilis* isolates from fox, raccoon and mink farms—the results show that the proportion of MDR isolates among all the *P. mirabilis* isolates was up to 73.58%, similar to that among the 32 *P. mirabilis* isolates from chicken carcasses in a poultry slaughterhouse in Brazil (78.13%) [47]. ESBL production is significantly associated with mortality in patients with bacteremia caused by *P. mirabilis* [49]. In this study, 51 of the 53 *P. mirabilis* isolates were resistant to beta-lactam antibiotics and most isolates exhibited two or more beta-lactamase genes. The most common ESBL-encoding gene among the 53 *P. mirabilis* isolates from foxes, raccoons and minks is *bla_{TEM}* (83.02%), which is spread worldwide and is now found in many different species of the order of *Enterobacterales* [17]. Carbapenem resistance rates are typically extremely low in clinical *P. mirabilis* isolates; however, approximately 71.70% of isolates from foxes, raccoons and minks were resistant to imipenem. Among the nine detected carbapenemase genes, the class D carbapenemase gene *bla_{OXA-24}* (15.09%) and the class B metallo-beta-lactamase *bla_{NDM}* (13.21%) were found for the first time in *P. mirabilis* isolates from foxes, raccoons and minks. Kang et al. found that 16.67% (6/54) of the *P. mirabilis* isolates from wild animals were resistant to meropenem, and they mainly carried the carbapenem-resistance genes *bla_{OXA-1}* and *bla_{NDM-1}* [19]. For the 35 *P. mirabilis* isolates obtained from 240 duck samples in Egypt, only three strains were carbapenem-resistant, two strains harbored the *bla_{NDM-1}* gene, and one strain carried the *bla_{KPC}* gene [50]. The presence of *bla_{NDM-1}* in *Proteus* spp. clinical isolates is still episodic in China and many other countries [17]. However, the *bla_{NDM-1}* gene had been identified on a genomic island in *P. mirabilis* recovered from the urine of a hospitalized patient in France in 2012 [51]. The *bla_{OXA-24}* gene, mainly detected in *Acinetobacter baumannii*, were shown to appear in reservoirs including livestock, companion animals, and the environment [17,52]. However, the *bla_{OXA-24}* gene is only detected in clinical strains of *P. mirabilis* in the United States [22]. To the best of our knowledge, *P. mirabilis* isolates harboring *bla_{OXA-24}* were identified for the first time from animal origins. Some other mechanisms including through porin mutation with or without decreased expression of penicillin binding proteins may contribute to the high imipenem-resistance rates of *P. mirabilis* isolates in this study. Moreover, the prevalence of beta-lactam-resistant *P. mirabilis* isolates from fur animals in this study and the similar antibiotic resistance profiling between *P. mirabilis* isolates from fur animals and farm environment suggested that MDR *P. mirabilis* isolates may likely spread from fur animals to the environment and potentially humans, posing a public health threat [53].

We also found increasing prevalence and diversity of some other antibiotic resistance genes in *P. mirabilis* from foxes, raccoons and minks. It was reported that *aac(6′)-Ib-cr*

conferring quinolones and aminoglycosides resistance was common in *Escherichia coli* [54]. Here we showed that the detection rate of *aac(6′)-Ib-cr* among 53 *P. mirabilis* isolates was up to 71.7%, similar to the 63.2% among 19 *P. mirabilis* isolates from a Chinese hospital by Hu et al. [55]. The detection rates of quinolone resistance genes *qnrA* (5.66%) and *qnrB* (22.64%) were higher than the 0% from chicken carcasses in Brazil by Sanches et al. [19]. For the aminoglycoside resistance of *P. mirabilis* isolates in this study, another frequently encountered resistance gene was *aadA*, which may result in high resistance rates to streptomycin in these *P. mirabilis* isolates. These results indicate that *aac(6′)-Ib-cr* and *aadA* gene might be common in *P. mirabilis* isolates in foxes, raccoons and minks of China. Florfenicol is mainly used in veterinary medicine, a key gene for florfenicol and chloramphenicol resistance, *floR*, coding for an efflux protein of 12 transmembrane segments, can spread among bacteria of the same and different species or genera through horizontal gene transfer [56]. In this study, we found that 90.57% of *P. mirabilis* isolates were resistant to florfenicol, of which 77.36% carried the *floR* gene. So, the emergence and dissemination of florfenicol resistance among *P. mirabilis* isolates will limit the use of this antimicrobial for treating bacterial infections.

All *P. mirabilis* isolates harbored *ureC*, *zapA*, *pmfA*, *atfA* and *mrpA* virulence genes, which were also prevalent in the *P. mirabilis* isolates from chicken carcasses [47] and pork meat [24]. Among the 10 virulence genes, the detection rate of *ucaA*, a major fimbrial subunit that can enhance the adhesion of *P. mirabilis*, was the lowest [57]. However, we showed that the virulence of a *ucaA*-positive strain was about 5-fold lower than that of a *ucaA*-negative strain in *G. mellonella* larvae. Moreover, the carriage rate of virulence-associated genes in these *P. mirabilis* isolates did not correlate with higher pathogenicity in a *G. mellonella* model. Three *P. mirabilis* isolates carrying all detected 10 virulence genes from different animal species showed different lethal abilities in *G. mellonella* larvae. These *P. mirabilis* isolates could multiply rapidly in the hemolymph of the larvae, similarly to *S. Enteritidis*, resulting in a fierce but ineffective inflammatory response triggered by high bacterial burdens and hemocyte depletion [58]. These results suggest that the carriage rate of virulence genes in *P. mirabilis* isolates can only reflect their epidemic, the pathogenicity of bacteria should be evaluated by appropriate animal models.

5. Conclusions

In the present study, we firstly reported the emergence of carbapenem-resistant *P. mirabilis* isolates harboring *bla_{OXA-24}* (15.09%) or *bla_{NDM}* (13.21%) from foxes, raccoons and minks. All *P. mirabilis* isolates harbored *ureC*, *zapA*, *pmfA*, *atfA* and *mrpA* genes, but the carriage rate of these virulence-associated genes in the isolates did not correlate with higher pathogenicity in a *G. mellonella* model. Moreover, phylogenetic analysis showed that the *P. mirabilis* isolates from farm environment samples may have close relatedness with that from animals.

Supplementary Materials: The following are available online at www.mdpi.com/article/10.3390/biology11020292/s1, Table S1. Characteristics of virulence genes of 53 *P. mirabilis* isolates from fur animal farms in this study. Table S2. Characteristics of antibiotic resistance genes of 53 *P. mirabilis* isolates from fur animal farms in this study.

Author Contributions: Conceptualization, G.H. G.W. and S.S.; methodology, G.H. and P.L.; investigation, P.L., Y.C. and L.C.; data curation, Y.C.; writing-original draft, P.L.; writing-review and editing, P.L., S.S. and G.H. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Shandong Agricultural Major Application Technology Innovation Project (SD2019XM009) and Shandong Provincial Modern Agricultural Industry Technology System (SDAIT-21-05).

Institutional Review Board Statement: The animal study was reviewed and approved by Animal Care and Use of Shandong Agricultural University (SDAUA-2018-027). Written informed consent was obtained from the owners for the participation of their animals in this study.

Informed Consent Statement: .Not applicable.

Data Availability Statement: The original contributions generated for this study are included in the article, further inquiries can be directed to the corresponding authors.

Acknowledgments: We thank Weixiang Fan for technical support and Yitian Zhou from University of Pennsylvania for English language editing. Also, we thank the anonymous referee for helpful comments and suggestions.

Conflicts of Interest: No conflict of interest exists in the submission of this manuscript, and manuscript is approved by all authors for publication.

Appendix A

Table A1. Primer sequence and fragment size of drug resistance genes.

Name	Gene	Sequence (5'-3')	Size (bp)	Reference
Beta-lactams	<i>bla_{TEM}</i>	F: ATAAAATTCTTGAAGACGAAA R: GACAGTTACCAATGCTTAATC	643	[27]
	<i>bla_{SHV}</i>	F: TTATCTCCCTGTTAGCCACC R: GATTGCTGATTTCGCTCGG	860	[27]
	<i>bla_{PSE}</i>	F: TAGGTGTTTCCGTTCTTG R: TCATTTCGCTCTTCCATT	150	[28]
	<i>bla_{OXA-1}</i>	F: TCAACTTTCAAGATCGCA R: GTGTGTTTAGAATGGTGA	591	[27]
	<i>bla_{OXA-23}</i>	F: GATCGGATTGGAGAACCAGA R: ATTTCTGACCGCATTTCCAT	501	[30]
	<i>bla_{OXA-24}</i>	F: TTCCCCTAACATGAATTTGT R: GTACTAATCAAAGTTGTGAA	1024	[30]
	<i>bla_{OXA-58}</i>	R: TGGATTGCACCTTCATCTTGG F: TGGCACGCATTTAGACCG R: AAACCCACATACCAACCC	507	[30]
	<i>bla_{OXA-48}</i>	F: GCGTGGTTAAGGATGAACAC R: CATCAAGTTCAACCCAACCG	438	[29]
	<i>bla_{CTX-M}</i>	F: CGCTTTGCGATGTGCAG R: ACCGCGATATCGTTGGT	550	[27]
	<i>bla_{KPC}</i>	F: GACGGAAAGCTTACAAAACTGACA R: CTTGTCATCCTTGTTAGGCG	259	[31]
	<i>bla_{NDM}</i>	F: GGTTTGGCGATCTGGTTTTC R: ATCCAGTTGAGGATCTGGGC	181	[31]
	<i>bla_{IMP}</i>	F: GGAATAGAGTGGCTTAATTCTC R: CAAGCTTCTATATTTGCGTCACC	275	[31]
	<i>bla_{VIM}</i>	F: GATGAGTTGCTTTTGATTGATACAGC R: CGCCCGAAGGACATCAA	153	[31]
Quinolones	<i>qnrA</i>	F: ATTTCTCACGCCAGGATTTG R: GATCGGCAAAGGTCAGGTCA	519	[27]
	<i>qnrB</i>	F: GATCGTGAAAGCCAGAAAGG R: ACGATGCCTGGTAGTTGTCC	513	[27]
	<i>qnrC</i>	F: GGTTGTACATTTATTGAATC R: TCCACTTTACGAGTTCT	666	[27]
	<i>qnrS</i>	F: ACGACATTCGTCAACTGCAA R: TAAATTGGCACCCCTGTAGGC	417	[27]

Aminoglycosides	<i>oqxA</i>	F: GATCAGTCAGTGGGATAGTTT R: TACTCGGCGTTAACTGATTA	670	[35]
	<i>aac(6′)-Ib-cr</i>	F: TTGCGATGCTCTATGAGTGGCTA R: CTCGAATGCCTGGCGTGTTT	482	[27]
	<i>aaC1</i>	F: ACCTACTCCCAACATCAGCC R: ATATAGATCTCACTACGCGC	528	[32]
	<i>aaC2</i>	F: ACTGTGATGGGATACGCGTC R: CTCCGTCAGCGTTTCAGCTA	482	[32]
	<i>aaC3</i>	F: CACAAGAACGTGGTCCGCTA R: AACAGGTAAGCATCCGCATC	185	[32]
	<i>aadA</i>	F: GTGGATGGCGGCCTGAAGCC R: AATGCCCAGTCGGCAGCG	535	[34]
	<i>aadB</i>	F: ATGGACACAACGCAGGTC R: TTAGGCCGCATATCGCGACC	495	[33]
	<i>aphA6</i>	F: ATGGAATTGCCCAATATTATTC R: TCAATTCAATTCATCAAGTTTTA	399	[33]
	<i>sul1</i>	F: CTTGATGAGAGCCGGCGG C R: GCAAGGCGGAAACCCGCGCC	238	[37]
	<i>sul2</i>	F: CGGCATCGTCAACATAAC C R: GTGTGCGGATGAAGTCAG	722	[36]
Folate pathway antagonists	<i>sul3</i>	F: AGATGTGATTGATTGGGAGC R: TAGTTGTTTCTGGATTAGAGCCT	443	[38]
	<i>cmlA</i>	F: TGTCAATTTACGGCATACTCG R: ATCAGGCATCCCATCCCAT	900	[37]
	<i>floR</i>	F: CACGTTGAGCCTCTATATGG R: ATGCAGAAGTAGAACGCGAC	890	[27]

Table A2. Primers and annealing temperatures of 10 virulence genes.

Gene	Sequence (5′→3′)	Fragment	Annealing Temperature
<i>ureC</i>	F: GTTATTCGTGATGGGATGGG R: ATAAAGGTGGTTACGCCAG	375 bp	52 °C
<i>mrpA</i>	F: ATTTCAAGGAAACAAAAGATG R: TTCTTACTGATAAGACATTG	410 bp	39 °C
<i>zapA</i>	F: ACCGCAGGAAAACATATAGCCC R: GCGACTATCTTCCGCATAATCA	493 bp	52 °C
<i>atfA</i>	F: CATAATTTCTAGACCTGCCCTAGCA R: CTGCTTGGATCCGTAATTTTAAACG	365 bp	49 °C
<i>atfC</i>	F: AGAAAGGGATCCTACAAATTAA R: TATAGCATGCATTTAAATTGCC	472 bp	49 °C
<i>ucaA</i>	F: GTAAAGTTGTTGCGCAAAC R: TTGAGCCACTGTGGATACA	365 bp	49 °C
<i>pmfA</i>	F: GGATCATCTATAATGAAACTG R: CTGATAATCAACTTGGAAAGTT	534 bp	52 °C
<i>rsbA</i>	F: TCGATTTCAAGTGTGTTGGCCAT R: TCGATTTCAAGTGTGTTGGCCAT	1647 bp	55 °C
<i>rsmA</i>	F: TAGCGAGTGTTGACGAGTGG R: AGCGAGGTGAAGAACGAGAA	562 bp	56 °C
<i>hmpA</i>	F: CCAGTGAATTAACGGCAGGT R: CGTGCCCAAGTAATGGCTAAT	654 bp	55 °C

References

- Drzewiecka, D. Significance and roles of *Proteus* spp. bacteria in natural environments. *Microbial Ecology* **2016**, *72*, 741–758.
- Jacobsen, S.; Stickler, D.; Mobley, H.; Shirliff, M. Complicated catheter-associated urinary tract infections due to *Escherichia coli* and *Proteus mirabilis*. *Clinical microbiology reviews* **2008**, *21*, 26–59.
- Jain, S.; Gaiind, R.; Kothari, C.; Sehgal, R.; Shamweel, A.; Thukral, S.S.; Chellani, H.K. VEB-1 extended-spectrum β -lactamase-producing multidrug-resistant *Proteus mirabilis* sepsis outbreak in a neonatal intensive care unit in India: clinical and diagnostic implications. *JMM case reports* **2016**, *3*, e005056, doi:10.1099/jmmcr.0.005056.
- Cao, B.; Wang, M.; Liu, L.; Zhou, Z.; Wen, S.; Rozalski, A.; Wang, L. 16S-23S rDNA internal transcribed spacer regions in four *Proteus* species. *Journal of microbiological methods* **2009**, *77*, 109–118, doi:10.1016/j.mimet.2009.01.024.
- Wang, Y.; Zhang, S.; Yu, J.; Zhang, H.; Yuan, Z.; Sun, Y.; Zhang, L.; Zhu, Y.; Song, H. An outbreak of *Proteus mirabilis* food poisoning associated with eating stewed pork balls in brown sauce, Beijing. *Food Control* **2009**, *21*, 302305.
- Gong, Z.; Shi, X.; Bai, F.; He, X.; Zhang, H.; Li, Y.; Yu, W.; Lin, Y.; Qiu, Y.; Chen, Q.; et al. Characterization of a novel diarrheagenic strain of *Proteus mirabilis* associated with food poisoning in China. *Frontiers in Microbiology* **2019**, *2810*.
- Shi, X.; Lin, Y.; Qiu, Y.; Li, Y.; Jiang, M.; Chen, Q.; Jiang, Y.; Yuan, J.; Cao, H.; Hu, Q.; et al. Comparative screening of digestion tract toxic genes in *Proteus mirabilis*. *PLOS ONE* **2016**, *11*, e0151873.
- Pattanayak, S.; Kumar, P.R.; Sahoo, M.K.; Paul, A.; Sahoo, P.K. First field-based evidence of association of *Proteus mirabilis* causing large scale mortality in Indian major carp farming. *Aquaculture* **2018**, *495*, 435–442.
- Cabassi, C.S.; Taddei, S.; Predari, G.; Galvani, G.; Ghidini, F.; Schiano, E.; Cavarani, S. Bacteriologic findings in ostrich (*struthio camelus*) eggs from farms with reproductive failures. *Avian Diseases* **2004**, *48*, 716–722.
- Milagro, F.-D.; Monica, C.; Alexandra, G.-A.M.; Pulchérie, G.; Paula, S. Occurrence of *Proteus mirabilis* associated with two species of venezuelan oysters. *Revista do Instituto de Medicina Tropical de Sao Paulo* **2007**, *49*, 355–359.
- Sun, Y.; Wen, S.; Zhao, L.; Xia, Q.; Wang, H. Association among biofilm formation, virulence gene expression, and antibiotic resistance in *Proteus mirabilis* isolates from diarrhetic animals in Northeast China. *BMC Veterinary Research* **2020**, *16*.
- Malayeri, H.Z.; Jamshidi, S.; Salehi, T.Z. Identification and antimicrobial susceptibility patterns of bacteria causing otitis externa in dogs. *Veterinary Research Communications* **2010**, *34*, 435–444.
- Papp-Wallace, K.M.; Endimiani, A.; Taracila, M.A.; Bonomo, R.A. Carbapenems: past, present, and future. *Antimicrob Agents Chemother* **2011**, *55*, 4943–4960, doi:10.1128/AAC.00296-11.
- Ghafeurian, S.; Sadeghifard, N.; Soheili, S.; Sekawi, Z. Extended spectrum beta-lactamases: definition, classification and epidemiology. *Curr Issues Mol Biol* **2015**, *17*, 11–21.
- Tibbetts, R.; Frye, J.G.; Marschall, J.; Warren, D.; Dunne, W. Detection of KPC-2 in a clinical isolate of *Proteus mirabilis* and first reported description of carbapenemase resistance caused by a KPC beta-lactamase in *P. mirabilis*. *J Clin Microbiol* **2008**, *46*, 3080–3083, doi:10.1128/JCM.00979-08.
- Cabral, A.B.; Maciel, M.A.; Barros, J.F.; Antunes, M.M.; Lopes, A.C. Detection of *blaKPC-2* in *Proteus mirabilis* in Brazil. *Rev Soc Bras Med Trop* **2015**, *48*, 94–95, doi:10.1590/0037-8682-0152-2014.
- Girlich, D.; Bonnin, R.A.; Dortet, L.; Naas, T. Genetics of acquired antibiotic resistance genes in *Proteus* spp. *Front Microbiol* **2020**, *11*, 256, doi:10.3389/fmicb.2020.00256.
- Xie, X.; Zhang, J.; Wang, H.N.; Lei, C.W. Whole genome sequence of a New Delhi metallo-beta-lactamase 1-producing *Proteus mirabilis* isolate SNYG35 from broiler chicken in China. *J Glob Antimicrob Resist* **2021**, *24*, 266–269, doi:10.1016/j.jgar.2020.12.023.
- Kang, Q.; Wang, X.; Zhao, J.; Liu, Z.; Ji, F.; Chang, H.; Yang, J.; Hu, S.; Jia, T.; Wang, X.; et al. Multidrug-resistant *Proteus mirabilis* isolates carrying *blaOXA-1* and *blaNDM-1* from wildlife in China: increasing public health risk. *Integrative zoology* **2021**, *16*, 798–809, doi:10.1111/1749-4877.12510.
- Lei, C.W.; Zhang, A.Y.; Liu, B.H.; Wang, H.N.; Yang, L.Q.; Guan, Z.B.; Xu, C.W.; Zhang, D.D.; Yang, Y.Q. Two novel *Salmonella* genomic island 1 variants in *Proteus mirabilis* isolates from swine farms in China. *Antimicrob Agents Chemother* **2015**, *59*, 4336–4338, doi:10.1128/AAC.00120-15.
- Algammal, A.M.; Hashem, H.R.; Alfifi, K.J.; Hetta, H.F.; Sheraba, N.S.; Ramadan, H.; El-Tarabili, R.M. *atpD* gene sequencing, multidrug resistance traits, virulence-determinants, and antimicrobial resistance genes of emerging XDR and MDR-*Proteus mirabilis*. *Sci Rep* **2021**, *11*, 9476, doi:10.1038/s41598-021-88861-w.
- Leulmi, Z.; Kandouli, C.; Mihoubi, I.; Benlabed, K.; Lezzar, A.; Rolain, J.M. First report of *blaOXA-24* carbapenemase gene, *armA* methyltransferase and *aac(6)-Ib-cr* among multidrug-resistant clinical isolates of *Proteus mirabilis* in Algeria. *J Glob Antimicrob Resist* **2019**, *16*, 125–129, doi:10.1016/j.jgar.2018.08.019.
- Chen, L.; Al Laham, N.; Chavda, K.D.; Mediavilla, J.R.; Jacobs, M.R.; Bonomo, R.A.; Kreiswirth, B.N. First report of an OXA-48-producing multidrug-resistant *Proteus mirabilis* strain from Gaza, Palestine. *Antimicrob Agents Chemother* **2015**, *59*, 4305–4307, doi:10.1128/AAC.00565-15.
- Silva, S.M.; Caroline, R.d.S.; Carvalho, S.L.; Hugo, M.V.; Gabriel, L.B.M.; Henrique, M.G.G.; Henrique, D.d.O.B.; Kenji, N.E.; Carla, F.G.L.; Carolina, V.E.; et al. *Proteus mirabilis* from community-acquired urinary tract infections (UTI-CA) shares genetic similarity and virulence factors with isolates from chicken, beef and pork meat. *Microbial Pathogenesis* **2021**, *158*, 105098.
- Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing 27th ed CLSI supplement M100-S27. Clinical and Laboratory Standards Institute: PA, USA, **2017**.
- Clinical Laboratory Standards Institute. Performance Standards for Antimicrobial Disk Susceptibility Tests for Bacteria Isolated from Animals: CLSI Supplement VET01S; Replaces VET01-S2. Clinical and Laboratory Standards Institute: PA, USA, **2015**.

27. Ahmed, A.M.; Yusuke, M.; Maiko, S.; Akito, M.; Hitoshi, W.; Yukio, F.; Tadashi, S. Zoo animals as reservoirs of gram-negative bacteria harboring integrons and antimicrobial resistance genes. *Applied and environmental microbiology* **2007**, *73*, 6686–6690.
28. Puah, S.-M.; Puthuchear, S.D.; Liew, F.-Y.; Chua, K.-H. *Aeromonas aquariorum* clinical isolates: antimicrobial profiles, plasmids and genetic determinants. *International Journal of Antimicrobial Agents* **2013**, *41*, 281–284.
29. Poirel, L.; Walsh, T.R.; Cuvillier, V.; Nordmann, P. Multiplex PCR for detection of acquired carbapenemase genes. *Diagnostic microbiology and infectious disease* **2011**, *70*, 119–123, doi:10.1016/j.diagmicrobio.2010.12.002.
30. Hou, C.; Yang, F. Drug-resistant gene of blaOXA-23, blaOXA-24, blaOXA-51 and blaOXA-58 in *Acinetobacter baumannii*. *International journal of clinical and experimental medicine* **2015**, *8*, 13859–13863.
31. Oliveira, M.; Nunes, M.; Barreto Crespo, M.T.; Silva, A.F. The environmental contribution to the dissemination of carbapenem and (fluoro)quinolone resistance genes by discharged and reused wastewater effluents: The role of cellular and extracellular DNA. *Water research* **2020**, *182*, 116011, doi:10.1016/j.watres.2020.116011.
32. Navajas-Benito, E.V.; Alonso, C.A.; Sanz, S.; Olarte, C.; Martínez-Olarte, R.; Hidalgo-Sanz, S.; Somalo, S.; Torres, C. Molecular characterization of antibiotic resistance in *Escherichia coli* strains from a dairy cattle farm and its surroundings. *Journal of the Science of Food and Agriculture* **2017**, *97*, 362–365.
33. Sarhaddi, N.; Soleimanpour, S.; Farsiani, H.; Mosavat, A.; Dolatabadi, S.; Salimizand, H.; Jamehdar, S.A. Elevated prevalence of multidrug-resistant *Acinetobacter baumannii* with extensive genetic diversity in the largest burn centre of northeast Iran. *Journal of Global Antimicrobial Resistance* **2017**, *8*, 60–66, doi:10.1016/j.jgar.2016.10.009.
34. Nguyen, V.C.; Lijun, Z.; Thi, T.T.V.; Hoang, S.H.P.; Tran, N.T.; Qi, H.; Rui, Z. Association between the phenotypes and genotypes of antimicrobial resistance in *Haemophilus parasuis* isolates from swine in Quang Binh and Thua Thien Hue Provinces, Vietnam. *Engineering* **2020**, *6*, 40–48, doi:10.1016/j.eng.2019.10.014.
35. Ping, L.X.; Jing, X.; Lei, Y.; Liang, L.; Jian, S.; Hong, L.Y.; Xia, J.H. Characterization of CTX-M-14-producing *Escherichia coli* from food-producing animals. *Frontiers in Microbiology* **2015**, 1136.
36. Chu, C.; Chiu, C.; Wu, W.; Chu, C.; Liu, T.; Ou, J. Large drug resistance virulence plasmids of clinical isolates of *Salmonella enterica* serovar Choleraesuis. *Antimicrobial agents and chemotherapy* **2001**, *45*, 2299–2303.
37. Guerra, B.; Soto, S.; Argüelles, J.; Mendoza, M. Multidrug resistance is mediated by large plasmids carrying a class 1 integron in the emergent *Salmonella enterica* serotype [4,5,12:i:-]. *Antimicrobial agents and chemotherapy* **2001**, *45*, 1305–1308.
38. Sheng, C.; Shaohua, Z.; G, W.D.; M, S.C.; Ran, L.; Hanchun, Y.; F, M.P.; Sherry, A.; Jianghong, M. Characterization of multiple-antimicrobial-resistant *salmonella* serovars isolated from retail meats. *Applied and environmental microbiology* **2004**, *70*, 1–7.
39. Dashti, A.A.; Jadaon, M.M.; Abdulsamad, A.M.; Dashti, H.M. Heat treatment of bacteria: a simple method of DNA extraction for molecular techniques. *Kuwait Med J* **2009**, *41*, 117–122.
40. YouQin, Y.; ChunHua, H.; YiLong, C.; Yun, S.; XiaoWei, H.; DeHui, M. Comparative study on pathogenicity in mice and carriage of virulence genes of *Proteus mirabilis* isolates from various origins. *Chinese Journal of Biologicals* **2019**, *32*, 289–294.
41. Hernandez, R.J.; Hesse, E.; Dowling, A.J.; Coyle, N.M.; Feil, E.J.; Gaze, W.H.; Vos, M. Using the wax moth larva *Galleria mellonella* infection model to detect emerging bacterial pathogens. *PeerJ* **2019**, *6*, e6150, doi:10.7717/peerj.6150.
42. Peleg, O.; Eliopoulos, G.; Moellering, R. *Galleria mellonella* as a model system to study *Acinetobacter baumannii* pathogenesis and therapeutics. In Proceedings of the Infectious Diseases Society of America 2008 Annual Meeting, 2008; pp. 2605–2609.
43. Finney, D.J. The median lethal dose and its estimation. *Archives of toxicology* **1985**, *56*, 215–218, doi:10.1007/bf00295156.
44. Yang, B.; Wang, Y.; Qian, P.Y. Sensitivity and correlation of hypervariable regions in 16S rRNA genes in phylogenetic analysis. *BMC bioinformatics* **2016**, *17*, 135, doi:10.1186/s12859-016-0992-y.
45. Kumar, S.; Stecher, G.; Tamura, K. MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Molecular biology and evolution* **2016**, *33*, 1870–1874, doi:10.1093/molbev/msw054.
46. Van, B.T.P.; Charles, B.; Marius, G.; Grenfell, B.T.; A, L.S.; P, R.T.; Aude, T.; Ramanan, L. Global trends in antimicrobial use in food animals. *Proceedings of the national academy of sciences of the United States of America* **2015**, *112*, 5649–5654.
47. Sanches, M.S.; Baptista, A.A.S.; Souza, M.; Menck-Costa, M.F.; Koga, V.L.; Kobayashi, R.K.T.; Rocha, S.P.D. Genotypic and phenotypic profiles of virulence factors and antimicrobial resistance of *Proteus mirabilis* isolated from chicken carcasses: potential zoonotic risk. *Brazilian Journal of Microbiology* **2019**, *50*, 685–694.
48. Thi, N.N.; Niwat, C.; J, C.-M.J. Antimicrobial resistance in bacterial poultry pathogens: a review. *Frontiers in veterinary science* **2017**, *4*, 126.
49. Ahn, J.Y.; Ann, H.W.; Jeon, Y.; Ahn, M.Y.; Oh, D.H.; Kim, Y.C.; Kim, E.J.; Song, J.E.; Jung, I.Y.; Kim, M.H.; et al. The impact of production of extended-spectrum beta-lactamases on the 28-day mortality rate of patients with *Proteus mirabilis* bacteremia in Korea. *BMC Infect Dis* **2017**, *17*, 327, doi:10.1186/s12879-017-2431-8.
50. Algammal, A.M.; Hashem, H.R.; Alfifi, K.J.; Hetta, H.F.; SSheraba, N.; Hazem, R.; M, E.R. atpD gene sequencing, multidrug resistance traits, virulence-determinants, and antimicrobial resistance genes of emerging XDR and MDR-*Proteus mirabilis*. *Scientific reports* **2021**, *11*.
51. Girlich, D.; Dortet, L.; Poirel, L.; Nordmann, P. Integration of the bla_{NDM-1} carbapenemase gene into *Proteus* genomic island 1 (PGI1-PmPEL) in a *Proteus mirabilis* clinical isolate. *J Antimicrob Chemother* **2015**, *70*, 98–102, doi:10.1093/jac/dku371.
52. Hammoudi Halat, D.; Ayoub Moubareck, C. The current burden of carbapenemases: review of significant properties and dissemination among Gram-negative bacteria. *Antibiotics (Basel)* **2020**, *9*, doi:10.3390/antibiotics9040186.

53. Sofia, M.; George, S.; Karageorgopoulos, D.E.; Mavros, M.N.; Diamantis, K.; Falagas, M.E. In vitro antimicrobial susceptibility to isepamicin of 6,296 *Enterobacteriaceae* clinical isolates collected at a tertiary care university hospital in Greece. *Antimicrobial agents and chemotherapy* **2012**, *56*, 3067–3073.
54. Pasom, W.; Chanawong, A.; Lulitanond, A.; Wilailuckana, C.; Kenprom, S.; Puang-Ngern, P. Plasmid-mediated quinolone resistance genes, *aac(6')-Ib-cr*, *qnrS*, *qnrB*, and *qnrA*, in urinary isolates of *Escherichia coli* and *Klebsiella pneumoniae* at a teaching hospital, Thailand. *Japanese journal of infectious diseases* **2013**, *66*, 428–432, doi:10.7883/yoken.66.428.
55. Hu, Y.; Cai, J.; Zhang, R.; Zhou, H.; Sun, Q.; Chen, G. Emergence of *Proteus mirabilis* harboring *bla_{KPC-2}* and *qnrD* in a Chinese Hospital. *Antimicrobial agents and chemotherapy* **2012**, *56*, 2278–2282.
56. Lu, J.; Zhang, J.; Xu, L.; Liu, Y.; Li, P.; Zhu, T.; Cheng, C.; Lu, S.; Xu, T.; Yi, H.; et al. Spread of the florfenicol resistance *floR* gene among clinical *Klebsiella pneumoniae* isolates in China. *Antimicrobial resistance and infection control* **2018**, *7*, 127, doi:10.1186/s13756-018-0415-0.
57. Rafael, P.; Paola, S.; Ana, U.; J, M.D.; Pablo, Z. *Proteus mirabilis* uroepithelial cell adhesin (UCA) fimbria plays a role in the colonization of the urinary tract. *Pathogens and disease* **2013**, *67*, 104–107, doi:10.1111/2049-632X.12027.
58. Insua, J.L.; Llobet, E.; Moranta, D.; Pérez-Gutiérrez, C.; Tomás, A.; Garmendia, J.; Bengoechea, J.A. Modeling *Klebsiella pneumoniae* pathogenesis by infection of the wax moth *Galleria mellonella*. *Infection and immunity* **2013**, *81*, 3552–3565, doi:10.1128/iai.00391-13.