

Search for R-Parity-Violation-Induced Charged Lepton Flavor Violation at Future Lepton Colliders

Xunye Cai ¹, Jingshu Li ¹, Ran Ding ¹, Meng Lu ¹, Zhengyun You ^{1,*} and Qiang Li ^{2,*}

¹ School of Physics, Sun Yat-Sen University, Guangzhou 510275, China

² State Key Laboratory of Nuclear Physics and Technology, School of Physics, Peking University, Beijing 100871, China

* Correspondence: youzhy5@mail.sysu.edu.cn (Z.Y.); qliphy0@pku.edu.cn (Q.L.)

Abstract: Interest in searches for Charged Lepton Flavor Violation (CLFV) has continued in the past few decades since the observation of CLFV would indicate a new physics Beyond the Standard Model (BSM). As several future lepton colliders with high luminosity have been proposed, the search for CLFV will reach an unprecedented level of precision. Many BSM models allow CLFV processes at the tree level, such as the R-parity-violating (RPV) Minimal Supersymmetric Standard Model (MSSM), which is a good choice for benchmarking. In this paper, we perform a detailed fast Monte Carlo simulation study on RPV-induced CLFV processes at future lepton colliders, including a 240 GeV circular electron positron collider (CEPC) and a 6 or 14 TeV Muon Collider. As a result, we found that the upper limits on the τ -related RPV couplings will be significantly improved, while several new limits on RPV couplings can be set, which are inaccessible by low-energy experiments.

Keywords: R-parity violation; charged lepton flavor violation; future lepton colliders



Citation: Cai, X.; Li, J.; Ding, R.; Lu, M.; You, Z.; Li, Q. Search for R-Parity-Violation-Induced Charged Lepton Flavor Violation at Future Lepton Colliders. *Universe* **2024**, *10*, 243. <https://doi.org/10.3390/universe10060243>

Academic Editor: Kechen Wang and Ying-nan Mao

Received: 31 March 2024

Revised: 22 May 2024

Accepted: 27 May 2024

Published: 31 May 2024



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Although the Standard Model (SM) has achieved great success in the field of particle physics, it is still an incomplete theory. In the SM, lepton numbers are global $U(1)$ symmetries; and thus the electron lepton number L_e , muon lepton number L_μ , and tau lepton numbers L_τ are separately conserved, as is the total lepton number $L = L_e + L_\mu + L_\tau$. However, it is not consistent with the discovery of neutrino oscillations and non-zero neutrino masses, demonstrating that these symmetries are accidental and that there could be a lepton-flavor-violating short-range interaction among the charged leptons [1]. Therefore, Charged Lepton Flavor Violation (CLFV) processes [2–4] are expected to occur. However, even in the SM extended with a non-zero mass neutrino, CLFV rates are typically suppressed by a factor of $G_F^2 m_\nu^4 \sim 10^{-50}$, which is well below the sensitivity of the experiment and should be unobservable.

Nevertheless, many models of physics Beyond the Standard Model (BSM) introduce new sources of CLFV, such as Supersymmetry (SUSY) [5,6], Z' boson [7], leptoquark [8], quantum black hole (QBH) in low-scale gravity [9], two-Higgs-doublet model (2HDM) [10], and R-parity-violating (RPV) Minimal Supersymmetric Standard Model (MSSM) [11,12]. These models can give rise to sizeable CLFV rates that may be detectable in the next generation of collider experiments. Any such detection of CLFV would be clear evidence for the existence of New Physics (NP) and shed light on the probe of BSM physics.

For the reasons above, the search for CLFV has attracted great interest in recent decades. Numerous experiments have been performed or will be constructed with different approaches, including muon-based experiments, such as $\mu^- N \rightarrow e^- N$ at Mu2e [13,14] and COMET [15], $\mu \rightarrow e\gamma$ at MEG-II [16,17], $\mu \rightarrow eee$ at Mu3e [18,19], and high energy colliders like LEP and LHC looking for CLFV decays of mesons [20–23], μ , τ [24], Z bosons [25–27], and Higgs bosons [28–30]. In the next decades, several proposed lepton colliders, such as the Circular

Electron Positron Collider (CEPC) [31], the Future Circular Collider (FCC)-ee [32], the International Linear Collider (ILC) [33], the Compact Linear Collider (CLIC) [34] and the Muon Collider [35], will be ideal facilities to further probe CLFV processes [36–40] since they have cleaner environments with low backgrounds and higher luminosity than hadron colliders.

In a previous work [41], the potential to search for CLFV signals induced by the Z' model has been studied at the future lepton colliders. In this paper, we focus on the search for CLFV at CEPC and the Muon Collider with RPV-MSSM assumed. The Z' model assumes the existence of an extra Z' boson that couples to different lepton flavors, while RPV-MSSM is another interesting new physics model based on SUSY. The rest of this paper is organized as follows. In Section 2, we will give a brief introduction to RPV-MSSM and its present research status. Section 3 discusses the details of the fast Monte Carlo simulation we performed at CEPC and Muon Collider. In Section 4, we present the numerical results of RPV coupling limits and compare them with current and prospective experimental limits from low-energy μ and τ experiments. Lastly, we close with a conclusion of this paper in Section 5.

2. R-Parity Violating MSSM

MSSM is one of the promising candidates for BSM physics. In the MSSM, renormalizability and gauge invariance do not forbid all the coupling terms that cause lepton number and baryon number violation. It can be prevented by introducing a $\mathbb{Z}_2$ symmetry called R-parity [42].

$$R_p := (-1)^{3B+L+2S}, \quad (1)$$

where B , L , and S denote the baryon number, lepton number, and spin of the particle, respectively. All the SM particles have an R-parity of $+1$, while all the SUSY particles have an R-parity of -1 . One of the motivations for this symmetry is to ensure the stability of the lightest supersymmetric particle, which is a possible dark matter candidate [43–45]. R-parity conservation in MSSM will result in a large transverse missing energy signature at the collider experiment [46–51], but no such signals have been observed so far. Furthermore, RPV-MSSM can give a substantial contribution to muon $g - 2$ calculation through bilinear and trilinear terms in the RPV superpotential [52–54]. Allowing R-parity to be broken becomes acceptable and will give rise to a series of phenomenological consequences; thus, R-parity violation has been extensively studied in phenomenological theory [55–62] and in collider experiments [63–66].

When R-parity is broken, the R-parity violating superpotential must be included:

$$W_{RPV} = \frac{1}{2}\lambda_{ijk}L_iL_jE_k^c + \lambda'_{ijk}L_iQ_jD_k^c + \frac{1}{2}\lambda''_{ijk}U_i^cD_j^cD_k^c + \mu_iL_iH_u, \quad (2)$$

where L , E , Q , U , and D are superfields of lepton, charged lepton, quark, up quark, and down quark respectively; H_u is one of the Higgs superfields; λ_{ijk} , λ'_{ijk} , λ''_{ijk} are Yukawa couplings; and i, j, k denote the three generations. Gauge invariance enforces the antisymmetry of two indices for these couplings, $\lambda_{ijk} = -\lambda_{jik}$ and $\lambda''_{ijk} = -\lambda''_{ikj}$. We can write down the RPV part of the interaction Lagrangian in terms of component fields.

$$\begin{aligned} \mathcal{L}_I = & -\frac{1}{2}\lambda_{ijk}(\tilde{\nu}_{iL}\tilde{l}_{kR}l_{jL} + \tilde{l}_{jL}\tilde{l}_{kR}\nu_{iL} + \tilde{l}_{kR}^*\tilde{\nu}_{iR}^c l_{jL} - (i \leftrightarrow j)) \\ & -\lambda'_{ijk}(\tilde{\nu}_{iL}\tilde{d}_{kR}d_{jL} + \tilde{d}_{jL}\tilde{d}_{kR}\nu_{iL} + \tilde{d}_{kR}^*\tilde{\nu}_{iR}^c d_{jL} \\ & -\tilde{l}_{iL}\tilde{d}_{kR}u_{jL} - \tilde{u}_{jL}\tilde{d}_{kR}l_{iL} - \tilde{d}_{kR}^*\tilde{l}_{iR}^c u_{jL}) + \text{h.c.}, \end{aligned} \quad (3)$$

where $\lambda'' = 0$ is assumed for simplicity since non-zero λ'' corresponds to baryon number violation, which is irrelevant to the search for CLFV. l, ν, u, d denote the field of charged lepton, neutrino, up quark, and down quark, respectively. The tilde over a field represents its superpartner field. The superscripts c and $*$ represent charge conjugation and complex conjugation. The subscripts L and R represent left-handed and right-handed fields.

The interaction Lagrangian can allow for several CLFV processes at the tree level. In this work, we focus on CLFV processes produced by the above $\lambda_{ijk}\tilde{\nu}_{iL}\bar{L}_{kR}l_{jL}$ term, which can contribute to lepton collision CLFV processes at lepton colliders. There have been some studies similar to this paper performed to search for CLFV based on RPV-MSSM, but with different CLFV processes like lepton and meson decay and were studied at different experimental facilities like LHC and COMET [67–74].

3. Simulation and Analysis Framework

In this manuscript, we focus on the CLFV search based on RPV-MSSM and perform the simulation at a 240 GeV electron–positron collider, i.e., the CEPC with an integrated luminosity of 5 ab^{-1} and a 6 or 14 TeV Muon Collider with an integrated luminosity of 4 ab^{-1} .

3.1. Event Simulation

The CLFV signal processes studied in this manuscript include $ee \rightarrow e\mu$, $ee \rightarrow e\tau$, $ee \rightarrow \mu\tau$ at 240 GeV CEPC and $\mu\mu \rightarrow e\mu$, $\mu\mu \rightarrow e\tau$, $\mu\mu \rightarrow \mu\tau$ at 6 or 14 TeV Muon Collider. Figure 1 gives some examples of Feynman diagrams for these CLFV signal processes. The main background processes for each signal process are summarized in Table 1, where the WW and $\tau\tau$ background mean $\ell\ell \rightarrow WW$ or $\tau\tau$, with both W or τ decaying into the corresponding charged leptons in the final state. For the τ -related signal channel like $ee \rightarrow e\tau$, the $\tau\tau$ background process has only one τ decaying into the charged lepton, while the other τ goes through hadronic decay and is reconstructed with the jet collection in DELPHES and likewise for the $H\nu\bar{\nu}(H \rightarrow \tau\tau)$ and $H\nu\bar{\nu}(H \rightarrow WW)$ background processes.




Figure 1. Feynman diagrams of CLFV signal processes, namely the s-channel process (a) and the t-channel process (b), both propagated by s-neutrino.

Table 1. Summary of the CLFV signal and background processes.

Signal Process	Background Processes
$ee \rightarrow e\mu$	$WW, H\nu\bar{\nu}(H \rightarrow \tau\tau), H\nu\bar{\nu}(H \rightarrow WW), \tau\tau$
$ee \rightarrow e\tau$	$WW, H\nu\bar{\nu}(H \rightarrow \tau\tau), \tau\tau$
$ee \rightarrow \mu\tau$	$WW, H\nu\bar{\nu}(H \rightarrow \tau\tau), \tau\tau$
$\mu\mu \rightarrow e\mu$	$WW, WW\nu\bar{\nu}, H\nu\bar{\nu}(H \rightarrow \tau\tau), H\nu\bar{\nu}(H \rightarrow WW), \tau\tau$
$\mu\mu \rightarrow e\tau$	$WW, WW\nu\bar{\nu}, H\nu\bar{\nu}(H \rightarrow \tau\tau), H\nu\bar{\nu}(H \rightarrow WW), \tau\tau$
$\mu\mu \rightarrow \mu\tau$	$WW, WW\nu\bar{\nu}, H\nu\bar{\nu}(H \rightarrow \tau\tau), H\nu\bar{\nu}(H \rightarrow WW), \tau\tau$

Using the UFO [75] model published on the FeynRules model database for RPV extension of MSSM [76], both signal and background process events are generated with MadGraph5_aMC@NLO version 3.4.2 [77,78], and then we perform parton shower and

hadronization with PYTHIA8 version 3.0.6 [79]. In particular, the initial-state radiation (ISR) effect [80] was incorporated into the simulation. Lastly, we used DELPHES version 3.5.0 [81] for detector fast simulation with the default detector configuration cards of CEPC and the Muon Collider.

3.2. Event Selection and Analysis Method

The event selection criteria are described as follows. First, the events must include exactly two charged leptons in the final state with transverse momentum p_T and pseudo-rapidity η satisfying $p_T > 10 \text{ GeV}/c$ and $|\eta| < 2.5$. In particular, for the τ -related channel, τ goes through hadronic decay and is reconstructed with the jet collection in DELPHES, and the jets must satisfy $p_T > 20 \text{ GeV}/c$ and $|\eta| < 5$. These basic cuts applied on p_T and $|\eta|$ follow the default detector configuration cards of DELPHES, reflecting the tracking information provided by the detector. In addition, the final state leptons must meet the requirements of lepton flavor change and charge conservation. For example, in the $e^+e^- \rightarrow e^-\mu^+(e^+\mu^-)$ channel, all events must have only one e^- (e^+) and one μ^+ (μ^-).

For CEPC, the μ tracking efficiency ϵ is set to be 100% within $0.1 < |\eta| \leq 3$, and 0% for $|\eta| > 3$ or $|\eta| \leq 0.1$. For the Muon Collider, the μ tracking efficiency is $\epsilon \geq 90\%$ within $|\eta| \leq 2.5$, and 0% for $|\eta| > 2.5$. For the τ -related channel, as defined in DELPHES default cards of the detector configuration, the τ tagging efficiency is assumed to be 40% for the CEPC and 80% for the Muon Collider with $p_T > 10 \text{ GeV}/c$.

Moreover, we show the invariant mass distributions of final state di-leptons for different channels in Figure 2. To separate the signal from the backgrounds, as the filtering condition, the invariant masses cut is imposed at the value maximizing the quantity $S/\sqrt{S+B}$, where S denotes the event number of the signal and B denotes the event number of the background. Specifically, the invariant mass cuts of the final state di-leptons for the $ee \rightarrow e\mu$ (Figure 2a), $ee \rightarrow e\tau$ (Figure 2c) and $ee \rightarrow \mu\tau$ (Figure 2d) channel at CEPC, $\mu\mu \rightarrow e\mu$ (Figure 2b), $\mu\mu \rightarrow e\tau$ (Figure 2d) and $\mu\mu \rightarrow \mu\tau$ (Figure 2f) channel at the 6 TeV Muon Collider, as well as the case at the 14 TeV Muon Collider, are selected at 220 GeV, 160 GeV, 160 GeV for CEPC, 5.2 TeV, 4 TeV, 4.2 TeV for the 6 TeV Muon Collider, and 10 TeV, 9.5 TeV, 9.5 TeV for the 14 TeV Muon Collider, respectively, in order to maximize signal sensitivities from the backgrounds. After all cuts, we obtain histograms on the final state di-leptons p_T distributions for different channels shown in Figure 3, which can be exploited to set the upper limits on RPV coupling with the method described below.

For each process X , we define a per-event weight $n_{LX} = \sigma_X L / N_X$ to take into account the cross-section difference between the signal and background processes, where L denotes the integrated luminosity of the collider, σ_X denotes the cross-section of process X , and N_X denotes the number of generated events for process X . In our study, we simulate $N_X = 10^5$ events for each signal and background process. The signal and backgrounds yields are re-weighted according to their cross-section to be matched.

The test statistic Z is defined as follows,

$$Z = \sum_{i=1}^{\text{nbins}} Z_i, \quad (4)$$

$$\begin{cases} Z_i := 2[n_i - b_i + b_i \ln(b_i/n_i)] & 95\% \text{ C.L. Exclusion} \\ Z_i := 2[b_i - n_i + n_i \ln(n_i/b_i)] & 5\sigma \text{ Discovery.} \end{cases}$$

where b denotes the SM background yields, $n = s + b$ denotes the total yields including both signal and background, and s denotes the CLFV signal yields. In both cases, each Z_i is subjected to a χ^2 distribution with 1 degree of freedom by Wald's theorem [82], and thus, test statistic Z is subjected to χ^2 distribution with the number of degrees of freedom corresponding to the number of bins [83]. By summing each Z_i in p_T distribution histograms and finding the integral of χ^2 probability density function to match the corresponding significance, we can obtain the upper limits on RPV couplings. The results are summarized and plotted in the next section.




Figure 2. Invariant mass distributions of the final state di-leptons for $ee \rightarrow e\mu$ (a), $ee \rightarrow e\tau$ (c), and $ee \rightarrow \mu\tau$ (e) channel at CEPC, and $\mu\mu \rightarrow e\mu$ (b), $\mu\mu \rightarrow e\tau$ (d), and $\mu\mu \rightarrow \mu\tau$ (f) channel at the 6 TeV Muon Collider. A.U. refers to the Arbitrary Unit.




Figure 3. p_T distributions of the final state di-leptons for $ee \rightarrow e\mu$ (a), $ee \rightarrow e\tau$ (c), and $ee \rightarrow \mu\tau$ (e) channel at CEPC, and $\mu\mu \rightarrow e\mu$ (b), $\mu\mu \rightarrow e\tau$ (d) and $\mu\mu \rightarrow \mu\tau$ (f) channel at the 6 TeV Muon Collider. A.U. refers to the Arbitrary Unit.

4. Results

4.1. CEPC

Using the method described in Section 3.2, the 95% confidence level (C.L.) upper limit results of various couplings versus different s -neutrino masses obtained from the simulation at CEPC are presented in Figure 4. The current most stringent upper limit of these couplings from low-energy μ and τ experiments are also included for comparison. For a heavy lepton a decaying into leptons b and c and an anti-lepton $\bar{d}$ [69],

$$\begin{cases} \Gamma_{a \rightarrow bcd} = \frac{m_{la}^5}{6144\pi^3 m_{\tilde{\nu}_g}^4} \left(\lambda_{gdc}^2 \lambda_{gba}^2 + \lambda_{gcd}^2 \lambda_{gab}^2 + \lambda_{gdb}^2 \lambda_{gca}^2 + \lambda_{gbd}^2 \lambda_{gac}^2 \right) & b \neq c \\ \Gamma_{a \rightarrow bb\bar{d}} = \frac{m_{la}^5}{6144\pi^3 m_{\tilde{\nu}_{gL}}^4} \left(\lambda_{gdb}^2 \lambda_{gba}^2 + \lambda_{gbd}^2 \lambda_{gab}^2 \right). & b = c \end{cases} \quad (5)$$

by which we can convert the upper limits of the branching ratio obtained from the experiment to the upper limits of RPV couplings.




Figure 4. The 95% C.L. upper limit of RPV couplings $|\lambda_{311}\lambda_{312}|$ (equivalent to $|\lambda_{311}\lambda_{321}|$ and $|\lambda_{211}\lambda_{212}|$) from $ee \rightarrow e\mu$ simulation (a), $|\lambda_{211}\lambda_{213}|$ (equivalent to $|\lambda_{211}\lambda_{231}|$ and $|\lambda_{311}\lambda_{313}|$) from $ee \rightarrow e\tau$ simulation (b), $|\lambda_{211}\lambda_{232}|$ (equivalent to $|\lambda_{311}\lambda_{323}|$) from $ee \rightarrow \mu\tau$ simulation (c) versus different s-neutrino masses at CEPC.

As shown in Figure 4, although the $ee \rightarrow e\mu$ channel simulation gives a looser upper limit of couplings $|\lambda_{311}\lambda_{312}|$, $|\lambda_{311}\lambda_{321}|$ and $|\lambda_{211}\lambda_{212}|$ than the result obtained from the SINDRUM experiment [84] and prospect Mu3e experiment, the $ee \rightarrow e\tau$ and $ee \rightarrow \mu\tau$ channel simulation can give more stringent upper limits of couplings $|\lambda_{211}\lambda_{213}|$, $|\lambda_{211}\lambda_{231}|$, $|\lambda_{311}\lambda_{313}|$ and $|\lambda_{211}\lambda_{232}|$, $|\lambda_{311}\lambda_{323}|$ than the current best results from the Belle experiment [85]. However, with future experiments included, the prospective Belle II experiments will give more stringent upper limits of couplings than the $ee \rightarrow e\tau$ and $ee \rightarrow \mu\tau$ channels in our simulation results at CEPC.

4.2. Muon Collider

In the same manner, the 95% confidence level (C.L.) upper limit result of various couplings versus different s-neutrino masses obtained from simulation at the Muon Collider are presented in Figure 5, with the current most stringent upper limit of these couplings

available for comparison. $\mu\mu \rightarrow e\mu$ simulation can set new limits on couplings $|\lambda_{312}\lambda_{322}|$, $|\lambda_{321}\lambda_{322}|$ and $|\lambda_{121}\lambda_{122}|$, which have never been obtained from experiments yet since the process of $\mu \rightarrow \bar{\mu}\mu e$ cannot occur from μ decay.




Figure 5. 95% C.L. upper limit of RPV couplings $|\lambda_{312}\lambda_{322}|$ (equivalent to $|\lambda_{321}\lambda_{322}|$ and $|\lambda_{121}\lambda_{122}|$) from $\mu\mu \rightarrow e\mu$ simulation (a), $|\lambda_{122}\lambda_{131}|$ (equivalent to $|\lambda_{322}\lambda_{313}|$) from $\mu\mu \rightarrow e\tau$ simulation (b), $|\lambda_{122}\lambda_{123}|$ (equivalent to $|\lambda_{122}\lambda_{132}|$ and $|\lambda_{322}\lambda_{323}|$) from $\mu\mu \rightarrow \mu\tau$ simulation (c) versus different s-neutrino masses at the Muon Collider.

The simulation of the $\mu\mu \rightarrow e\tau$ and $\mu\mu \rightarrow \mu\tau$ channel gives more stringent upper limits of couplings $|\lambda_{122}\lambda_{131}|$, $|\lambda_{322}\lambda_{313}|$ and $|\lambda_{122}\lambda_{123}|$, $|\lambda_{122}\lambda_{132}|$, $|\lambda_{322}\lambda_{323}|$ than the current best results from the Belle experiment, and these results are even better than the prospect constraints from the Belle II experiment when the mass of s-neutrino is greater than about 2 TeV.

5. Conclusions and Outlook

In this work, we discuss the sensitivity and the potential for searching for CLFV at future lepton colliders based on RPV-MSSM. By performing fast Monte Carlo simulation of the process $ee \rightarrow e\mu$, $ee \rightarrow e\tau$, $ee \rightarrow \mu\tau$ at 240 GeV CEPC, and $\mu\mu \rightarrow e\mu$, $\mu\mu \rightarrow e\tau$, $\mu\mu \rightarrow \mu\tau$ at the 6 or 14 TeV Muon Collider with MadGraph5, PYTHIA8 and DELPHES, the 95% C.L. upper limits of RPV couplings versus different s-neutrino masses can be obtained. We found that $\mu\mu \rightarrow e\mu$ simulation can set new limits on couplings $|\lambda_{312}\lambda_{322}|$, $|\lambda_{321}\lambda_{322}|$ and $|\lambda_{121}\lambda_{122}|$, which have not yet been obtained by experiments. The $ee \rightarrow e\tau$ and $ee \rightarrow \mu\tau$, $\mu\mu \rightarrow e\tau$ and $\mu\mu \rightarrow \mu\tau$ channel simulation can give more stringent upper limits than the current best results from the Belle experiment. The $\mu\mu \rightarrow e\tau$ and $\mu\mu \rightarrow \mu\tau$ channel simulation results are even better than the results of the prospect Belle II experiment. These simulation results demonstrate that future lepton colliders have some superiority over the current collider experiments and low-energy μ and τ experiments in the search for CLFV.

Author Contributions: Conceptualization, Z.Y. and Q.L.; methodology, Z.Y., M.L. and Q.L.; software, X.C., M.L., J.L. and R.D.; validation, X.C., J.L. and R.D.; formal analysis, X.C.; investigation, X.C.; resources, Z.Y., M.L. and Q.L.; data curation, X.C.; writing—original draft preparation, X.C.; writing—review and editing, Z.Y., M.L. and Q.L.; visualization, X.C.; supervision, Z.Y. and Q.L.; project administration, Z.Y. and Q.L.; funding acquisition, Z.Y., M.L. and Q.L. All authors have read and agreed to the published version of the manuscript.

Funding: This work is supported in part by the National Natural Science Foundation of China under Grant No. 12150005, 12075004, 12175321, 12061141003; the National Key Research and Development Program of China under Grant No. 2018YFA0403900; National College Students Innovation and Entrepreneurship Training Program, Sun Yat-sen University; State Key Laboratory of Nuclear Physics and Technology, Peking University under Grant No. NPT2020KFY04, NPT2020KFY05.

Data Availability Statement: The data supporting the findings of this study will be available on reasonable request from the authors.

Conflicts of Interest: The authors declare no conflicts of interest.

Abbreviations

The following abbreviations are used in this manuscript:

CLFV	Charged Lepton Flavor Violation
SM	Standard Model
BSM	Beyond the Standard Model
RPV	R-parity Violation
MSSM	Minimal Supersymmetric Standard Model
CEPC	Circular Electron Positron Collider
SUSY	Supersymmetry
QBH	Quantum Black Hole
2HDM	Two-Higgs-Doublet Model
NP	New Physics
FCC	Future Circular Collider
ILC	International Linear Collider
CLIC	Compact Linear Collider
ISR	Initial-state Radiation
A.U.	Arbitrary Unit
C.L.	Confidence Level

References

1. Gonzalez-Garcia, M.C.; Nir, Y. Neutrino Masses and Mixing: Evidence and Implications. *Rev. Mod. Phys.* **2003**, *75*, 345–402. [\[CrossRef\]](#)
2. De Gouvea, A.; Vogel, P. Lepton Flavor and Number Conservation, and Physics Beyond the Standard Model. *Prog. Part. Nucl. Phys.* **2013**, *71*, 75–92. [\[CrossRef\]](#)
3. Davidson, S.; Echenard, B.; Bernstein, R.H.; Heeck, J.; Hitlin, D.G. Charged Lepton Flavor Violation. *arXiv* **2022**, arXiv:2209.00142v1. [\[CrossRef\]](#)
4. Bernstein, R.H.; Cooper, P.S. Charged Lepton Flavor Violation: An Experimenter’s Guide. *Phys. Rept.* **2013**, *532*, 27–64. [\[CrossRef\]](#)
5. Lee, I.H. Lepton Number Violation in Softly Broken Supersymmetry. *Phys. Lett. B* **1984**, *138*, 121–127. [\[CrossRef\]](#)
6. Lee, I.H. Lepton Number Violation in Softly Broken Supersymmetry. 2. *Nucl. Phys. B* **1984**, *246*, 120–142. [\[CrossRef\]](#)
7. Langacker, P. The Physics of Heavy Z' Gauge Bosons. *Rev. Mod. Phys.* **2009**, *81*, 1199–1228. [\[CrossRef\]](#)
8. Doršner, I.; Fajfer, S.; Greljo, A.; Kamenik, J.F.; Košnik, N. Physics of leptoquarks in precision experiments and at particle colliders. *Phys. Rept.* **2016**, *641*, 1–68. [\[CrossRef\]](#)
9. Gingrich, D.M. Quantum black holes with charge, colour, and spin at the LHC. *J. Phys. G* **2010**, *37*, 105008. [\[CrossRef\]](#)
10. Branco, G.C.; Ferreira, P.M.; Lavoura, L.; Rebelo, M.N.; Sher, M.; Silva, J.P. Theory and phenomenology of two-Higgs-doublet models. *Phys. Rep.* **2012**, *516*, 1–102. [\[CrossRef\]](#)
11. Choudhury, A.; Mondal, A.; Mondal, S. Status of R-parity violating SUSY. *Eur. Phys. J. Spec. Top.* **2024**, 1–22. [\[CrossRef\]](#)
12. Chemtob, M. Phenomenological constraints on broken R parity symmetry in supersymmetry models. *Prog. Part. Nucl. Phys.* **2005**, *54*, 71–191. [\[CrossRef\]](#)
13. Bartoszek, L. *Mu2e Technical Design Report*; U.S. Department of Energy Office of Scientific and Technical Information: Oak Ridge, TN, USA, 2014. [\[CrossRef\]](#)

14. Mu2e Collaboration. Mu2e Run I Sensitivity Projections for the Neutrinoless $\mu^- \rightarrow e^-$ Conversion Search in Aluminum. *Universe* **2023**, *9*, 54. [\[CrossRef\]](#)
15. Abramishvili, R.; Adamov, G.; Akhmetshin, R.R.; Allin, A.; Angélique, J.C.; Anishchik, V.; Aoki, M.; Aznabayev, D.; Bagaturia, I.; Ban, G.; et al. COMET Phase-I Technical Design Report. *PTEP Prog. Theor. Exp. Phys.* **2020**, *2020*, 033C01. [\[CrossRef\]](#)
16. Renga, F. [MEG-II Collaboration]. The search for lepton flavour violation with the MEG II experiment. *PoS* **2022**, *ICHEP2022*, 708. [\[CrossRef\]](#)
17. Baldini, A.M.; Baranov, V.; Biasotti, M.; Boca, G.; Cattaneo, P.W.; Cavoto, G.; Cei, F.; Chiappini, M.; Chiarello, G.; Corvaglia, A.; et al. The Search for $\mu^+ \rightarrow e^+ \gamma$ with 10–14 Sensitivity: The Upgrade of the MEG Experiment. *Symmetry* **2021**, *13*, 1591. [\[CrossRef\]](#)
18. Arndt, K.; Augustin, H.; Baesso, P.; Berger, N.; Berg, F.; Betancourt, C.; Bortoletto, D.; Bravar, A.; Brigg, K.; vom Bruch, D.; et al. Technical design of the phase I Mu3e experiment. *Nucl. Instrum. Methods A* **2021**, *1014*, 165679. [\[CrossRef\]](#)
19. Dittmeier, S. Searching for cLFV with the Mu3e experiment. *PoS* **2022**, *ICHEP2022*, 692. [\[CrossRef\]](#)
20. Love, W.; Savinov, V.; Lopez, A.; Mehrabyan, S.; Mendez, H.; Ramirez, J.; Huang, G.S.; Miller, D.H.; Pavlunin, V.; Sanghi, B.; et al. Search for Lepton Flavor Violation in Upsilon Decays. *Phys. Rev. Lett.* **2008**, *101*, 201601. [\[CrossRef\]](#)
21. Lees, J.P.; Poireau, V.; Tisser, V.; Grauges, E.; Palano, A.; Eigen, G.; Brown, D.N.; Kolomensky, Y.G.; Fritsch, M.; Koch, H.; et al. Search for Lepton Flavor Violation in $\Upsilon(3S) \rightarrow e^+ \mu^-$. *Phys. Rev. Lett.* **2022**, *128*, 091804. [\[CrossRef\]](#)
22. Ablikim, M. et al. [BESIII Collaboration]. Search for the lepton flavor violating decay $J/\psi \rightarrow e\mu$. *Sci. China Phys. Mech. Astron.* **2023**, *66*, 221011. [\[CrossRef\]](#)
23. Achasov, M.N.; Beloborodov, K.I.; Bergyugin, A.V.; Bogdanchikov, A.G.; Bukin, A.D.; Bukin, D.A.; Dimova, T.V.; Druzhinin, V.P.; Golubev, V.B.; Koop, I.A.; et al. Search for Lepton Flavor Violation Process $e^+ e^- \rightarrow e\mu$ in the Energy Region $\sqrt{s} = 984 - 1060$ MeV and $\phi \rightarrow e\mu$ Decay. *Phys. Rev. D* **2010**, *81*, 057102. [\[CrossRef\]](#)
24. Aubert, B.; Karyotakis, Y.; Lees, J.P.; Poireau, V.; Principe, E.; Prudent, X.; Tisser, V.; Tico, J.G.; Grauges, E.; Martinelli, M.; et al. Searches for Lepton Flavor Violation in the Decays $\tau^+ \rightarrow e^+ \gamma$ and $\tau^+ \rightarrow \mu^+ \gamma$. *Phys. Rev. Lett.* **2010**, *104*, 021802. [\[CrossRef\]](#) [\[PubMed\]](#)
25. Aad, G.; Abbott, B.; Abdallah, J.; Khalek, S.A.; Aben, R.; Abi, B.; Abolins, M.; AbouZeid, O.S.; Abramowicz, H.; Abreu, H.; et al. Search for the lepton flavor violating decay $Z \rightarrow e\mu$ in pp collisions at $\sqrt{s}$ TeV with the ATLAS detector. *Phys. Rev. D* **2014**, *90*, 072010. [\[CrossRef\]](#)
26. Akers, R. et al. [OPAL Collaboration]. A Search for lepton flavor violating Z^0 decays. *Z. Phys. C* **1995**, *67*, 555–564. [\[CrossRef\]](#)
27. Abreu, P. [DELPHI Collaboration]. Search for lepton flavor number violating Z^0 decays. *Z. Phys. C* **1997**, *73*, 243–251. [\[CrossRef\]](#)
28. Aad, G.; Abbott, B.; Abbott, D.C.; Abud, A.A.; Abeling, K.; Abhayasinghe, D.K.; Abidi, S.H.; AbouZeid, O.S.; Abraham, N.L.; Abramowicz, H.; et al. Searches for lepton-flavour-violating decays of the Higgs boson in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector. *Phys. Lett. B* **2020**, *800*, 135069. [\[CrossRef\]](#)
29. Sirunyan, A.M.; Tumasyan, A.; Adam, W.; Ambrogio, F.; Asilar, E.; Bergauer, T.; Brandstetter, J.; Brondolin, E.; Dragicevic, M.; Erö, J.; et al. Search for lepton flavour violating decays of the Higgs boson to $\mu\tau$ and $e\tau$ in proton-proton collisions at $\sqrt{s} = 13$ TeV. *J. High Energy Phys.* **2018**, *6*, 1–54. [\[CrossRef\]](#)
30. Franconi, L.; Haug, S.; Anders, J.K.; Fehr, A.; Ereditato, A. Search for the Higgs boson decays $H \rightarrow ee$ and $H \rightarrow e\mu$ in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. *Phys. Lett. B* **2020**, *801*, 135148. [\[CrossRef\]](#)
31. Group, C.S. CEPC Conceptual Design Report: Volume 1—Accelerator. *arXiv* **2018**, arXiv:1809.00285.
32. CERN FCC Web Site. Available online: <https://fcc.web.cern.ch> (accessed on 30 May 2024).
33. Barklow, T.; Brau, J.; Fujii, K.; Gao, J.; List, J.; Walker, N.; Yokoya, K. ILC Operating Scenarios. *arXiv* **2015**, arXiv:1506.07830. [\[CrossRef\]](#)
34. Charles, T.K.; Giansiracusa, P.J.; Lucas, T.G.; Rassool, R.P.; Volpi, M.; Balazs, C.; Afanaciev, K.; Makarenko, V.; Patapenka, A.; Zhuk, I.; et al. The Compact Linear Collider (CLIC)—2018 Summary Report; 2/2018. *arXiv* **2018**, arXiv:1812.06018. [\[CrossRef\]](#)
35. Delahaye, J.P.; Diemoz, M.; Long, K.; Mansoulié, B.; Pastrone, N.; Rivkin, L.; Schulte, D.; Skrinksky, A.; Wulzer, A. Muon Colliders. *arXiv* **2019**, arXiv:1901.06150. [\[CrossRef\]](#)
36. Li, T.; Schmidt, M.A. Sensitivity of future lepton colliders to the search for charged lepton flavor violation. *Phys. Rev. D* **2019**, *99*, 055038. [\[CrossRef\]](#)
37. Li, T.; Schmidt, M.A.; Yao, C.Y.; Yuan, M. Charged lepton flavor violation in light of the muon magnetic moment anomaly and colliders. *Eur. Phys. J. C* **2021**, *81*, 811. [\[CrossRef\]](#)
38. Homiller, S.; Lu, Q.; Reece, M. Complementary signals of lepton flavor violation at a high-energy muon collider. *J. High Energy Phys.* **2022**, *7*, 36. [\[CrossRef\]](#)
39. Bossi, F.; Ciafaloni, P. Lepton Flavor Violation at muon-electron colliders. *J. High Energy Phys.* **2020**, *10*, 33. [\[CrossRef\]](#)
40. Cirigliano, V.; Fuyuto, K.; Lee, C.; Mereghetti, E.; Yan, B. Charged Lepton Flavor Violation at the EIC. *JHEP* **2021**, *3*, 256. [\[CrossRef\]](#)
41. Li, J.; Wang, W.; Cai, X.; Yang, C.; Lu, M.; You, Z.; Qian, S.; Li, Q. A Comparative Study of Z' mediated Charged Lepton Flavor Violation at future lepton colliders. *JHEP* **2023**, *3*, 190. [\[CrossRef\]](#)
42. Barbier, R.; Bérat, C.; Besançon, M.; Chemtob, M.; Deandrea, A.; Dudas, E.; Fayet, P.; Lavignac, S.; Moreau, G.; Perez, E.; et al. R-parity violating supersymmetry. *Phys. Rep.* **2005**, *420*, 1–202. [\[CrossRef\]](#)
43. Kumar Barman, R.; Belanger, G.; Godbole, R.M. Status of low mass LSP in SUSY. *Eur. Phys. J. ST* **2020**, *229*, 3159–3185. [\[CrossRef\]](#)
44. Barman, R.K.; Bélanger, G.; Bhattacharjee, B.; Godbole, R.M.; Sengupta, R. Is Light Neutralino Thermal Dark Matter in the Phenomenological Minimal Supersymmetric Standard Model Ruled Out? *Phys. Rev. Lett.* **2023**, *131*, 011802. [\[CrossRef\]](#) [\[PubMed\]](#)

45. Chowdhury, D.; Patel, K.M.; Tata, X.; Vempati, S.K. Indirect Searches of the Degenerate MSSM. *Phys. Rev. D* **2017**, *95*, 075025. [\[CrossRef\]](#)
46. Aad, G. et al. [ATLAS Collaboration]. Search for direct pair production of sleptons and charginos decaying to two leptons and neutralinos with mass splittings near the W-boson mass in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector. *JHEP* **2023**, *6*, 031. [\[CrossRef\]](#)
47. Aad, G. et al. [ATLAS Collaboration]. Search for new phenomena in final states with photons, jets and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. *J. High Energy Phys.* **2023**, *7*, 21. [\[CrossRef\]](#)
48. Aad, G. et al. [ATLAS Collaboration]. Searches for new phenomena in events with two leptons, jets, and missing transverse momentum in 139 fb^{-1} of $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector. *Eur. Phys. J. C* **2023**, *83*, 515. [\[CrossRef\]](#)
49. Tumasyan, A. et al. [The CMS Collaboration]. Search for chargino-neutralino production in events with Higgs and W bosons using 137 fb^{-1} of proton-proton collisions at $\sqrt{s} = 13$ TeV. *J. High Energy Phys.* **2021**, *10*, 45. [\[CrossRef\]](#)
50. Tumasyan, A. et al. [The CMS Collaboration]. Search for higgsinos decaying to two Higgs bosons and missing transverse momentum in proton-proton collisions at $\sqrt{s} = 13$ TeV. *J. High Energy Phys.* **2022**, *5*, 14. [\[CrossRef\]](#)
51. Tumasyan, A.; Adam, W.; Andrejkovic, J.W.; Bergauer, T.; Chatterjee, S.; Damanakis, K.; Dragicevic, M.; Del Valle, A.E.; Fruehwirth, R.; Jeitler, M.; et al. Search for electroweak production of charginos and neutralinos at $s = 13 \text{ TeV}$ in final states containing hadronic decays of WW, WZ, or WH and missing transverse momentum. *Phys. Lett. B* **2023**, *842*, 137460. [\[CrossRef\]](#)
52. Martin, S.P.; Wells, J.D. Muon Anomalous Magnetic Dipole Moment in Supersymmetric Theories. *Phys. Rev. D* **2001**, *64*, 035003. [\[CrossRef\]](#)
53. Moroi, T. The Muon anomalous magnetic dipole moment in the minimal supersymmetric standard model. *Phys. Rev. D* **1996**, *53*, 6565–6575; Erratum in *Phys. Rev. D* **1997**, *56*, 4424. [\[CrossRef\]](#) [\[PubMed\]](#)
54. Chakraborty, A.; Chakraborty, S. Probing $(g - 2)_\mu$ at the LHC in the paradigm of R-parity violating MSSM. *Phys. Rev. D* **2016**, *93*, 075035. [\[CrossRef\]](#)
55. Köhler, D. Various Phenomenological Aspects of the R-Parity Violating MSSM. Ph.D. Thesis, Rheinische Friedrich-Wilhelms-Universität Bonn, Bonn, Germany, 2024.
56. Calibbi, L.; D'Eramo, F.; Junius, S.; Lopez-Honorez, L.; Mariotti, A. Displaced new physics at colliders and the early universe before its first second. *J. High Energy Phys.* **2021**, *5*, 234. [\[CrossRef\]](#)
57. Hu, Q.Y.; Li, X.Q.; Muramatsu, Y.; Yang, Y.D. R-parity violating solutions to the $R_{D^{(*)}}$ anomaly and their GUT-scale unifications. *Phys. Rev. D* **2019**, *99*, 015008. [\[CrossRef\]](#)
58. Bardhan, D.; Ghosh, D.; Sachdeva, D. $RK^{(*)}$ from RPV-SUSY s-neutrinos. *Nucl. Phys. B* **2023**, *986*, 116059. [\[CrossRef\]](#)
59. Zheng, M.D.; Chen, F.Z.; Zhang, H.H. Explaining anomalies of B-physics, muon $g - 2$ and W mass in R-parity violating MSSM with seesaw mechanism. *Eur. Phys. J. C* **2022**, *82*, 895. [\[CrossRef\]](#)
60. Zheng, M.D.; Chen, F.Z.; Zhang, H.H. The $W\ell\nu$ -vertex corrections to W-boson mass in the R-parity violating MSSM. *AAPPs Bull.* **2023**, *33*, 16. [\[CrossRef\]](#)
61. Zheng, M.D.; Zhang, H.H. Studying the $b \rightarrow s\ell^+\ell^-$ anomalies and $(g - 2)_\mu$ in R-parity violating MSSM framework with the inverse seesaw mechanism. *Phys. Rev. D* **2021**, *104*, 115023. [\[CrossRef\]](#)
62. Hu, Q.Y.; Yang, Y.D.; Zheng, M.D. Revisiting the B-physics anomalies in R-parity violating MSSM. *Eur. Phys. J. C* **2020**, *80*, 365. [\[CrossRef\]](#)
63. Karmakar, S. Searches for RPV SUSY in ATLAS, CMS and LHCb. *Proc. Sci.* **2023**, *LHCP2022*, 73. [\[CrossRef\]](#)
64. Aad, G.; Abbott, B.; Abbott, D.C.; Abed Abud, A.; Abeling, K.; Abhayasinghe, D.K.; Abidi, S.H.; Aboulhorma, A.; Abramowicz, H.; Abreu, H.; et al. [ATLAS Collaboration] A search for an unexpected asymmetry in the production of $e^+\mu^-$ and $e^-\mu^+$ pairs in proton-proton collisions recorded by the ATLAS detector at $\sqrt{s} = 13$ TeV. *Phys. Lett. B* **2022**, *830*, 137106. [\[CrossRef\]](#)
65. Dercks, D.; Dreiner, H.; Krauss, M.E.; Opferkuch, T.; Reinert, A. R-Parity Violation at the LHC. *Eur. Phys. J. C* **2017**, *77*, 856. [\[CrossRef\]](#)
66. Yamanaka, M. Search for R-parity violation from J-PARC and LHC. *JPS Conf. Proc.* **2015**, *8*, 025001. [\[CrossRef\]](#)
67. Aaboud, M.; Aad, G.; Abbott, B.; Abeloos, B.; Abhayasinghe, D.K.; Abidi, S.H.; AbouZeid, O.S.; Abraham, N.L.; Abramowicz, H.; Abreu, H.; et al. Search for lepton-flavor violation in different-flavor, high-mass final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector. *Phys. Rev. D* **2018**, *98*, 092008. [\[CrossRef\]](#)
68. Sato, J.; Yamanaka, M. A way to crosscheck μ -e conversion in the case of no signals of $\mu \rightarrow e\gamma$ and $\mu \rightarrow 3e$. *Phys. Rev. D* **2015**, *91*, 055018. [\[CrossRef\]](#)
69. Dreiner, H.K.; Kramer, M.; O'Leary, B. Bounds on R-parity violating supersymmetric couplings from leptonic and semi-leptonic meson decays. *Phys. Rev. D* **2007**, *75*, 114016. [\[CrossRef\]](#)
70. Choudhury, D.; Roy, P. New constraints on lepton nonconserving R-parity violating couplings. *Phys. Lett. B* **1996**, *378*, 153–158. [\[CrossRef\]](#)
71. Huitu, K.; Maalampi, J.; Raidal, M.; Santamaria, A. New constraints on R-parity violation from muon e conversion in nuclei. *Phys. Lett. B* **1998**, *430*, 355–362. [\[CrossRef\]](#)
72. Kim, J.E.; Ko, P.; Lee, D.G. More on R-parity and lepton family number violating couplings from muon(ium) conversion, and tau and π^0 decays. *Phys. Rev. D* **1997**, *56*, 100–106. [\[CrossRef\]](#)
73. Dreiner, H.K.; Polesello, G.; Thormeier, M. Bounds on broken R parity from leptonic meson decays. *Phys. Rev. D* **2002**, *65*, 115006. [\[CrossRef\]](#)

74. Littenberg, L.S.; Shrock, R. Implications of improved upper bounds on $|\Delta L| = 2$ processes. *Phys. Lett. B* **2000**, *491*, 285–290. [\[CrossRef\]](#)
75. Degrande, C.; Duhr, C.; Fuks, B.; Grellscheid, D.; Mattelaer, O.; Reiter, T. UFO—The Universal FeynRules Output. *Comput. Phys. Commun.* **2012**, *183*, 1201–1214. [\[CrossRef\]](#)
76. Fuks, B. Beyond the Minimal Supersymmetric Standard Model: From theory to phenomenology. *Int. J. Mod. Phys. A* **2012**, *27*, 1230007. [\[CrossRef\]](#)
77. Alwall, J.; Herquet, M.; Maltoni, F.; Mattelaer, O.; Stelzer, T. MadGraph 5: Going Beyond. *J. High Energy Phys.* **2011**, *6*, 128. [\[CrossRef\]](#)
78. Alwall, J.; Frederix, R.; Frixione, S.; Hirschi, V.; Maltoni, F.; Mattelaer, O.; Shao, H.S.; Stelzer, T.; Torrielli, P.; Zaro, M. The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations. *J. High Energy Phys.* **2014**, *7*, 79. [\[CrossRef\]](#)
79. Sjöstrand, T.; Ask, S.; Christiansen, J.R.; Corke, R.; Desai, N.; Ilten, P.; Mrenna, S.; Prestel, S.; Rasmussen, C.O.; Skands, P.Z. An introduction to PYTHIA 8.2. *Comput. Phys. Commun.* **2015**, *191*, 159–177. [\[CrossRef\]](#)
80. Frixione, S.; Mattelaer, O.; Zaro, M.; Zhao, X. Lepton collisions in MadGraph5_aMC@NLO. *arXiv* **2021**, arXiv:2108.10261. [\[CrossRef\]](#)
81. De Favereau, J.; Delaere, C.; Demin, P.; Giammanco, A.; Lemaître, V.; Mertens, A.; Selvaggi, M. DELPHES 3, A modular framework for fast simulation of a generic collider experiment. *J. High Energy Phys.* **2014**, *2*, 57. [\[CrossRef\]](#)
82. Wald, A. Tests of statistical hypotheses concerning several parameters when the number of observations is large. *Trans. Am. Math. Soc.* **1943**, *54*, 426–482. [\[CrossRef\]](#)
83. Cowan, G.; Cranmer, K.; Gross, E.; Vitells, O. Asymptotic formulae for likelihood-based tests of new physics. *Eur. Phys. J. C* **2011**, *71*, 1554; Erratum in *Eur. Phys. J. C* **2013**, *73*, 2501. [\[CrossRef\]](#)
84. Bellgardt, U.; Otter, G.; Eichler, R.; Felawka, L.; Niebuhr, C.; Walter, H.K.; Bertl, W.; Lordong, N.; Martino, J.; Egli, S. Search for the Decay $\mu^+ \rightarrow e^+ e^+ e^-$. *Nucl. Phys. B* **1988**, *299*, 1–6. [\[CrossRef\]](#)
85. Hayasaka, K.; Inami, K.; Miyazaki, Y.; Arinstein, K.; Aulchenko, V.; Aushev, T.; Bakich, A.M.; Bay, A.; Belous, K.; Bhardwaj, V.; et al. Search for Lepton Flavor Violating Tau Decays into Three Leptons with 719 Million Produced Tau+Tau- Pairs. *Phys. Lett. B* **2010**, *687*, 139–143. [\[CrossRef\]](#)

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.