universe

MDPI

Review

The Spherical Nucleus Puzzle and the Discovery of the New Spherical-like γ-Soft Spectra

Tao Wang

College of Physics, Tonghua Normal University, Tonghua 134000, China; suiyueqiaoqiao@163.com

Abstract

Since the 1950s, nuclear physicists have believed that we have a complete conceptual framework for understanding the low(est)-energy excitations of atomic nuclei. This perspective has persisted in contemporary nuclear structure research, but it now appears overly optimistic. In this review, I present two previously unexpected discoveries, one experimental and one theoretical. Although the spherical phonon excitation spectrum has been considered a typical paradigm of collective excitations in nuclear structure theories, it has not been supported by recent experiments. The result of this experimental discovery reveals a *new* γ-soft rotational mode which has never been predicted by previous theories. This mode differs from previous γ-soft ones and can be described by the newly proposed SU3-IBM theory, a new spherical-like γ-soft mode representing a specific shape phase.

Keywords: spherical nucleus puzzle; new spherical-like γ-soft spectra; SU3-IBM; Cd nuclei; Pd nuclei; SU(3) symmetry

1. Introduction

Over the past century, our understanding of the shapes of atomic nuclei has evolved from a spherical shape to various deformed shapes [1]. After the discovery of the atomic nucleus, this peculiar quantum many-body system at the center of the atom began to be regarded as a "droplet" with a clear boundary; hence, it would have a shape. Of course, at the beginning, it was thought to be a spherical shape (for the whole nuclei). Soon, as revealed by experiments, researchers realized that atomic nuclei can be spherical and also undergo deformation.

After the magic numbers of the nuclei were revealed by the spherical average field and the strong spin–orbit coupling [2,3], the magic nuclei were indeed considered to be spherical. Even today, this is almost true. The key change is our understanding of the collective excitations of the nucleus with valence nucleons. Starting in the 1950s, it was believed that researchers had a thorough grasp of the most fundamental and the most important quadrupole deformations—which govern the collective low(est)-energy excitations of atomic nuclei [4–7]. Yet new experimental data [8–14] and new emerging theories (SU3-IBM) [15–18] suggest that our previous understanding might have been nothing more than an illusion. Deformation has become more and more important since the 1950s, but the notion of a spherical shape has become less and less so. At the same time, our understanding of quadrupole deformations is not sufficient.

When there are a lot of valence nucleons, the residual interactions between them cause the nucleus to deform significantly, which is almost inevitable [6]. However, when there are not so many valence nucleons, it is almost an inevitable intuition that, considering

Check for updates

Academic Editors: Máté Csanád, Panos Christakoglou, Giuseppe Verde and You Zhou

Received: 28 January 2026
Revised: 25 February 2026
Accepted: 27 February 2026
Published: 24 March 2026

Copyright: © 2026 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license.

the spherical shape of the magic nucleus nearby, these nuclei should have a phonon excitation spectra of a spherical shape. So, for the nuclei, during the large period, there is the alternation of the single-particle excitation of the magic nucleus and the collective excitation of the non-magic nuclei, and during the small period, there is the alternation of the spherical phonon excitation and the prolate rotational excitation of the greatly deformed nuclei.

Vibrational and rotational modes appear to occur simultaneously and even intertwine [7]. This perspective was clearly illustrated in the following quote from a 2010 review article by Cejnar et al. [19] in *Reviews of Modern Physics* titled "Quantum phase transitions in the shapes of atomic nuclei":

> "As valence nucleons are added, configuration mixing is generated by the residual interactions, and collective behavior emerges. Typically, as we discuss below, this situation leads to $R_{4/2} \geq 2.0$. ..... We will see shortly that this corresponds to a model of a nucleus that can undergo small amplitude quadrupole (angular momentum 2) oscillations about a spherical equilibrium shape. The line with $R_{4/2} = 3.33$ will be seen to correspond to nonspherical (deformed) nuclei that rotate according to the eigenvalue expression for a quantum mechanical symmetric top." ($R_{4/2}$ is the energy ratio of the 4_1^+ state to the 2_1^+ state).

The experimental research conducted in the past 20 years has begun to change the view of collective evolutions, which is fundamental. The most important change is that the existence of spherical phonon excitation spectra was not confirmed experimentally [8–14]. The phonon excitation spectra are the simplest energy spectra of the geometric model [7], and the lowest energy levels at the beginning of experimental confirmation seem to confirm the existence of this spectra. However, when energy levels are higher, systematic deviations occur [12]. On the other hand, the shape coexistence leads to a misunderstanding of the actual energy spectra of atomic nuclei [16,17].

In 2016, Heyde and Wood [20] made a prescient prediction based on the experimental results:

> "The emerging picture of nuclear shapes is that quadrupole deformation is fundamental to achieving a unified view of nuclear structure. While it has now long been recognized that many nuclei are deformed, the reference frame for nuclear structure discussion has been spherical shapes. We would argue that *a shift in perspective is needed: sphericity is a special case of deformation*. Thus, we argue that the reference frame must be fundamentally one of a deformed many-body system."

And in 2019, Garrett et al. [13]. continued their opinion:

> "Rather than proceeding from spherical, closed-shell nuclei through a region of spherical vibrators before encountering deformation, deformation and shape coexistence may be confronted immediately, even at the closed shell."

These ideas based on the new experiments provide a new description of the low(est)-energy excitations of the nuclear structure for the first time, and some results that were considered correct before are not so.

Since 2019, I have recognized this as a new collective excitation mode which exhibits similarities to the phonon excitation spectra of the spherical shape, while its B(E2) values closely resemble those of the γ-soft rotation (this paper was published in 2022) [15]. Remarkably, this hypothesis has been confirmed over the past six years [16–18]. A new extended interacting boson model with SU(3) higher-order interactions (SU3-IBM) was proposed by me, and it is well fitted to the experimental results. In this review, I comment on the new understanding, which proves the prediction of Heyde and Wood: "a shift in perspective is needed: sphericity is a special case of deformation". The spectrum previ-

ously thought to be excited by spherical phonon vibration is actually a new spherical-like γ-soft phase. This is a story of great coincidence, but it is precisely the history of science's continuous progresses and repetitions.

The rest of this review is organized as follows. First, some basic concepts, ideas and theories are introduced briefly so that researchers who are not familiar with this topic can quickly grasp the key aspects of the change. Next, I introduce the main point of the experimental discoveries, which is to point out that a new collective excitation mode has been found. Finally, I introduce the newly proposed SU3-IBM theory and its fitting works, particularly on ^{106}Pd, which perfectly aligns experimental and theoretical results, confirming the existence of the new spherical-like γ-soft phase.

2. Theoretical Foundations

2.1. The IBM, Shapes and Shape Evolutions

The atomic nucleus is a quantum many-body system composed of protons and neutrons, with the number of protons or neutrons ranging from a few to hundreds, which makes understanding the properties of the atomic nucleus extremely challenging. In the construction of a nuclear Hamiltonian, it is obvious that the proton and neutron are the basic degrees of freedom.

After the interpretation of the magic numbers, the protons and neutrons of the nucleus are divided into two parts. One part is the inert proton core and neutron core, the other part is the valence nucleons, and there are residual interactions between them which produce the low-energy collective excitations of the nucleus.

In the ground state of atomic nuclei, nucleons are prone to pairing due to attractive interactions, similar to the electron pairs that cause superconductivity in solids [21]. Intuitively, however, these nucleon pairs cannot be treated as bosons [22–25].

In 1975, Arima and Iachello proposed a counterintuitive model known as the interacting boson model (IBM) [26–28]. Even today, while this model has indeed achieved tremendous success, this approach to understanding remains somewhat perplexing. In the IBM, the valence nucleon pair is treated as a boson, thus the low-energy states of atomic nuclei can be regarded as an interacting boson system. Therefore, the boson number N is half of the number of valence nucleons. These are the IBM's fundamental assumptions, and the latest theoretical advancements have provided a completely unexpected confirmation of these basic premises [29]. However, this confirmation of the boson hypothesis is even more perplexing. The pairing of nucleons is a complex process, and it is evident that bosons do not exist in atomic nuclei.

This implies that the pairing effect in nuclei may bring some previously unrecognized emergent phenomena, like the ones in high-temperature superconductors [25,30–33].

In the simplest IBM (IBM-1), the proton and neutron are not distinguished, and only the s bosons with angular momentum $L = 0$ and the d bosons with angular momentum $L = 2$ are considered. The constructed Hamiltonian has U(6) symmetry. Thus, the IBM is an algebraic model that allows the application of many familiar group theory techniques. Given that the degrees of freedom of the bosons are only an approximation of the nucleon pairs, it was thought that although it could describe the evolution of the collective excitation of the nuclei well, it could not provided detailed information. This is also the reason why the previous satisfactory agreement between theory and experiment was relatively good.

The IBM has become the language for classifying the collective excitations of atomic nuclei. In previous geometric models [7], the shapes of atomic nuclei are classified by specific solutions, whereas in the IBM, they are determined by specific subgroup limits. The U(6) symmetry has three subgroup limits: (1) The U(5) symmetry limit describes the spherical phonon vibration; (2) the SU(3) symmetry limit describes the rotation of the

prolate shape; and (3) the O(6) symmetry limit describes the γ-soft rotation mode. Around 2000, a new $\overline{\mathrm{SU(3)}}$ symmetry limit was proposed that describes the rotation of the oblate shape [34].

A simple and effective Hamiltonian that describes these shapes is the Q-consistent formalism: follows [35–38]

$$\hat{H}_1 = c[(1-\eta)\hat{n}_d - \frac{\eta}{N}\hat{Q}_\chi \cdot \hat{Q}_\chi], \tag{1}$$

where c, η, and χ are three parameters ($0 \leq \eta \leq 1$); N is the boson number; and $\hat{Q}_\chi = [d^\dagger \times \tilde{s} + s^\dagger \times \tilde{d}]^{(2)} + \chi[d^\dagger \times \tilde{d}]^{(2)}$ is the general quadrupole operator ($-\frac{\sqrt{7}}{2} \leq \chi \leq \frac{\sqrt{7}}{2}$). This Hamiltonian can describe various quadrupole shapes. If $\eta = 0$, it describes the spherical shape (the U(5) symmetry limit). If $\eta = 1$ and $\chi = -\frac{\sqrt{7}}{2}$, it describes the prolate shape (the SU(3) symmetry limit). If $\eta = 1$ and $\chi = 0$, it describes the γ-soft rotation (the O(6) symmetry limit). If $\eta = 1$ and $\chi = \frac{\sqrt{7}}{2}$, it describes the oblate shape (the $\overline{\mathrm{SU(3)}}$ symmetry limit). These symmetry limit points can be seen in Figure 1.




Figure 1. Phase diagram of the IBM-1. This figure can be also found in [39–41].

With this simple model, various shape phase transitions can be studied [19,35–48]. Since 2000, these studies have received a lot of attentions with the study of quantum phase transition and the discoveries of critical point dynamical symmetries which are the approximate exact solutions of the Bohr Hamiltonian [7]. E(5) presents the critical point of the spherical to the γ-soft shape phase transition [49], and X(5) describes the critical point of the spherical to the prolate shape phase transition [50]. The locations of the E(5) and X(5) points can be seen in Figure 1.

The aforementioned works represent our fundamental understanding of the evolutions of the low-energy collective excitation modes in nuclei (quadrupole deformations, shape evolutions and critical descriptions), see Figure 1.

However, the extension to the oblate shape, i.e., the introduction of the $\overline{\mathrm{SU(3)}}$ symmetry limit [34], resulted in a significant conflict between experimental data and theoretical predictions [51]. The problem was not taken seriously at the time. Otherwise, the new SU3-IBM could have been proposed at that time. When considering the oblate shape, some special mathematical structures of the IBM are revealed, that is, the O(6) symmetry limit is not a shape phase, but a critical point of the shape transition from the prolate shape to the oblate shape [51].

In realistic even–even nuclei, the oblate shape is rare, which is the prolate dominance phenomenon. However, in $\hat{H}_1$, this does not seem to be the case. The spectra of the prolate

shape and the oblate shape are the same for the same N, but this mirror symmetry is not found in realistic nuclei. In Ref. [51], the energy ratio $R_{4/2} = E_{4_1^+}/E_{2_1^+}$ of the 4_1^+ and 2_1^+ states of the realistic nuclei in the Hf-Hg region is 3.33 for the prolate shape, while it is 2.55 for the oblate shape. (This difference was not given enough attentions at that time.) Clearly the mirror symmetry in theory does not exist in realistic nuclei, and the realistic prolate and oblate nuclei are asymmetric [29,52].

The actual experimental results show that the previous construction of the Hamiltonian of the IBM-1 is too simple, which leads to the mirror symmetry that does not exist in reality. In this review, I emphasize a new γ-soft spectrum. To better understand this spectrum, familiarity with the energy spectra of the U(5) symmetry limit and the O(6) symmetry limit is essential.

Since 1950, researchers in nuclear structure have believed in the existence of spherical phonon vibration excitation, not only because it is a simple solution of the geometric model, but also because the experimental data seem to support this. From the perspective of hindsight, this is a rare coincidence. For the lowest energy levels, the γ-soft spectra are very similar to the phonon spectra of the spherical shape. What the researchers have observed are actually the γ-soft spectra, though they have been misinterpreting it as the spherical phonon spectra.

Figure 2 shows the spherical harmonic phonon spectra with equal spacing for $N = 6$. For the ground 0_1^+ state, it is a condensate state of 6 s bosons. For the first excited 2_1^+ state, it has 1 d boson and 5 s bosons, which corresponds to the one phonon state of the geometric model. For the two phonon excitations, there are three degenerate 4_1^+, 2_2^+, 0_2^+ states (two-phonon triplet states), which can be easily found in realistic nuclei. For the three phonon excitations, there are five degenerate 6_1^+, 4_2^+, 3_1^+, 2_3^+, 0_3^+ states (three-phonon quintet states). As will be seen later, these quintet states are the focus of misunderstanding.




Figure 2. Energy spectra of the U(5) symmetry limit in $\hat{H}_1$ (the spherical shape).

The energy of the (misunderstood) single-phonon excitation is high (about 0.5 MeV) because these nuclei are close to the magic number. Thus, it is difficult to identify these higher multi-phonon excited states in experiments. It is a key factor that it was not realized that these spectra are not due to phonon excitation. The triplet 4_1^+, 2_2^+, 0_2^+ states were first observed in the experiments, and they were mistaken for the spherical phonon excitation spectrum.

Prior to the IBM's advent, γ-softness was discussed in the geometric model [53], where the potential energy in Hamiltonian was independent of the shape variable γ that describe the triaxiality. After proposing the IBM, Arima and Iachello discovered the new O(6)

symmetry limit [54] beyond the U(5) symmetry limit and the SU(3) symmetry limit, which precisely characterizes γ-softness [55].

The geometric model quantizes the shapes, and when considering the quadrupole deformations, the potential energy is a function of the shape variables β and γ. β is a deformation variable that describes the deviation from sphericity. If the potential energy is independent of γ, the model exhibits the O(5) symmetry.

Figure 3 shows the O(6) γ-soft spectra for $N = 6$. Compared with Figure 2, the O(6) γ-soft spectra exhibit a slight increase in distance of the left levels of the 0_6^+ state (the degeneracy of these states are not changed due to the O(5) symmetry), while the right levels shows an overall upward shift, especially the 0_3^+, 2_4^+ states. Thus, the lowest part of the O(6) γ-soft spectra is very different from that of the spherical phonon spectra. The 4_1^+, 2_2^+ states are degenerate, and 6_1^+, 4_2^+, 3_1^+, 0_2^+ states are degenerate. However, for the highest levels in the U(5) symmetry limit and the O(6) symmetry limit, they are difficult to distinguish.




Figure 3. Energy spectra of the O(6) symmetry limit in $\hat{H}_1$ (the γ-soft rotation).

When η in $\hat{H}_1$ changes from 0 to 1, shape phase transition from the spherical shape to the γ-soft rotation can be described [28,56]. However, some conceptual conflicts will appear. In $\hat{H}_1$, O(6) γ-softness is in fact not a shape phase, it is a critical point of the shape phase transition from the prolate shape to the oblate shape. The reason for this that is the construction of $\hat{H}_1$ is too simple. As the whole transition has the O(5) symmetry, if the angular momenta are the same, then the energy level evolutions display the level-crossing phenomena. This point was discussed in [38].

In previous studies, due to the simplicity of the adopted theories and insufficient experimental data, it was mistakenly concluded that both the U(5) and O(6) symmetry energy spectra exist, which is a rather intriguing coincidence.

2.2. Rigid Triaxility and the SU(3) Symmetry Mapping

Since the birth of quantum mechanics, triaxial rotation has been a basic problem, and it has following Hamiltonian:

$$\hat{H}_2 = \sum \frac{\hat{L}_i^2}{2I_i}, (i = 1, 2, 3) \tag{2}$$

where $\hat{L}_i$ is the i-th component of angular momentum, and I_i is the corresponding moment of inertia. For $\hat{H}_2$, angular momentum $\hat{L}^2$ is a conserved quantity.

For atomic nuclei, it is evident that the rigid triaxial shapes must be considered. If the nucleus is rigid triaxial, then the moments of inertia of rotation in each direction are different, and the rotational effect should be described by $\hat{H}_2$. This was discussed by Davydov et al. based on the geometric model [57,58]. In quantum mechanics, the spherical body does not rotate, and a prolate or oblate body can rotate around an axis perpendicular to its symmetry axis; thus, $\hat{H}_2$ is not needed before.

In the previous understanding of the low-lying collective excitations of the nucleus, the rigid triaxial deformations are not important. For even–even nuclei, in the low-lying spectra, it seems difficult to see the shadow of rigid triaxial deformations. The main collective excitation modes of the nuclear quadrupole deformations are spherical phonon vibration excitation, prolate rotation excitation and γ-soft rotation mode, while the oblate rotation excitation is rare, and the rigid triaxial deformations are only discovered recently [59–62].

Thus, for a long time, $\hat{H}_1$ served as a Hamiltonian that effectively characterizes both the realistic shapes of atomic nuclei and their shape phase transitions. These triaxial deformations are almost forgotten. $\hat{H}_1$, except for the introduction of the mirror symmetry of the prolate shape and the oblate shape, seems to be consistent with the realistic experimental data. It seemed that large-deformed nuclei can indeed be described by the prolate rotational spectra, and γ-soft nuclei can be characterized by the O(6) symmetry limit. Nuclei near the magic numbers can also be modeled by the spherical phonon excitation spectra. All these models appear reasonable, and $\hat{H}_1$ seems particularly plausible.

This is very similar to the previous descriptions of the planetary orbits based on the geocentric theory of circular motion. It was not until Kepler deduced the elliptical orbit based on more accurate observations of Tycho Brahe that the heliocentric theory was confirmed.

Recently, I have come to realize that rigid triaxialities play a decisive role in the study of the quadrupole shapes of nuclei based on more-detailed experimental data, and the introduction of $\hat{H}_2$ becomes more important. The breakthrough is that the spherical phonon excitation spectrum has not been proved by the experiments.

In 1981, Van Isacker and Chen realized that $\hat{H}_1$ cannot describe any rigid triaxial deformation [63]. To describe the rigid triaxial deformations, the third-order interactions must be introduced. They considered the 6 d interaction $[d^\dagger d^\dagger d^\dagger]^{(L)} \cdot [\tilde{d}\tilde{d}\tilde{d}]^{(L)}$. This interaction has been used for a long time to describe the phenomena related to the rigid triaxiality. However, there are some fundamental difficulties. (1) Clearly, this is a higher-order interaction related to the U(5) symmetry. This high-order interaction seems reasonable when the spherical phonon excitations exist, but it is not suitable now. (2) This higher-order interaction, while capable of inducing a rigid triaxial deformation with $\gamma = 30^\circ$ in the large-N limit, exhibits a peculiar energy spectrum for the finite N value that fails to align with the observed nuclear spectra. (3) This higher-order interaction cannot account for other triaxial deformation with γ values that are not 30°.

Of course, there is another way to describe the rigid triaxiality, namely the SU(3) symmetry mapping of the rigid triaxial rotor [64–67].

Elliott was the first to introduce the SU(3) symmetry into the discussions of nuclear structures [68–70]. In the explanation of the magic numbers, the three-dimensional harmonic oscillator potential and the strong spin–orbit coupling are used [2,3]. The potential of three-dimensional simple harmonic oscillator has U(3) symmetry. When the strong spin–orbit coupling is added, the SU(3) symmetry of higher-energy shells is broken, but the SU(3) symmetry of light nuclei is still preserved. Elliott observed this phenomenon, particularly when he discovered that certain light nuclei exhibit characteristic rotational spectra. He then introduced the SU(3) symmetry interaction to explain the properties of light nuclei, and for the first time, he linked the single-particle behavior with the collective excitation.

This is an important breakthrough. In this work, the SU(3) symmetry and the rotation of the prolate shape are linked together for the first time.

However, for the medium and heavy nuclei, there are also significant rotational spectra [67]. This is interpreted as the emergence of the SU(3) symmetry in higher-energy shells, leading to the discovery of pseudo-SU(3) symmetry [71,72]. Higher-energy shells generally lack SU(3) symmetry, but when divided into two parts, the lower part may exhibit SU(3) symmetry, resulting in a rotational spectrum [73,74]. The upper part exhibits O(M) symmetry (M is the number of the single particle orbitals in this part), which appears to be associated with γ-softness.

Recently, Bonatsos et al. presented an updated perspective, revealing that, for higher-energy shells, removing the two highest-energy single-particle orbits results in the lower shells exhibiting the SU(3) symmetry, termed proxy-SU(3) symmetry [75–77]. This changes the ideas induced by the pseudo-SU(3) symmetry, and the whole energy shell can achieve a good approximation of SU(3) symmetry; even a holes-dominated situation can have the SU(3) symmetry.

In Elliott's study, a straightforward correspondence exists between the angular momentum operators ($\hat{L}_i$) and the quadrupole moment operators ($\hat{Q}_j$), which can be mapped to the eight generators of SU(3) symmetry. Therefore, the Hamiltonian they form possesses SU(3) symmetry. In general nuclear structure studies, this correspondence is not considered. In previous studies, this correspondence was only used when discussing the prolate rotation.

If the atomic nucleus has triaxial deformation, its rotation must be described using $\hat{H}_2$. In the SU(3) symmetry mapping, this corresponds to [64–67]

$$\hat{H}_2 = a\hat{L}^2 + b[\hat{L} \times \hat{Q} \times \hat{L}]^{(0)} + c[(\hat{L} \times \hat{Q})^1 \times (\hat{L} \times \hat{Q})^1]^{(0)} \tag{3}$$

where a, b and c are the three transformation parameters. This transformation is very enlightening, which means that the rotation of the rigid triaxiality and the higher-order interactions have direct relationships. Thus, if the rigid triaxiality is necessary in the description of the shapes of the nuclei, then the higher-order interactions are necessary in the SU(3) symmetry mapping. The SU(3) third- and fourth-order interactions appear to be important.

In Ref. [78], this was discussed in detail, not only the rotational effect but also the ground deformation. Thus, the Hamiltonian that can describe the rigid triaxial deformation in the SU(3) symmetry mapping is as follows:

$$\begin{aligned} \hat{H}_3 \quad = \quad & -t_1\hat{C}_2[SU(3)] + t_2\hat{C}_3[SU(3)] + t_3\hat{C}_2^2[SU(3)] \\ & + a\hat{L}^2 + b[\hat{L} \times \hat{Q} \times \hat{L}]^{(0)} + c[(\hat{L} \times \hat{Q})^1 \times (\hat{L} \times \hat{Q})^1]^{(0)} \end{aligned} \tag{4}$$

where $\hat{C}_2[SU(3)]$, $\hat{C}_3[SU(3)]$ are the SU(3) second- and third-order Casimir operators. Ref. [79] further extended the discussions on $\hat{H}_3$. However, in [78,79], the ground deformations are not clearly discussed. They use the combination of $\hat{C}_2[SU(3)]$ and $\hat{C}_2^2[SU(3)]$ to derive any rigid triaxial deformation. Although $\hat{C}_3[SU(3)]$ was also mentioned, it received no attention.

At that time, discussions of the rigid triaxial deformation and its rotation were important, especially this SU(3) symmetry mapping, but in fact such ideas are not really connected with realistic nuclei.

Recent pivotal experiments supporting the existence of the rigid triaxiality revealed that the ^{238}U [80] and ^{154}Sm nuclei [81], previously thought to have the prolate shape, actually possess small rigid triaxiality. These pivotal experiments not only confirm that the large-deformed nuclei are rigid triaxial, but they also validate the correctness of the

SU3-IBM [82,83]. Previous research on the SU(3) symmetry mapping of the rigid triaxial rotation has become critically important. Since 2019, Otsuka et al. has also argued that these large-deformed nuclei are in fact with small rigid triaxiality [84–86].

2.3. The SU3-IBM

Prior to Fortunato et al.'s discussions of the $\hat{C}_3[SU(3)]$ interaction [87], the SU(3) third-order Casimir operator had long been overlooked. In the first paper discussing the SU(3) higher-order interactions in the IBM [64], this item was found to merely alter the $\hat{C}_2[SU(3)]$ scale as a perturbation, thus appearing of little practical value.

This oversight has left us unaware of something extremely important. The reason is that in the study of nuclear structures, it has been always assumed that higher-order interactions exist but can only play a perturbation role. However, from the rigid triaxial deformation and its rotation in $\hat{H}_3$, we can see that the higher-order interactions are actually important (from the prolate shape to any rigid triaxial deformation). The key issue here is that for a long time, the experimental energy spectra failed to demonstrate the utility of the rigid triaxial rotations. A major error here is the misinterpretation of the energy spectra near the previously assumed magic numbers, which was mistakenly assumed to be a spherical excitation spectrum [7]. The misunderstanding has led to a misinterpretation of the collective excitation of the entire atomic nuclei [19].

The existence of magic numbers implies that the interactions between nucleons in the nucleus correspond to a large mean field. Therefore, the remaining part of the correlation interactions seems to be determined by the two-body items. However, there seem to be some new possibilities induced by the pairing interaction which have not **been** explained in the IBM.

If there are more possibilities of two-body interaction, these interactions may bring about quantum cancellation, which may lead to the cancellation of the residual interaction between two-body interactions in the IBM. In this way, there may be strong three-body and four-body interactions in the nuclei near the magic numbers.

The small rigid triaxiality of ^{238}U and ^{154}Sm has decisive consequences, which is that these large-deformed nuclei have rigid triaxiality, so the description of these large-deformed nuclei in the IBM also needs the three-body and four-body interactions. This was never considered in previous IBM research, and even in other nuclear structure models, which has posed fundamental challenges. This means that the three-body and four-body interactions are necessary for the collective excitation of the nucleus. This leads to a fundamental change in the view of collective excitation of atomic nuclei. At least in the IBM, third-order and fourth-order interactions must be considered.

Since 2019, our understanding of collective nuclear excitations has undergone a fundamental transformation. The previously overlooked rigid triaxial deformations and their rotations have now become central topics in nuclear structure, marking a Copernican revolution in the field. Especially in the IBM, $\hat{H}_1$ has become unreasonable, and higher-order interactions must be considered. This may not be obvious in other models.

In 2019, I became aware of the experimental works on spherical nuclei [8–12], which profoundly influenced my understanding of collective excitation in atomic nuclei (before the Otsuka et al.'s work [84] and the Garrett et al.'s work in 2019 [13]). I realized that if spherical nuclei do not exist, then the previous understanding was flawed [19]. If only the two-body interactions are considered, then the new results cannot be obtained. Only when the three-body and four-body interactions are introduced can new results can be obtained. (They do not generate a perturbation effect.) Therefore, I proposed the SU3-

IBM [15] (the first paper was published three years later, after the second paper on the B(E2) anomaly [88]), whose Hamiltonian is

$$\begin{aligned}\hat{H} &= c\{(1-\eta)\hat{n}_d + \eta[-\frac{\hat{C}_2[SU(3)]}{2N} + \alpha\frac{\hat{C}_3[SU(3)]}{2N^2} \\ &+\beta\frac{\hat{C}_2^2[SU(3)]}{2N^3} + \gamma\frac{\Omega}{2N^2} + \delta\frac{\Lambda}{2N^3}]\},\end{aligned} \tag{5}$$

where c, η, α, β, γ, δ are fitting parameters. Ω is $[\hat{L}\times\hat{Q}\times\hat{L}]^{(0)}$ and Λ is $[(\hat{L}\times\hat{Q})^{(1)}\times(\hat{L}\times\hat{Q})^{(1)}]^{(0)}$. $\hat{C}_2[SU(3)]$, $\hat{C}_3[SU(3)]$ are the SU(3) second- and third-order Casimir operators, respectively. During the fitting process, the $\hat{L}^2$ interaction is also incorporated. The SU3-IBM is a new extension of the IBM. In the SU3-IBM, the SU(3) symmetry is the most important and dominates all the quadrupole deformations [15,16]. Historically, SU(3) symmetry has played a very important role in the research on nuclear structure [64,65,67,68,71–73,75,78,79]. Now, it becomes more important, or even fundamental.

In Fortunato et al.'s seminal work [87], the phase diagram of the simplest SU3-IBM was discussed (here $\beta=\gamma=\delta=0$), see Figure 1 (a) in [18]. Compared with $\hat{H}_1$, the key is to replace the $\overline{\mathrm{SU(3)}}$ symmetry limit by describing the oblate shape with the SU(3) third-order Casimir operator $\hat{C}_3[SU(3)]$. Thus, they found a new evolutional path from the prolate shape to the oblate shape. Then, an analytic discussion on the prolate to the oblate shape was given by Zhang et al. [89]. However, unfortunately, the discussion did not continue.

In Zhang et al.'s seminal work [89], two prominent results are found. One is the asymmetry between the prolate shape and the oblate shape, and the other is the boson number odd–even effect at the side of the oblate shape. However, the latter is not considered to exist in realistic nuclei. In this paper, when evolving from the prolate shape to the oblate shape, a first-order phase transition point at $\alpha_s=\frac{3N}{2N+3}$ was confirmed. It was also pointed out this point is a multiple phase coexistence point with the SU(3) irreducible representation (λ,μ) satisfying $\lambda+2\mu=2N$.

The two works [87,89] lay the theoretical foundation for the SU(3) third-order interaction, which is no longer negligible. However, these works lack further in-depth discussions. Otherwise, the SU3-IBM could have been proposed to describe the new characteristics of realist atomic nuclei at that time.

In 2019, I realized that $\hat{H}$ could describe a new spherical-like γ-soft structure [15], which could resolve the perplexing spherical nucleus puzzle discovered experimentally [8–14]. The result was revolutionary, changing our understanding of the collective nature of the nucleus. (This will be discussed in the last two sections.)

Subsequently, the SU3-IBM was found to be able to explain the B(E2) anomaly [88,90–101] discovered in [102–105], to explain the prolate–oblate shape asymmetric evolution [52,87,89,106], to produce the features of the γ-soft nucleus ^{196}Pt at a better level [107], and to produce the E(5)-like spectra in ^{82}Kr [108]. Recently, this model has successfully produced the boson number odd–even effect in $^{196-204}$Hg [29], which is a unique prediction of the SU3-IBM [89]. We have also recently confirmed the ^{166}Er and ^{154}Sm with small rigid triaxiality in the SU3-IBM [82,83]. The shape phase transition from the new γ-soft phase to the prolate shape has also been found [18]. These results prove that the SU3-IBM can give more accurate descriptions of the collective behaviors in nuclei than previous theories and can provide a unified picture for understanding the collective excitations in nuclear structure.

Previously, the IBM was viewed as an approximate framework, making the current outcomes unpredictable. In the SU3-IBM, the SU(3) symmetry is upgraded to a fundamental one, which enables the theory to reveal a wealth of experimental details previously undetectable, representing a pivotal breakthrough in this new framework. This means that the SU(3) symmetry, a whole feature, is revealed in the low-lying excitations governed by

pairing force, which is a key discovery not found in all previous nuclear structure theories. The IBM is not an algebraic correspondence of the geometric model, but a true effective theory for describing the low-lying excitations of atomic nucleus.

3. Experimental Discoveries of the Spherical Nucleus Puzzle

3.1. Cd Nuclei

In 2008, Garrett, Green and Wood published a seminal paper entitled "Breakdown of vibrational motion in the isotopes $^{110-116}$Cd" [8], which marked the experimental discovery of the spherical nucleus puzzle. In this paper, they concluded: "it is possible that the Cd isotopes may not represent vibrational systems and that the essential physics of their motion has been missing."

As described earlier, researchers in the field of nuclear structure have been convinced since 1950 that the spherical phonon excitation spectrum must appear where collectivity begins. The harmonic vibration of the spherical shape has been regarded as one of the paradigms of the low-energy collective quadrupole excitations of atomic nuclei, which is the beginning of the collectivity and has been proved by experimental data. Thus, Ref. [8] was incredible when it first appeared. (Unfortunately, I did not follow up on the related research studies until 2019.) In 2010, Garrett and Wood continued this analysis with a rotational point [9].

In 2012, Garrett et al. presented some critical evidence in another seminal paper supporting this shift of opinions [10]. Near the magic numbers, when the number of valence nucleons is small, not only collectivity but also shape coexistence can appear [109]. Shape coexistence is universal in Cd nuclei. For the normal states, it was discovered long ago that the B(E2) transition probabilities between these energy levels differ significantly from the ones in the spherical phonon spectra, while exhibiting behavior similar to that of the γ-soft nuclei. However, because of the shape coexistence, this inconsistency is explained as the strong coupling between the normal states and the intruder states, which leads to serious distortion of the transition behaviors. In [10], these misunderstandings were corrected, and the coupling strength between the normal states and the intruder states is actually weak. Thus, the B(E2) values are not a distorted phenomenon, and these normal states represent the emergence of γ-soft behavior. They concluded "The decay pattern of the nonintruder states is suggestive of a γ-soft rotor, or O(6) nucleus, rather than a vibrational, or U(5), pattern".

Here, the experimental results have already denied the spherical phonon excitation spectrum, at least for the Cd nuclei (whose energy spectra are very similar), but what it is remains unclear. Clearly, this is like the spherical energy spectrum, but the B(E2) values resemble that of a γ-soft nucleus. It appears to be absent in both the earlier IBM and other nuclear structure models.

The approach based on outdated understanding has persisted for 60 to 70 years, so such an unusual outcome naturally leaves people somewhat perplexed. The spherical phonon vibration mode, which is the first mode studied when learning the geometric model, is especially hard to imagine not existing.

These two papers [8,10] ushered in a new era in nuclear structure research, as their findings were entirely contrary to previous ideas. However, this point, at the beginning, was not well realized. In the history of scientific development, such anomalies always appear, and the subsequent research groups are indifferent.

In 2012, Batchelder et al. provided the decisive spectral signal in a seminal paper [11]. They found that in ^{120}Cd, the 0_3^+ state is absent near the nearly degenerate 6_1^+, 4_2^+,3_1^+ and 2_3^+ (at the previous three-phonon level); see Figure 4a. However, in this paper, they did not propose that this might be a new excitation mode.




Figure 4. Partial experimental low-lying spectra in ^{120}Cd and ^{112}Cd. These figures are from [16].

The reason that the low-energy excitation modes of the Cd nuclei are misunderstood as the spherical phonon excitation modes and have been persisting for 60–70 years is the shape coexistence. It not only contributed to our misinterpretation of the distortion in B(E2) behaviors, but also resulted in a critical misrecognition. In ^{102}Cd, see Figure 4b, the first group intruder states (red) was recognized early, but the second group intruder states (blue) was not recognized until 2019. For a long time, this blue 0^+ state was mistakenly identified as the 0^+ state in the three-phonon excitation of the normal states, leading to the misconception that the normal states here refer to the spherical phonon excitation spectra.

This is a fascinating coincidence, yet the misconception about the collective excitations in nuclei persists to this day.

When the entire Cd isotopes are considered, the overall results become more apparent; see Figure 5. The shape coexistence is obviously not the isolated behavior of a single nucleus, but it appears on the evolution of the energy levels of the whole isotopes. As this is an intruder state behavior involving interactions with multiple nucleons, it exhibits a characteristic parabolic shape, as seen in the evolution of the first green 0^+ states in Figure 5.




Figure 5. Evolution of the part low-lying spectra of $^{108-120}$Cd. This figure is from [16].

The two-phonon triple states 4_1^+, 2_2^+, 0_2^+ (blue) (with previous phonon excitation language) are clear for the whole $^{108-120}$Cd. The five states 6_1^+, 4_2^+,3_1^+, 2_3^+ (red) and the second green 0^+ state have been mistaken as the three phonon quintuple states for a long time, although the second green 0^+ state also shows a clear parabolic feature, but it was not until ^{120}Cd that this became truly clear. The second green 0^+ states are in fact the bandhead of the second intruder states [13,14].

3.2. Other Nuclei

In [12], Garrett et al. present other nuclei showing similar behaviors in the Cd nuclei, such the Te, Pd, Ru, Mo, and Sr nuclei. Therefore, the spherical nucleus puzzle is a more common phenomenon. This reveals that the collective behaviors of nuclei near magic numbers were generally misunderstood in the past.

4. Theoretical Discoveries

4.1. New Spherical-like γ-Soft Spectra

Theoretical discovery is another coincidence. In early 2019, I became aware of the challenge of the spherical nucleus puzzle. Unlike previous studies, I believed that a new collective excitation pattern had emerged here that had never been theoretically discovered before. (This point was also not given by previous experiments.) This may be related to my ongoing research on the IBM's higher-order interactions, and I have also developed the Fortran programs incorporating these interactions [38].

Therefore, a straightforward response was to employ the entire program to tackle the challenge of the spherical nucleus puzzle, reproducing the peculiar energy spectrum of the normal states in nuclei in Figure 5 (at that time the spectra in ^{120}Cd). The finding was astonishing, confirming the existence of a new spherical-like γ-soft mode. When I came to this result, I had not noticed Refs. [87,89]. After making this discovery, I reviewed their works and realized that the SU(3) three-order interaction plays a decisive role here.

Figure 6 presents the new spherical-like γ-soft spectra. For $\hat{H}$, the parameters are $\eta = 0.5$, $\alpha = \frac{3N}{2N+3}$, $\beta = \gamma = \delta = 0$. Therefore, the parameters of this energy spectrum are definite for any N. If $\eta = 1.0$, $\alpha = \frac{3N}{2N+3}$, $\beta = \gamma = \delta = 0$, then this is the SU(3) degenerate point, which is found in [89]. In Figure 6, the obvious degeneracy of energy levels is shown [15], such as 4_1^+, 2_2^+ or 6_1^+, 4_2^+, 3_1^+. This is an unexpected outcome, as there exists

no identical subgroup between the U(5) symmetry limit and the SU(3) degenerate point (SU(3) symmetry). Logically, degeneracy should not occur in this evolutional region. Thus the IBM still has underlying algebraic structures that remain unclear to us. The reason for this degeneracy has not yet been found, but it is now known that this hidden symmetry is closely related to the actual collective excitations of the atomic nuclei.




Figure 6. Energy spectra of the new spherical-like γ-soft mode for $N = 6$ (**a**) and $N = 7$ (**b**). These figures are from [17,107].

From Figure 6, it is clear that 4_1^+, 2_2^+ and 0_2^+ states are nearly degenerate, and in particular there is really no 0_3^+ state near the 6_1^+, 4_2^+, 3_1^+ and 2_2^+ states. This is exactly what the experiments found. In [15,16], we applied this spectrum to fit the Cd data, and the fitting results were indeed excellent. Some details are remarkably consistent, such as the anomalous evolution trend of the electric quadrupole moment $Q_{2_1^+}$ in the first 2^+ state.

As shown in Figure 7, the evolutional trend of $Q_{2_1^+}$ in Cd isotopes differs from that of its neighboring nuclei (Te, Pd, Ru, and Xe). When the number of bosons is between 6 and 9, this value decreases with increasing N. To the best of my knowledge, this feature has never been discussed in previous studies, as it has not been observed in earlier nuclear structure models. In our paper the results are given spontaneously. (The quantitative gap originates from the weak coupling between the normal states and the intruder states.) This feature is also analyzed in detail in a recent paper [18], which explains the different evolutionary trends of Cd and Pd nuclei. A model that can spontaneously provide experimental details significantly enhances its credibility.




Figure 7. (**a**) The evolutional behaviors of the experimental $Q_{2_1^+}$ values in Te, Pd, Cd, Ru, and Xe nuclei [110]. (**b**) The evolutional behaviors of the theoretical (red) and experimental (black) results for $^{108-116}$Cd. These figures are from [16].

This unexpected discovery convinced me that the new spherical-like γ-soft spectrum provided by the SU3-IBM is indeed the genuine solution to the spherical nucleus puzzle. This leads me to seek a conclusion that is absolutely certain and can prove it.

4.2. ^{106}*Pd*

In the study of Cd nuclei, we have only discussed the lowest energy levels, which is insufficient. From Figure 6, compared with the spectra in Figures 2 and 3, we can find that this spectrum is very like the spherical phonon spectrum, but for the higher energy levels, which are very different. The excessive number of energy levels in the Cd nucleus, combined with the shape coexistence, makes it more challenging to investigate higher-energy-level characteristics. So a new nucleus is needed, and the coupling between the normal state and the intruder state can be almost ignored. Thus, its normal states correspond to the new γ-soft spectra we need to determine.

I have discovered the following reference nucleus: ^{106}Pd.

Figure 8 shows the experimental and theoretical results of the low-lying levels of the normal states in ^{106}Pd, up to the 10_1^+ state and under 4000 keV. The degree of agreement between theory and experiment is striking. Clearly, the higher-energy group levels 8_1^+, 6_2^+, 5_1^+, 4_3^+, 4_4^+, 2_4^+ and 0_3^+ and 10_1^+, 8_1^+, 7_1^+, 6_3^+, 5_2^+, 6_4^+, 4_5^+, 3_2^+, 2_5^+ and 0_4^+ really exist and

can be easily found. (The order of labeling in the experiment agrees with Ref. [111].) The theoretical five lowest 0^+ states all have experimental correspondences.




Figure 8. The experimental (**a**) and theoretical (**b**) results of the low-lying levels of the normal states in ^{106}Pd. These figures are from [16].

Table 1 shows the absolute $B(E2)$ values for $E2$ transitions from the low-lying states in ^{106}Pd. Qualitatively, the results of the theory and experiment fit at a good level, and there is no complete inconsistency. For the strong $E2$ transitions, they almost fit very well. The value of $B(E2; 0_2^+ \rightarrow 2_2^+)$ (19(+7-3) W.u.) seems smaller than the theoretical result (84.6 W.u.), but in ^{108}Pd, it is 47(+5-11) W.u., and in $^{112-116}$Cd, they are 99(16), 127(16) and even 3.0×10^4(8) W.u., respectively. Overall, the new theory fits better than the IBM-2 (the protons and neutrons are distinguished) [112,113]. In the IBM-2, the value of $B(E2; 2_3^+ \rightarrow 0_2^+)$ (12.3 W.u.) is much smaller than the experimental result (39(4) W.u.). In ^{108,110}Pd, they are 35(+14-15) and 160(40), respectively. Clearly, the values of $B(E2; 4_2^+ \rightarrow 3_1^+)$, $B(E2; 2_5^+ \rightarrow 3_1^+)$, $B(E2; 2_5^+ \rightarrow 2_3^+)$ and $B(E2; 0_5^+ \rightarrow 2_3^+)$ are all smaller. I look forward to more precise experiments with ^{106}Pd.

Table 1. Absolute $B(E2)$ values in W.u. for $E2$ transitions from the low-lying states in ^{106}Pd. The theory has effective charge $e = 2.027$ (W.u.)$^{1/2}$. [a] From Ref. [110]; [b] from Ref. [114]; [c] from Ref. [111]; [d] from Ref. [112,113]. This table is from [17].

L_i	L_f	Exp. 1 [a]	Exp. 2 [b]	Exp. 3 [c]	Theo.	IBM-2 [d]
2_1^+	0_1^+	44.3(15)	42(4)	44.3(15)	44.3	40.0
2_2^+	2_1^+	44(4)	39(4)	43.7^{+58}_{-50}	49.9	46.0
	0_1^+	1.17(10)	0.87^{+10}_{-9}	1.18^{+15}_{-13}	0.31	0.52
0_2^+	2_1^+	35(8)	43^{+6}_{-9}	35(8)	35.0	44.7
	2_2^+		19^{+7}_{-3}		**84.6**	**27.6**
4_1^+	2_1^+	76(11)	71(7)	76(11)	65.4	64.5
	2_2^+		0.7^{+72}_{-3}		6.2	0.98
3_1^+	2_1^+			0.444^{+57}_{-50}	0.50	0.92
	2_2^+			16.2^{+30}_{-26}	50.2	30.1
	4_1^+			6.0^{+11}_{-60}	7.21	9.38
2_3^+	0_1^+		0.14(2)	0.147^{+25}_{-20}	0.03	0.23
	2_1^+		0.52^{+10}_{-7}	0.52^{+13}_{-10}	0.25	0.13
	2_2^+		10.2^{+22}_{-15}	10^{+2}_{-10}	5.57	2.80
	0_2^+		39(4)	38^{+13}_{-11}	**37.8**	**12.3**
	4_1^+		5.3^{+25}_{-14}	10.6^{+51}_{-42}	**22.7**	**4.70**
4_2^+	2_1^+		0.007^{+6}_{-3}		2.87	0.001
	2_2^+	35(6)	35^{+5}_{-4}	34.5^{+71}_{-54}	28.0	35.2
	4_1^+	21(3)	23^{+3}_{-2}	22.1^{+51}_{-39}	17.4	25.9
	3_1^+			33^{+11}_{-33}	**31.8**	**4.61**
6_1^+	4_1^+	88(9)	89^{+10}_{-13}	88.3^{+97}_{-79}	80.2	75.7
2_5^+	0_1^+			0.062^{+25}_{-21}	0.00	0.00
	2_1^+			0.167^{+83}_{-69}	0.01	0.01
	2_2^+			3.2^{+50}_{-25}	0.04	7.63
	0_2^+			5.4^{+21}_{-19}	0.14	0.00
	3_1^+			54^{+23}_{-20}	**16.9**	**4.67**
	2_3^+			46^{+31}_{-22}	**44.5**	**12.9**
0_5^+	2_1^+				0.57	0.39
	2_2^+				0.04	0.00
	2_3^+				**60.5**	**23.2**
4_4^+	4_1^+			0.43^{+88}_{-36}	0.07	0.39
	2_3^+			15^{+16}_{-11}	5.61	
0_6^+	2_1^+				1.82	
	2_2^+				3.32	
	2_3^+				5.14	
2_8^+	0_1^+			0.153^{+70}_{-53}	0.001	
	2_1^+			0.00003^{+434}_{-3}	0.00	
	2_2^+			0.02^{+11}_{-2}	0.06	
	2_3^+			0.08^{+61}_{-8}	0.04	
	2_5^+				12.5	
8_1^+	6_1^+	105(23)	107^{+13}_{-26}		57.2	
σ					11.2	15.5

Table 2 shows the quadrupole moments of the 2_1^+, 4_1^+, 6_1^+, 2_2^+, 4_2^+ and 2_3^+ states. Clearly, the new theory also agrees with the experimental results very well and better than the calculation results in [112,113].

Table 2. The values of the quadrupole moments of some low-lying states in ^{106}Pd in eb. [a] From Ref. [110]; [b] from Ref. [114]; [c] from Ref. [112,113]. This table is from [17].

	Exp.1 [a]	**Exp.2** [b]	**Theo.**	**IBM-2** [c]
$Q_{2_1^+}$	−0.51(7)	−0.55(5)	−0.47	−0.42
$Q_{4_1^+}$		-0.77^{+5}_{-8}	−0.97	−0.60
$Q_{6_1^+}$		-1.11^{+17}_{-10}	−1.43	−0.70
$Q_{2_2^+}$		0.39^{+5}_{-4}	0.31	0.28
$Q_{4_2^+}$		-0.23^{+14}_{-4}	−0.13	−0.03
$Q_{2_3^+}$		-0.47^{+7}_{-18}	−0.61	−0.23
σ			0.175	0.232

From Tables 1 and 2, the deviation of the IBM-2 is about 1.36 times that of the new theory. The IBM-2 is insufficient to understand the systematic behaviors in the Cd nuclei [8,10], the B(E2) anomaly [102–105] and the prolate–oblate shape asymmetric evolutions [52]. Therefore, it is insufficient for understanding the collective behavior of atomic nuclei. This fit directly confirms the existence of the new spherical-like γ-soft spectra and denies the possibility of the existence of the spherical nucleus in the Cd-Pd nuclei region.

Table 3 shows the possible two-times relationships of the 0^+ states in other candidates with the new spherical-like γ-soft spectra. The possible 0_3^+ state, whose energy is nearly twice that of the 0_2^+ state, is shown. Its energy is also near that of 5_1^+ and 8_1^+ states. These will be discussed in future and with the configuration mixing calculations [8,10].

Table 3. Possible two-times relationships of the 0^+ states in other candidates with spherical-like spectra. ^{106}Pd is also listed. The unit is keV.

	^{108}**Cd**	^{110}**Cd**	^{112}**Cd**	^{114}**Cd**	^{108}**Pd**	^{106}**Pd**	^{102}**Ru**
0_2^+	1913	1731	1433	1306	1053	1134	944
possible 0_3^+	(3800)	3427	2834				
possible 0_3^+		(3489)	2883	2637	(2141)	2278	1968
5_1^+	2565	2927	2666		2084	2366	2219
8_1^+	3683	3275	2881	(2669)	2548	2963	2706

Therefore, the confirmation of ^{106}Pd's energy spectrum is of paramount importance, as it validates that the new spherical-like γ-soft spectrum is indeed the definitive solution of the spherical nucleus puzzle. In the following studies, this new spectrum will be applied to other nuclei which were previously considered as spherical [12].

4.3. Shape Phase Transition from the New Spherical-like γ-Soft Phase to the Prolate Shape

In a recent paper [18], we further explored this new spherical-like γ-soft mode. In the IBM's earlier model ($\hat{H}_1$), the O(6) gamma-softness is a critical point from the prolate to the oblate shape phase transition, which seems somewhat peculiar. In reality, typical γ-soft nuclei such as Xe, Ba, Os, and Pt nuclei do not exhibit this characteristic. More importantly, there is no mirror symmetry between the energy spectrum of the prolate shape and the oblate shape. In the SU3-IBM, this new γ-soft mode represents a real phase. The SU3-IBM predicts a novel shape phase transition from the new γ-soft phase to the

prolate shape (a phenomenon not observed in previous nuclear structure studies), and we have identified key experimental data to support this prediction. ^{108}Pd may be the critical nucleus. If the large-deformed nucleus has small rigid triaxiality, then this new γ-soft phase to the small rigid triaxial shape phase transition may be more realistic, which will be studied in the future.

4.4. Other New Results

In [14,115,116], the deformation of the ground state in the Cd nuclei was found, but the characteristics of the energy spectrum cannot be fully confirmed. In [117,118], the spherical nucleus puzzle was expected to be resolved by the U(5) partial dynamical symmetry, but it was denied by [14]. In ^{106}Pd, the detailed spectra information was derived, which was realized by previous theories.

5. Some Discussions

Here, I mainly emphasize a novel γ-soft excitation mode which has never been discovered in previous nuclear structure studies to the best of my knowledge. This suggests that a key factor was overlooked in previous studies. Here, I expand the scope of my discussion to focus on the large-deformed nuclei. Recent experiments [80,81] and theories [82–86] have revealed that the large-deformed nuclei are not of the prolate shape previously thought but rather have small, rigid triaxial shapes, which almost confirms the correctness and uniqueness of the SU3-IBM in the IBM framework.

The experimental results of the low-lying excitations of the large-deformed nuclei are basically consistent. Here, I observe, consistent with Otsuka et al.'s perspective [84–86], that this 2_2^+ state represents a rotational excitation in one rigid triaxial deformation, which somewhat differs from previous interpretations with γ vibrational excitation of the prolate shape [7].

The key lies in the 0_2^+ state, which has been discussed by researchers previously [119–127]. Here, it represents a new excitation with a rigid triaxial configuration, as explicitly addressed in recent studies [82,83].

Thus, in the SU3-IBM, a novel and definitive explanation for the large-deformed nuclei has been developed, though further research studies are required to advance this field.

The higher-order interactions have long been investigated, yet their connections to realistic nuclei remained unclear before the discovery of the SU3-IBM. The reason for this lies in the fact that insufficient experimental data in earlier studies led to misinterpretations. The recent discovery of the proxy-SU(3) symmetry provides a solid foundation for further understanding the role of these higher-order interactions in the shell model [128,129].

Recent advances have been made in our understanding of the shape of the magic nucleus [130], but these conclusions are not yet fully established. It has been found that some magic nuclei are not spherical [131–133]. A more realistic understanding was made recently [134].

The key problem in scientific studies is the consistency between the theoretical results and the experimental results. The studies cannot ignore the details of the new experimental results and just stay at a rough comparison. This is contrary to the basic principles of scientific research. In this review, I only discuss the most critical new γ-soft mode without providing a comprehensive review. I highlight the key aspects of how to discuss these findings from a broader perspective of scientific understanding. This is crucial, as it indicates that the current understanding of nuclear structure is incomplete.

Beginning with the discovery of the neutron in 1932, the research on nuclear structure has extended for over 90 years. The systematic deterioration of experimental results and theoretical predictions is not accidental, but reflects the inadequacies of our understanding

based on existing theories. This needs more attentions and further studies from nuclear structure researchers. This is the purpose of this review.

6. Conclusions

The long-standing paradigm of spherical ground states and harmonic quadrupole vibrations has been experimentally and theoretically falsified. The low-lying spectra formerly labeled "spherical" are in reality the lowest excitations of a new spherical-like γ-soft phase that emerges naturally when the SU(3) higher-order interactions (three- and four-body interactions in the *s*–*d* boson space) are retained in the IBM. This has led to some fundamental changes, just as Heyde and Wood said: "a shift in perspective is needed: sphericity is a special case of deformation".

Removing the spherical shape does not merely delete one row from the nuclear chart. It forces the entire community to adopt the SU(3) higher-order interactions as the default language for low-lying energy collective phenomena and to re-measure every nucleus against a γ-soft ruler rather than a spherical one, just as Heyde and Wood said: "the reference frame must be fundamentally one of a deformed many-body system".

In this review, I present a new view on the quadrupole deformations in nuclei. The spherical phonon vibrational excitations have been believed in by nuclear structure researchers because of the simple construction of the previous theories and the misunderstanding of the experimental data. Garrett et al.'s experimental works revealed a fact that starkly contradicted the traditional notions, proving the previous understanding to be incorrect. In the experiments, it is also revealed that the large-deformed nuclei are not of the prolate shape as previously understood, but that they have small, rigid triaxiality. This ultimately led to the failure of our previous understanding framework for the collective excitations in nuclei.

The rigid triaxial deformation, which was considered to be unimportant before, has become the core of this new understanding. The SU3-IBM model I proposed has been well fitted to many anomalies, especially the new spherical-like γ-soft spectra which have been experimentally confirmed in ^{106}Pd. These results cannot be regarded as accidental. Therefore, in the nucleus, the pairing interactions bring about a new effect that was not expected before, and the low-lying phenomena of nuclei can be described by the interacting system of many bosons with definite number N. Arima and Iachello's insights carry deeper significance.

Here, the three-body interactions play a dominant role. This does not exist in the previous understanding of nuclear structure. We do not know what the reason for such an effective description is, but what is certain now is that such a description is valid and correct. The IBM is an effective description of the low-lying collective excitations of nuclei, it is more effective than previous our understanding, and it has special consequences. In future studies, starting from the effective field theories [135,136] and the UV-free scheme [137], we aim to uncover the rationale behind this approach and gain a deeper understanding of the collective behaviors of nuclei.

A more comprehensive review of the various anomalies discussed in the SU3-IBM will be given in future.

Funding: This research received no external funding.

Data Availability Statement: No new data were generated or analyzed in this review.

Conflicts of Interest: The author declares no conflicts of interest.

Abbreviations

The following abbreviations are used in this manuscript:

IBM	the interacting boson model
SU3-IBM	the interacting boson model with SU(3) higher-order interactions

References

1. Verney, D. History of the concept of nuclear shape. *Eur. Phys. J. A* **2025**, *61*, 82. [CrossRef]
2. Mayer, M.G. On closed shells in nuclei. ii. *Phys. Rev.* **1949**, *75*, 1969–1970. [CrossRef]
3. Haxel, O.; Jensen, J.H.D.; Suess, H.E. On the magic numbers in nuclear structure. *Phys. Rev.* **1949**, *75*, 1766. [CrossRef]
4. Bohr, A. Rotational motion in nuclei. *Rev. Mod. Phys.* **1976**, *48*, 365–374. [CrossRef]
5. Mottelson, B. Elementary modes of excitation in the nucleus. *Rev. Mod. Phys.* **1976**, *48*, 375–384. [CrossRef]
6. Rainwater, J. Background for the spheroidal nuclear model proposal. *Rev. Mod. Phys.* **1976**, *48*, 385–391. [CrossRef]
7. Bohr, A.; Mottelson, B.R. *Nuclear Structure*; Volume II: Nuclear Deformations; Addison-Wesley/W. A. Benjamin, Inc.: Reading, MA, USA , 1975.
8. Garrett, P.E.; Green, K.L.; Wood, J.L. Breakdown of vibrational motion in the isotopes $^{110-116}$Cd. *Phys. Rev. C* **2008**, *78*, 044307. [CrossRef]
9. Garrett, P.E; Wood, J.L. On the robustness of surface vibrational modes: Case studies in the Cd region. *J. Phys. G Nucl. Part. Phys.* **2010**, *37*, 064028. [CrossRef]
10. Garrett, P.E.; Bangay, J.; Diaz Varela, A.; Ball, G.C.; Cross, D.S.; Demand, G.A.; Finlay, P.; Garnsworthy, A.B.; Green, K.L.; Hackman, G.; et al. Detailed spectroscopy of ^{110}Cd: Evidence for weak mixing and the emergence of γ-soft behavior. *Phys. Rev. C* **2012**, *86*, 044304. [CrossRef]
11. Batchelder, J.C.; Brewer, N.T.; Goans, R.E.; Grzywacz, R.; Griffith, B.O.; Jost, C.; Korgul, A.; Liu, S.H.; Paulauskas, S.V.; Spejewski, E.H.; et al. Low-lying collective states in ^{120}Cd populated by β decay of ^{120}Ag: Breakdown of the anharmonic vibrator model at the three-phonon level. *Phys. Rev. C* **2012**, *86*, 064311. [CrossRef]
12. Garrett, P.E.; Wood, J.L.; Yates, S.W. Critical insights into nuclear collectivity from complementary nuclear spectroscopic methods. *Phys. Scr.* **2018**, *93*, 063001. [CrossRef]
13. Garrett, P.E.; Rodríguez, T.R.; Diaz Varela, A.; Green, K.L.; Bangay, J.; Finlay, A.; Austin, R.A.E.; Ball, G.C.; Bandyopadhyay, D.S.; Bildstein, V.; et al. Multiple shape coexistence in ^{110,112}Cd. *Phys. Rev. Lett.* **2019**, *123*, 142502. [CrossRef]
14. Garrett, P.E.; Rodríguez, T.R.; Diaz Varela, A.; Green, K.L.; Bangay, J.; Finlay, A.; Austin, R.A.E.; Ball, G.C.; Bandyopadhyay, D.S.; Bildstein, V.; et al. Shape coexistence and multiparticle-multihole structures in ^{110,112}Cd. *Phys. Rev. C* **2020**, *101*, 044302.
15. Wang, T. New γ-soft rotation in the interacting boson model with SU(3) higher-order interactions. *Chin. Phys. C* **2022**, *46*, 074101. [CrossRef]
16. Wang, T.; Chen, X.; Zhang, Y. Spherical-like spectra for the description of the normal states of $^{108-120}$Cd in the SU3-IBM and the $Q_{2_1^+}$ anomaly. *Chin. Phys. C* **2025**, *49*, 014107. [CrossRef]
17. Wang, T. Typical new spherical-like γ-soft spectra in ^{104,106,108}Pd. *Phys. Rev. C* **2025**, *112*, 034301. [CrossRef]
18. Zhao, D.H.; Wu, Y.X.; Gong, L.; Yin, Z.Y.; Kang, X.S.; Wang, T. Double shape quantum phase transitions in the SU3-IBM: New γ-soft phase and the shape phase transition from the new γ-soft phase to the prolate shape. *arXiv* **2025** , arXiv:2504.06571.
19. Cejnar, P.; Jolie, J.; Casten, R.F. Quantum phase transitions in the shapes of atomic nuclei. *Rev. Mod. Phys.* **2010**, *82*, 2155–2212. [CrossRef]
20. Heyde, K.; Wood, J.L. Nuclear shapes: From earliest ideas to multiple shape coexisting structures. *Phys. Scr.* **2016**, *91*, 083008. [CrossRef]
21. Bardeen, J.; Cooper, L.N.; Schrieffer, J.R. Theory of superconductivity. *Phys. Rev.* **1952**, *108*, 1175–1204. [CrossRef]
22. Dean, D.J.; Hjorth-Jensen, M. Pairing in nuclear systems: From neutron stars to finite nuclei. *Rev. Mod. Phys.* **2003**, *75*, 607–656. [CrossRef]
23. Strinati, G.C.; Pieri, P.; Röpke, G.; Schuck, P.; Urban, M. The BCS–BEC crossover: From ultra-cold Fermi gases to nuclear systems. *Phys. Rep.* **2018**, *738*, 1–76. [CrossRef]
24. Hung, N.Q.; Dang, N.D.; Moretto, L.G. Pairing in excited nuclei: A review. *Rep. Prog. Phys.* **2019**, *82*, 056301. [CrossRef] [PubMed]
25. Chen, Q.J.; Wang, Z.Q.; Boyack, R.; Yang, S.L.; Levin, K. When superconductivity crosses over: From BCS to BEC. *Rev. Mod. Phys.* **2024**, *96*, 025002. [CrossRef]
26. Arima, A.; Iachello, F. Collective nuclear states as representations of a SU(6) group. *Phys. Rev. Lett.* **1975**, *35*, 1069. [CrossRef]
27. Iachello, F.; Arima, A. *The Interacting Boson Model*; Cambridge University Press: Cambridge, UK, 1987.
28. Casten, R.F.; Warner, D.D. The interacting boson approximation. *Rev. Mod. Phys.* **1988**, *60*, 389–469. [CrossRef]
29. Wang, T.; Zhou, C.X.; Fortunato, L. The boson number hypothesis and the boson number odd-even effect in $^{196-204}$Hg. *arXiv* **2025**, arXiv:2412.14881v2.

30. Demler, E.; Hanke, W.; Zhang, S.C. SO(5) theory of antiferromagnetism and superconductivity. *Rev. Mod. Phys.* **2004**, *76*, 909–974. [CrossRef]
31. Lee, P.A.; Nagaosa, N.; Wen, X.G. Doping a Mott insulator: Physics of high-temperature superconductivity. *Rev. Mod. Phys.* **2006**, *78*, 17–85. [CrossRef]
32. Fradkin, E.; Kivelson, S.A.; Tranquada, J.M. Theory of intertwined orders in high temperature superconductors. *Rev. Mod. Phys.* **2015**, *87*, 457–482. [CrossRef]
33. Fradkin, E. Intertwined orders and the physics of high temperature superconductors. *Particles* **2025**, *8*, 70. [CrossRef]
34. Jolie, J.; Casten, R.F.; von Brentano, P.; Werner, V. Quantum phase transition for γ-soft nuclei. *Phys. Rev. Lett.* **2001**, *87*, 162501. [CrossRef] [PubMed]
35. Cejnar, P.; Jolie, J. Quantum phase transitions studied within the interacting boson model. *Phys. Rev. E* **2000**, *61*, 6237–6247. [CrossRef]
36. Cejnar, P.; Heinze, S.; Jolie, J. Ground-state shape phase transitions in nuclei: Thermodynamic analogy and fnite-N effects. *Phys. Rev. C* **2003**, *68*, 034326. [CrossRef]
37. Iachello, F.; Zamfir, N.V. Quantum phase transitions in mesoscopic systems. *Phys. Rev. Lett.* **2004**, *92*, 212501. [CrossRef]
38. Pan, F.; Wang, T.; Huo, Y.S.; Draayer, J.P. Quantum phase transitions in the consistent-Q Hamiltonian of the interacting boson model. *J. Phys. G Nucl. Part. Phys.* **2008**, *35*, 125105. [CrossRef]
39. Warner, D. A triple point in nuclei. *Nature* **2002**, *420*, 614–615. [CrossRef]
40. Casten, R.F. Shape phase transitions and critical-point phenomena in atomic nuclei. *Nat. Phys.* **2006**, *2*, 811–820. [CrossRef]
41. Casten, R.F.; McCutchan, E.A. Quantum phase transitions and structural evolution in nuclei. *J. Phys. G Nucl. Part. Phys.* **2007**, *34*, R285–R320. [CrossRef]
42. Bonatsos, D.; McCutchan, E.A. Shape phase transitions in geometrical and algebraic nuclear collective models. *Nucl. Phys. News* **2009**, *19*, 13–17. [CrossRef]
43. Casten, R.F. Quantum phase transitions and structural evolution in nuclei. *Prog. Part. Nucl. Phys.* **2009**, *62*, 183–209. [CrossRef]
44. Cejnar, P.; Jolie, J. Quantum phase transitions in the interacting boson model. *Prog. Part. Nucl. Phys.* **2009**, *62*, 210–256. [CrossRef]
45. Jolos, R.V.; Kolganova, E.A. Phase transitions in atomic nuclei. *Phys.-Usp.* **2021**, *64*, 325–343. [CrossRef]
46. Fortunato, L. Quantum phase transitions in algebraic and collective models of nuclear structure. *Prog. Part. Nucl. Phys.* **2021**, *121*, 103891. [CrossRef]
47. Cejnar, P.; Stránský, P.; Macek, M.; Kloc, M. Excited-state quantum phase transitions. *J. Phys. A Math. Theor.* **2021**, *54*, 133001. [CrossRef]
48. Bonatsos, D.; Martinou, A.; Peroulis, S.K.; Mertzimekis, T.J.; Minkov, N. Prolate-oblate shape transitions and O(6) symmetry in even–even nuclei: A theoretical overview. *Phys. Scr.* **2024**, *99*, 062003. [CrossRef]
49. Iachello, F. Dynamic symmetries at the critical point. *Phys. Rev. Lett.* **2000**, *85*, 3580–3583. [CrossRef]
50. Iachello, F. Analytic description of critical point nuclei in a spherical-axially deformed shape phase transition. *Phys. Rev. Lett.* **2001**, *87*, 052502. [CrossRef]
51. Jolie, J.; Linnemann, A. Prolate-oblate phase transition in the Hf-Hg mass region. *Phys. Rev. C* **2003**, *68*, 031301(R). [CrossRef]
52. Wang, T.; He, B.C.; Li, D.K.; Zhou, C.X. Prolate-oblate asymmetric shape phase transition in the interacting boson model with SU(3) higher-order interactions. *Phys. Rev. C* **2023**, *78*, 064322. [CrossRef]
53. Wilets, L.; Jean, M. Surface oscillations in even-even nuclei. *Phys. Rev.* **1956**, *102*, 788–796. [CrossRef]
54. Arima, A.; Iachello, F. New symmetry in the sd boson model of nuclei: The group O(6). *Phys. Rev. Lett.* **1978**, *40*, 385–387. [CrossRef]
55. Cizewski, J.A.; Casten, R.F.; Smith, G.J.; Stelts, M.L.; Börner, H.G.; Davidson, W.F. Evidence for a new symmetry in nuclei: The structure of ^{196}Pt and the O(6) limit. *Phys. Rev. Lett.* **1978**, *40*, 167–170. [CrossRef]
56. Leviatan, A.; Novoselsky, A.; Talmi, I. O(5) symmetry in IBA-1. The O(6)-U(5) transition region. *Phys. Lett. B* **1986**, *172*, 144–148. [CrossRef]
57. Davydov, A.S.; Filippov, G.F. Rotational states in even atomic nuclei. *Nucl. Phys.* **1958**, *8*, 237–249. [CrossRef]
58. Davydov, A.S.; Rostovsky, V.S. Relative transition probabilities between rotational levels of non-axial nuclei. *Nucl. Phys.* **1959**, *12*, 58–68. [CrossRef]
59. Toh, Y.; Chiara, C.J.; McCutchan, E.A.; Walters, W.B.; Janssens, R.V.F.; Carpenter, M.P.; Zhu, S.; Broda, R.; Fornal, B.; Kay, B.P.; et al. Evidence for rigid triaxial deformation at low energy in ^{76}Ge. *Phys. Rev. C* **2013**, *87*, 041304(R). [CrossRef]
60. Forney, A.M.; Walters, W.B.; Chiara, C.J.; Janssens, R.V.F.; Ayangeakaa, A.D.; Sethi, J.; Harker, J.; Alcorta, M.; Carpenter, M.P.; Gürdal, G.; et al. Novel $\Delta J = 1$ sequence in ^{78}Ge: Possible evidence for triaxiality. *Phys. Rev. Lett.* **2018**, *120*, 212501. [CrossRef]
61. Ayangeakaa, A.D.; Janssens, V.F.R.; Zhu, S.; Little, D.; Henderson, J.; Wu, C.Y.; Hartley, D.J.; Albers, M.; Auranen, K.; Bucher, B.; et al. Evidence for rigid triaxial deformation in ^{76}Ge from a model-independent analysis. *Phys. Rev. Lett.* **2019**, *123*, 102501. [CrossRef]

62. Bonatsos, D.; Martinou, A.; Peroulis, S.K.; Petrellis, D.; Vasileiou, P.; Mertzimekis, T.J.; Minkov, N. Triaxial shapes in even–even nuclei: A theoretical overview. *Atoms* **2025**, *13*, 47. [CrossRef]
63. Van Isacker, P.; Chen, J.Q. Classical limit of the interacting boson Hamiltonian. *Phys. Rev. C* **1981**, *24*, 684–689. [CrossRef]
64. Vanden Berghe, G.; De Meyer, H.E.; Van Isacker, P. Symmetry-conserving higher-order interaction terms in the interacting boson mode. *Phys. Rev. C* **1985**, *32*, 1049–1052. [CrossRef]
65. Leschber, Y.; Draayer, J.P. Algebraic realization of rotational dynamics. *Phys. Lett. B* **1987**, *190*, 1–6. [CrossRef]
66. Castaños, O.;Draayer, J.P.; Leschber, Y. Quantum rotor and its SU(3) realization. *Comput. Phys. Comm.* **1988**, *52*, 71–84. [CrossRef]
67. Kota, V.K.B. *SU(3) Symmetry in Atomic Nuclei*; Springer Nature: Singapore, 2020.
68. Elliott, J.P. Collective motion in the nuclear shell model I. classification schemes for states of mixed configurations. *Proc. Roy. Soc.* **1958**, *A*245, 128–145. [CrossRef]
69. Elliott, J.P. Collective motion in the nuclear shell model II. the introduction of intrinsic wavefunctions. *Proc. Roy. Soc.* **1958**, *245*, 562–581.
70. Harvey, M. The nuclear SU(3) model. *Adv. Nucl. Phys.* **1968**, *1*, 67–182.
71. Hecht, K.T.; Adler, A. Generalized seniority for favored $J = 0$ pairs in mixed configurations. *Nucl. Phys. A* **1969**, *137*, 129–143. [CrossRef]
72. Arima, A.; Harvey, M.; Schimizu, K. Pseudo *LS* coupling and pseudo *SU*3 coupling schemes. *Phys. Lett.* **1969**, *B30*, 517–522. [CrossRef]
73. Draayer, J.P.; Weeks, K.J. Shell-model description of the low-energy structure of strongly deformed nuclei. *Phys. Rev. Lett.* **1983**, *51*, 1422–1425. [CrossRef]
74. Draayer, J.P.; Weeks, K.J. Towards a shell model description of the low-energy structure of deformed nuclei I. even-even systems. *Ann. Phys.* **1984**, *156*, 41–67. [CrossRef]
75. Bonatsos, D.; Assimakis, I.E.; Minkov, N.; Martinou, A.; Cakirli, B.; Casten, R.F.; Blaum, K. Proxy-SU(3) symmetry in heavy deformed nuclei. *Phys. Rev. C* **2017**, *95*, 064325. [CrossRef]
76. Bonatsos, D.; Assimakis, I.E.; Minkov, N.; Martinou, A.; Sarantopoulou, S.; Cakirli, R.B.; Casten, R.F.; Blaum, K. Analytic predictions for nuclear shapes, prolate dominance, and the prolate-oblate shape transition in the proxy-SU(3) model. *Phys. Rev. C* **2017**, *95*, 064326. [CrossRef]
77. Bonatsos, D. Prolate over oblate dominance in deformed nuclei as a consequence of the SU(3) symmetry and the Pauli principle. *Eur. Phys. J.* **2017**, *A*53, 148–149. [CrossRef]
78. Smirnov, Y.F.; Smirnova, N.A.; Van Isacker, P. SU(3) realization of the rigid asymmetric rotor within the interacting boson model. *Phys. Rev. C* **2000**, *61*, 041302(R). [CrossRef]
79. Zhang, Y.; Pan, F.; Dai, L.R.; Draayer, J.P. Triaxial rotor in the SU(3) limit of the interacting boson model. *Phys. Rev. C* **2014**, *90*, 044310. [CrossRef]
80. STAR Collaboration. Imaging shapes of atomic nuclei in high-energy nuclear collisions. *Nature* **2024**, *635*, 67–72.
81. Kleemann, J.; Pietralla, N.; Friman-Gayer, U.; Isaak, J.; Papst, O.; Prifti, K.; Werner, V.; Ayangeakaa, A.D.; Beck, T.; Colò, G.; et al. Gamma decay of the ^{154}Sm isovector giant dipole resonance: Smekal-Raman scattering as a novel probe of nuclear ground-state deformation. *Phys. Rev. Lett.* **2025**, *134*, 022503. [CrossRef]
82. Zhou, C.X.; Shang, X.; Wang, T. Different triaxial shapes in the energy spectra of ^{154}Sm. *Phys. Lett. B* **2026**, *874*, 140286. [CrossRef]
83. Zhou, C.X.; Shang, X.; Wang, T. Rigid triaxiality has the SU(3) symmetry: ^{166}Er as an example. *submitted*.
84. Otsuka, T.; Tsunoda, Y.; Abe, T.; Shimizu, N.; Van Duppen, P. Underlying structure of collective bands and self-organization in quantum systems. *Phys. Rev. Lett.* **2019**, *123*, 222502. [CrossRef]
85. Tsunoda, Y.; Otsuka, T. Triaxial rigidity of ^{166}Er and its Bohr-model realization. *Phys. Rev. C* **2021**, *103*, L021303. [CrossRef]
86. Otsuka, T.; Tsunoda, Y.; Shimizu, N.; Utsuno, Y.; Abe, T.; Ueno, H. Prevailing triaxial shapes in atomic nuclei and a quantum theory of rotation of composite object. *Eur. Phys. J. A* **2025**, *61*, 126. [CrossRef]
87. Fortunato, L.; Alonso, C.E.; Arias, J.M.; Garcıa-Ramos, J.E.; Vitturi, A. Phase diagram for a cubic-Q interacting boson model Hamiltonian: Signs of triaxiality. *Phys. Rev. C* **2011**, *84*, 014326. [CrossRef]
88. Wang, T. A collective description of the unusually low ratio $B_{4/2} = B(E2;4_1^+ \to 2_1^+)/B(E2;2_1^+ \to 0_1^+)$. *EPL* **2020**, *129*, 52001. [CrossRef]
89. Zhang, Y.; Pan, F.; Liu, Y.X.; Luo, Y.A.; Draayer, J.P. Analytically solvable prolate-oblate shape phase transitional description within the SU(3) limit of the interacting boson model. *Phys. Rev. C* **2012**, *85*, 064312. [CrossRef]
90. Zhang, Y.; He, Y.W.; Karlsson, D.; Qi, C.; Pan, F.; Draayer, J.P. A theoretical interpretation of the anomalous reduced E2 transition probabilities along the yrast line of neutron-deficient nuclei. *Phys. Lett. B* **2022**, *834*, 137443. [CrossRef]
91. Wang, T. B(E2) anomaly cannot be explained with O(6) higher-order interactions. *Phys. Rev. C* **2023**, *107*, 064303. [CrossRef]
92. Zhang, Y.; Wang, S.N.; Pan, F.; Qi, C.; Draayer, J.P. Triaxial rotor modes in finite-N boson systems. *Phys. Rev. C* **2024**, *110*, 024303. [CrossRef]

93. Pan, F.; Zhang, Y.; Wu, Y.X.; Dai, L.R.; Draayer, J.P. B(E2) anomaly along the yrast line in neutron-deficient $A \approx 170$ even-even nuclei induced by a triaxial rotor term. *Phys. Rev. C* **2024**, *110*, 054324. [CrossRef]
94. Teng, W.; Zhang, Y.; Qi, C. A novel approach for the anomalous collectivity in neutron-deficient Os isotopes. *Chin. Phys. C* **2025**, *49*, 014102. [CrossRef]
95. Zhang, Y.; Teng, W. B(E2) anomaly and triaxial deformation in the interacting boson model. *Phys. Rev. C* **2025**, *111*, 014324. [CrossRef]
96. Teng, W.; Wang, S.N.; Zhang, Y.; Zhao, X.Z.; Deng, X.; Li, X.T. B(E2) anomaly and triaxial deformation within a two-fluid SU(3) symmetry. *Chin. Phys. C* **2025**, *49*, 084106. [CrossRef]
97. Cheng, Y.X.; Zhao, D.H.; Shao, Y.Y.; Gong, L.; Wang, T.; Kang, X.S. SU(3) analysis for B(E2) anomaly. *Chin. Phys. C* **2025**, *49*, 104105. [CrossRef]
98. Teng, W.; Zhang, Y.; Wang, S.N.; Pan, F.; Qi, C.; Draayer, J.P. Anomalous collective modes in atomic nuclei within the proton-neutron interacting boson model. *Phys. Lett. B* **2025**, *865*, 139487. [CrossRef]
99. Teng, W.; Wang, S.N.; Zhao, X.Z.; Zhang, Y. The IBM description of the B(E2) anomaly: Dynamical triaxiality and configuration mixing. *Nucl. Phys. A* **2025**, *1063*, 123214. [CrossRef]
100. Zhang, C.G.; Jin, S.C.; Wang, T.; Wang, T. $B(E2; 2_1^+ \rightarrow 0_1^+)$ anomaly in ^{166}Os. *Chin. Phys. C* **2026**, *50*, 03410.
101. Wang, T.; Cheng, Y.X.; Li, D.K.; Kang, X.S.; Jin, S.C.; Wang, T.; Zhang, Z.Q.; Zhang, C.G.; Zhang, Z.X. Level-anticrossing and new relationship in the B(E2) anomaly. *arXiv* **2025**, arXiv:2503.22100v2.
102. Grahn, T.; Stolze, S.; Joss, D.T.; Page, R.D.; Sayğı, B.; O'Donnell, D.; Akmali, M.; Andgren, K.; Bianco, L.; Cullen, D.M.; et al. Excited states and reduced transition probabilities in ^{168}Os. *Phys. Rev. C* **2016**, *94*, 044327. [CrossRef]
103. Sayğı, B.; Joss, D.T.; Page, R.D.; Grahn, T.; Simpson, J.; O'Donnell, D.; Alharshan, G.; Auranen, K.; Bäck, T.; Boening, S.; et al. Reduced transition probabilities along the yrast line in ^{166}W. *Phys. Rev. C* **2017**, *96*, 021301(R). [CrossRef]
104. Cederwall, B.; Doncel, M.; Aktas, Ö.; Ertoprak, A.; Liotta, R.; Qi, C.; Grahn, T.; Cullen, D.M.; Hodge, D.; Giles, M.; et al. Lifetime measurements of excited states in ^{172}Pt and the variation of quadrupole transition strength with angular momentum. *Phys. Rev. Lett.* **2018**, *121*, 022502. [CrossRef]
105. Goasduff, A.; Ljungvall, J.; Rodríguez, T.R.; Bello Garrote, F.L.; Etile, A.; Georgiev, G.; Giacoppo, F.; Grente, L.; Klintefjord, M.; Kuşoğlu, A.; et al. B(E2) anomalies in the yrast band of ^{170}Os. *Phys. Rev. C* **2019**, *100*, 034302. [CrossRef]
106. Teng, W.; Wang, S.N.; Zhang, Y. Understanding Xe isotopes near $A = 130$ through the prolate-oblate shape phase transition. *Phys. Rev. C* **2025**, *112*, 054317. [CrossRef]
107. Wang, T.; He, B.C.; Zhou, C.X.; Li, D.K.; Fortunato, L. Emerging γ-soft-like spectrum in ^{196}Pt in the SU3-IBM (I). *Chin. Phys. C* **2024**, *48*, 094102. [CrossRef]
108. Zhou, C.X.; Wang, T. E(5)-like emerging γ softness in ^{82}Kr. *Phys. Rev. C* **2023**, *108*, 024309. [CrossRef]
109. Heyde, K.; Wood, J.L. Shape coexistence in atomic nuclei. *Rev. Mod. Phys.* **2011**, *83*, 1467–1521. [CrossRef]
110. ENSDF. Available online: https://www.nndc.bnl.gov/ensdf/ (accessed on 28 January 2026).
111. Prados-Estévez, F.M.; Peters, E.E.; Chakraborty, A.; Mynk, M.G.; Bandyopadhyay, D.; Boukharouba, N.; Choudry, S.N.; Crider, B.P.; Garrett, P.E.; Hicks, S.F.; et al. Collective quadrupole behavior in ^{106}Pd. *Phys. Rev. C* **2017**, *95*, 034328. [CrossRef]
112. Giannatiempo, A.; Nannini, A.; Sona, P. Interacting boson approximation-2 analysis of the Pd and Ru chains. I. Mixed symmetry states of $F_{max}-1$ character in even palladium isotopes. *Phys. Rev. C* **1998**, *58*, 3316–3334. [CrossRef]
113. Giannatiempo, A. Vibrational-γ bands in even $^{104-118}$Pd isotopes. *Phys. Rev. C* **2018**, *98*, 034305. [CrossRef]
114. Svensson, L.E.; Fahlander, C.; Hasselgren, L.; Bäcklin, A.; Westerberg, L.; Cline, D.; Czosnyka, T.; Wu, C.Y.; Diamond, R.M.; Kluge, H. Multiphonon vibrational states in ^{106,108}Pd. *Nucl. Phys. A* **1995**, *584*, 547–572. [CrossRef]
115. Nomura, K.; Jolie, J. Structure of even-even cadmium isotopes from the beyond-mean-field interacting boson model. *Phys. Rev. C* **2018**, *98*, 024303. [CrossRef]
116. Nomura, K.; Karakatsanis, K.E. Collective-model description of shape coexistence and intruder states in cadmium isotopes based on a relativistic energy density functional. *Phys. Rev. C* **2022**, *106*, 064317. [CrossRef]
117. Leviatan, A.; Gavrielov, N.; García-Ramos, J.E.; Van Isacker, P. Quadrupole phonons in the cadmium isotopes. *Phys. Rev. C* **2018**, *98*, 031302. [CrossRef]
118. Gavrielov, N.; García-Ramos, J.E.; Van Isacker, P.; Leviatan, A. Persistent vibrational structure in $^{110-116}$Cd. *Phys. Rev. C* **2023**, *108*, L031305. [CrossRef]
119. Sharpey-Schafer, J.F.; Mullins, S.M.; Bark, R.A.; Gueorguieva, E.; Kau, J.; Komati, F.; Lawrie, J.J.; Maine, P.; Minkova, A.; Murray, S.H.T.; et al. Shape transitional nuclei: What can we learn from the yrare states? or hello the double vacuum; goodbye β-vibrations! *AIP Conf. Proc.* **2008**, *1012*, 19–25. [CrossRef]
120. Sharpey-Schafer, J.F.; Bark, R.A.; Bvumbi, S.P.; Lawrie, E.A.; Lawrie, J.J.; Madiba, T.E.; Majola, S.N.T.; Minkova, A.; Mullins, S.M.; Papka, P.; et al. A double vacuum, configuration dependent pairing and lack of β-vibrations in transitional nuclei: Band structure of N = 88 to N = 91 Nuclei. *Nucl. Phys. A* **2010**, *834*, 45c–49c. [CrossRef]

121. Sharpey-Schafer, J.F.; Mullins, S.M.; Bark, R.A.; Kau, J.; Komati, F.; Lawrie, E.A.; Lawrie, J.J.; Madiba, T.E.; Maine, P.; Minkova, A.; et al. Congruent band structures in ^{154}Gd: Configuration-dependent pairing, a double vacuum and lack of β-vibrations. *Eur. Phys. J. A* **2011**, *47*, 5. [CrossRef]
122. Sharpey-Schafer, J.F.; Madiba, T.E.; Bvumbi, S.P.; Lawrie, E.A.; Lawrie, J.J.; Minkova, A.; Mullins, S.M.; Papka, P.; Roux, D.G.; Timár, J. Blocking of coupling to the 0_2^+ excitation in ^{154}Gd by the [505]11/2$^-$ neutron in ^{155}Gd. *Eur. Phys. J. A* **2011**, *47*, 6. [CrossRef]
123. Sharpey-Schafer, J.F.; Bark, R.A.; Bvumbi, S.P.; Dinoko, T.R.S.; Majola, S.N.T. "Stiff" deformed nuclei, configuration dependent pairing and the β and γ degrees of freedom. *Eur. Phys. J. A* **2019**, *55*, 15. [CrossRef]
124. Sharpey-Schafer, J.F. A proton pairing isomer in ^{174}Hf: Reinterpretation of data and support for the split prolate–oblate pairing approximation for deformed nuclei. *Eur. Phys. J. A* **2023**, *59*, 183. [CrossRef]
125. Sharpey-Schafer, J.F. Decline and fall of nuclear β and γ vibrations. In *Proceedings of South African Institute of Physics (SAIP)*; South African Institute of Physics: Pretoria, South Africa , 2023; p. 218.
126. Garrett, P.E. Characterization of the β vibration and 0_2^+ states in deformed nuclei. *J. Phys. G Nucl. Part. Phys.* **2001**, *27*, R1. [CrossRef]
127. Aprahamian, A.; Lee, K.; Lesher, S.R.; Bijker, R. The nature of 0^+ excitations in deformed nuclei. *Prog. Part. Nucl. Phys.* **2025**, *143*, 104173. [CrossRef]
128. Bonatsos, D.; Assimakis, I.E.; Martinou, A.; Peroulis, S.K.; Sarantopoulou, S.; Minkov, N. Proxy-SU(3) symmetry for heavy deformed nuclei: Nuclear spectra. *Bulg. J. Phys.* **2019**, *46*, 325–336.
129. Castanos, O.; Draayer, J.P.; Leschber, Y. Shape variables and the shell model. *Z. Phys. A* **1988**, *329*, 33–43.
130. Poves, A.; Nowacki, F.; Alhassid, Y. Limits on assigning a shape to a nucleus. *Phys. Rev. C* **2020**, *101*, 054307. [CrossRef]
131. Togashi, T.; Tsunoda, Y.; Otsuka, T.; Shimizu, N.; Honma, M. Novel shape evolution in Sn isotopes from magic numbers 50 to 82. *Phys. Rev. Lett.* **2018**, *121*, 062501. [CrossRef]
132. Taniuchi, R.; Santamaria, C.; Doornenbal, P.; Obertelli, A.; Yoneda, K.; Authelet, G.; Baba, H.; Calvet, D.; Château, F.; Corsi, A.; et al. ^{78}Ni revealed as a doubly magic stronghold against nuclear deformation. *Nature* **2019**, *569*, 53–58. [CrossRef]
133. Henderson, J.; Heery, J.; Rocchini, M.; Siciliano, M.; Sensharma, N.; Ayangeakaa, A.D.; Janssens, R.V.F.; Kowalewski, T.M.; Abhishek, A.D.; Stevenson, P.D.; et al. Deformation and collectivity in doubly magic ^{208}Pb. *Phys. Rev. Lett.* **2025**, *134*, 062502. [CrossRef]
134. Ding, C.R.; Wang, C.C.; Yao, J.M.; Hergert, H.; Liang, H.Z.; Bogner, S.K. From spin to pseudospin symmetry: The Origin of magic numbers in nuclear structure. *Phys. Rev. Lett.* **2026**, *136*, 052501. [CrossRef]
135. Hammer, H.W.; König, S.; van Kolck, U. Nuclear effective field theory: Status and perspectives. *Rev. Mod. Phys.* **2020**, *92*, 025004. [CrossRef]
136. Coello Pérez, E.A.; Papenbrock, T. Effective field theories for collective excitations of atomic nuclei. *J. Phys. G Nucl. Part. Phys.* **2025**, *52*, 033001. [CrossRef]
137. Jia, L.B. Tamed loops: A way to obtain finite loop results without UV divergences. *Commun. Theor. Phys.* **2026**, *78*, 015201. [CrossRef]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.