

Article

Nonsingular Phantom Cosmology in Five-Dimensional $f(R, T)$ Gravity

Rakesh Ranjan Sahoo ¹ , Kamal Lochan Mahanta ² and Saibal Ray ^{3,*} 
¹ Department of Physics, Odisha University of Technology and Research, Techno Campus, Mahalaxmi Vihar, Bhubaneswar 751029, India

² Department of Mathematics, C.V. Raman Global University, Vidya Nagar, Mahura, Janla, Bhubaneswar 752054, India

³ Centre for Cosmology, Astrophysics and Space Science (CCASS), GLA University, Mathura 281406, India

* Correspondence: saibal.ray@gla.ac.in

Abstract: We obtain exact solutions to the field equations for five-dimensional locally rotationally symmetric (LRS) Bianchi type-I spacetime in the $f(R, T)$ theory of gravity, where specifically, the following three cases are considered: (i) $f(R, T) = \mu(R + T)$, (ii) $f(R, T) = R\mu + RT\mu^2$, and (iii) $f(R, T) = R + \mu R^2 + \mu T$, where R and T , respectively, are the Ricci scalar and trace of the energy-momentum tensor. It is found that the equation of state (EOS) parameter w is governed by the parameter μ involved in the $f(R, T)$ expressions. We fine-tune the parameter μ to obtain the effect of phantom energy in the model. However, we also restrict this parameter to obtain a stable model of the universe.

Keywords: phantom energy; LRS Bianchi type-I; $f(R, T)$ theory; 5d spacetimes



Citation: Sahoo, R.R.; Mahanta, K.L.; Ray, S. Nonsingular Phantom Cosmology in Five-Dimensional $f(R, T)$ Gravity. *Universe* **2022**, *8*, 573. <https://doi.org/10.3390/universe8110573>

Academic Editors: Stefano Bellucci, Sergei D. Odintsov and Lorenzo Iorio

Received: 6 August 2022

Accepted: 26 October 2022

Published: 30 October 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Researchers have found it impossible to prevent the addition of dimensions and the unification of forces in nature. It is notable that space and time can be seen as a combined fourth component of special relativity, according to Minkowski [1,2]. In a similar way, Maxwell combined the theories of electricity and magnetism. The next step in this regard was to combine electromagnetism with general relativity (GR) [3]. Over the years, many researchers have been trying to construct unified field theories that geometrize all the fundamental forces of nature. The geometrization of gravity by the general theory of relativity (GR) motivated scientists to propose a higher-dimensional theory that can unify gravitation and electromagnetism [4,5]. Essentially, it has been considered that gravity and electromagnetism are coupled via an additional dimension in the higher-dimensional models in general relativity.

In their interesting work, Chodos and Detweiler [6] showed the evolution of a 5d vacuum universe into a cogent four-dimensional one. Alvarez and Gavela [7] discussed the cosmological scenario in which the dynamical compactification of the higher dimensions produces an abundance of entropy in the universe. Further, they pointed out the possibility of solving the flatness and horizon problems in this scenario.

Using the Kaluza–Klein (KK) theory [4,5], later on, Marciano [8] investigated the mechanism of the evolution of the fundamental constants throughout cosmic time. For this purpose, he derived unique relationships between the low-energy couplings, as well as the masses and propounded that a time variation in any of these parameters can render the proof of higher dimensions. He also reviewed available experimental bounds and, therefore, urged the requirement of new measurements. On the other hand, Gegenberg and Das [9] constructed 5d cosmological models with a real massless non-self interacting scalar field source and pointed out that non-trivial solutions to the field equations occur only when the homogeneous and isotropic three-space has non-positive constant curvature.

It is also interesting to note that Lorenz-Petzold [10] obtained exact solutions to the higher-dimensional field equations in a vacuum, as well as the perfect fluid case along with a non-vanishing cosmological constant.

Wesson [11] considered the higher-dimensional spacetimes with a new challenge by pointing out that “the space part of its metric varies with time in the same way as the de Sitter solution of the conventional four-dimensional theory” and formulated his so-called five-dimensional gravitational theory. Under this theory, Grøn [12] successfully obtained vacuum, radiation, and matter-dominated cosmological models. These models describe an inflationary universe in the variable rest-mass theory as was proposed by Wesson (vide the proposition and application of five-dimensional cosmological principle by Wesson in the Book [13]).

Several authors [14–16] have discussed the KK extension of the FRW cosmological models. In higher dimensions, anisotropic generalizations of these models are available in the literature [17–19], whereas inhomogeneous cosmologies in $5d$ have been studied by other authors [20–22]. A few exact solutions to the Einstein field equations in KK spacetime were obtained by various authors and showed that those reproduce, as well as extend the known solutions of the four dimensions [23–25]. The exact solution to the Einstein field equations, which is Ricci and Riemannian flat in $5d$, was obtained by Liko and Wesson [26]. Interestingly, this solution in $4d$ represents a cosmological model for the early vacuum-dominated universe. Some noteworthy works where variable G and Λ have been studied [27,28] have immense consequences in KK cosmology and higher-dimensional geometry, e.g., Pahwa [29] constructed a homogeneous, anisotropic $4 + d$ cosmological model and, hence, studied the late-time acceleration of the universe.

Higher-dimensional cosmology in various alternative theories of gravity can also be found in the literature, which originated due to a few drawbacks of Einstein’s general relativity (specifically, GR has failed to explain the late-time cosmic acceleration phenomena) and, hence, to comply with the observational evidence. Therefore, one possible technique to justify the observational data [30–37] is the modification of GR. Harko et al. [38] obtained the gravitational field equations in the metric formalism and the equations of motion for the test particles. This theory is called the $f(R, T)$ theory of gravity, where the Lagrangian is an arbitrary function of R and T being the Ricci scalar and trace of the energy–momentum, respectively. Under $f(R, T)$ gravity, various authors have studied different mathematical aspects, as well as physical applications of the theory [39–63].

We design the present article as follows: We provide the basic equations, as well as the Einstein field equations under the cosmological system in Section 2. In Section 3, we have the exact solutions sets under three specific cases, whereas in Section 4, the behavior of the models is presented and analyzed. The results and their discussion are presented in Section 5 to provide some concluding remarks along with salient features. In the current work, units are expressed using the natural system with $G = c = \hbar = 1$.

2. Modified Einstein Field Equations

In their theory, Harko et al. [38] considered three explicit functional forms of $f(R, T)$, which is an arbitrary function of the Ricci scalar R and of the trace of the stress–energy tensor T , as follows:

$$f(R, T) = \begin{cases} R + 2f(T), \\ f_1(R) + f_2(T), \\ f_1(R) + f_2(R)f_3(T). \end{cases} \quad (1)$$

One can therefore obtain several theoretical models for each choice of $f(R, T)$. However, in the present work, we consider the second and third case, i.e., $f(R, T) = f_1(R) + f_2(T)$ and $f(R, T) = f_1(R) + f_2(R)f_3(T)$, for constructing cosmological models through the $5d$ locally rotationally symmetric (LRS) Bianchi type-I spacetime metric in the form

$$ds^2 = dt^2 - A(t)^2 dx^2 - B(t)^2 (dy^2 + dz^2) - F(t)^2 dn^2, \quad (2)$$

where A , B , and F are functions of the time coordinate only and the extra dimension n is the space like coordinate.

Now, the gravitational field equations can be provided as [38]

$$f_R(R, T)R_{ij} - \frac{1}{2}g_{ij}f(R, T) + (g_{ij}\square - \nabla_i\nabla_j)f_R(R, T) = 8\pi T_{ij} - f_T(R, T)T_{ij} - f_T(R, T)\Theta_{ij}, \quad (3)$$

where $f_R(R, T) = \frac{\partial f(R, T)}{\partial R}$ and $f_T(R, T) = \frac{\partial f(R, T)}{\partial T}$ are the partial derivative with respect to R and T , respectively, $\square = \nabla_i\nabla^i$, ∇_i denotes covariant derivatives, and $\Theta_{ij} = -2T_{ij} - g_{ij}P$.

Various authors have studied the choice of Lagrangian density for matter, i.e., $\mathcal{L}_m = P$ [64–66], and have addressed the problem of Lagrangian density for perfect fluids where P is the pressure. In this case, we assume that the entire universe is filled with a perfect fluid. The choice for the matter Lagrangian density is thus made to be $\mathcal{L}_m = -P$ [67].

Here, we consider the source of gravitation as the perfect fluid. Therefore, the energy–momentum tensor is taken as

$$T_{ij} = (P + \rho)u_i u_j - P g_{ij}, \quad (4)$$

together with the comoving coordinates

$$g_{ij}u^i u^j = 1. \quad (5)$$

In the above equations, ρ and u_i are the energy density and five-velocity vector of the cosmic fluid distribution, respectively.

3. Solutions to the Field Equations

3.1. $f(R, T) = \mu R + \mu T$

Let us consider here the second case, i.e., $f(R, T) = f_1(R) + f_2(T)$ with $f_1(R) = \mu R$, $f_2(T) = \mu T$, where

$$R = \frac{2A_{tt}}{A} + \frac{4A_t B_t}{AB} + \frac{2A_t F_t}{AF} + \frac{4B_{tt}}{B} + \frac{4B_t F_t}{BF} + \frac{2B_t^2}{B^2} + \frac{2F_{tt}}{F},$$

$T = \rho(t) - 4P(t)$, and μ is an arbitrary constant.

Now, Equation (3) becomes

$$G_{ij} = \left[\frac{8\pi + \mu}{\mu} \right] T_{ij} + \left[P + \frac{T}{2} \right] g_{ij}. \quad (6)$$

Here, $G_{ij} = R_{ij} - \frac{1}{2}g_{ij}R$ is the Einstein tensor. For the line element (2), the explicit form of the field Equation (6) using (4) and (5) can be obtained as

$$-\frac{2\mu A_t B_t}{AB} - \frac{\mu A_t F_t}{AF} - \frac{2\mu B_t F_t}{BF} - \frac{\mu B_t^2}{B^2} + \mu P - \frac{3}{2}\mu\rho - 8\pi\rho = 0, \quad (7)$$

$$\frac{2\mu B_{tt}}{B} + \frac{2\mu B_t F_t}{BF} + \frac{\mu B_t^2}{B^2} + \frac{\mu F_{tt}}{F} - 2\mu P - 8\pi P + \frac{1}{2}\mu\rho = 0, \quad (8)$$

$$\frac{\mu A_{tt}}{A} + \frac{\mu A_t B_t}{AB} + \frac{\mu A_t F_t}{AF} + \frac{\mu B_{tt}}{B} + \frac{\mu B_t F_t}{BF} + \frac{\mu F_{tt}}{F} - 2\mu P - 8\pi P + \frac{1}{2}\mu\rho = 0, \quad (9)$$

$$\frac{\mu A_{tt}}{A} + \frac{2\mu A_t B_t}{AB} + \frac{2\mu B_{tt}}{B} + \frac{\mu B_t^2}{B^2} - 2\mu P - 8\pi P + \frac{1}{2}\mu\rho = 0. \quad (10)$$

Here and in what follows, the suffix “ t ” after a field variable represents an ordinary differentiation with respect to the time “ t ”.

In order to derive the exact solution of the field Equations (7)–(10), we take the following scale transformations [68]:

$$\begin{aligned} A(t) &= e^{\alpha(\tau)}, \\ B(t) &= e^{\gamma(\tau)}, \\ F(t) &= e^{\lambda(\tau)} \\ dt &= AB^2 F d\tau. \end{aligned} \quad (11)$$

Now, the field Equations (7)–(10) using (11) reduce to

$$2\mu P - (3\mu + 16\pi)\rho + e^{-2(\alpha+2\gamma+\lambda)}[-2\mu\lambda'(\alpha' + 2\gamma') - 2\mu\gamma'(2\alpha' + \gamma')] = 0 \quad (12)$$

$$\mu(\rho - 2e^{-2(\alpha+2\gamma+\lambda)}[\lambda'(\alpha' + 2\gamma') + \gamma'(2\alpha' + \gamma') - 2\gamma'' - \lambda'']) - 4(\mu + 4\pi)P = 0 \quad (13)$$

$$e^{2(\alpha+2\gamma+\lambda)}[\mu\rho - 4(\mu + 4\pi)P] + 2\mu[\alpha'' - \alpha'(2\gamma' + \lambda') + \gamma'' - \gamma'(\gamma' + 2\lambda') + \lambda''] = 0, \quad (14)$$

$$\mu(\rho - 2e^{-2(\alpha+2\gamma+\lambda)}[-\alpha'' + \lambda'(\alpha' + 2\gamma') + \gamma'(2\alpha' + \gamma') - 2\gamma'']) - 4(\mu + 4\pi)P = 0, \quad (15)$$

where the prime stands for $\frac{d}{d\tau}$.

It is to be noted that there are five unknowns α , γ , λ , P , and ρ involved in the above four equations. Therefore, for obtaining exact solutions of Equations (12)–(15), we need to consider some interplaying relationships between any two parameters, such as [69–71]

$$\lambda = m\gamma, \quad (16)$$

where $m \neq 0$ is a parameter, the value of which can be chosen suitably depending on the physical situation.

Now, solving Equations (12)–(15), we obtain the solutions as

$$\gamma(\tau) = k_1\tau + k_2, \quad (17)$$

$$\lambda(\tau) = m(k_1\tau + k_2), \quad (18)$$

$$\alpha(\tau) = k_3\tau + k_4, \quad (19)$$

$$\rho = -\frac{2\mu(3\mu + 8\pi)(c_1 e^{-2\tau c_2})}{(\mu + 8\pi)(5\mu + 16\pi)}, \quad (20)$$

$$P = -\frac{4\mu(\mu + 4\pi)(c_1 e^{-2\tau c_2})}{(\mu + 8\pi)(5\mu + 16\pi)}, \quad (21)$$

where k_1 , k_2 , k_3 , and k_4 are the integration constant and $c_1 = k_1[2k_1m + k_1 + k_3(m + 2)]$ and $c_2 = k_1(m + 2) + k_3$. Without loss of generality, we take $k_2 = k_4 = 0$.

3.2. $f(R, T) = R + \mu R^2 + \mu T$

In this case, we consider $f_1(R) = R + \mu R^2$ and $f_2(T) = \mu T$. Therefore, Equation (3) becomes

$$G_{ij} + 2\mu R R_{ij} - \frac{1}{2}\mu R^2 g_{ij} + 2\mu(g_{ij}\square - \nabla_i \nabla_j)R = (8\pi + \mu)T_{ij} + \left(P + \frac{T}{2}\right)\mu g_{ij}. \quad (22)$$

For (2), the field equations in $f(R, T)$ theory using (11) and (22) become

$$\begin{aligned} & -2(\gamma')^2(8\mu(7\alpha'' + 12\gamma'' + 7\lambda'' - 12(\lambda')^2) + e^{2(\alpha+2\gamma+\lambda)}) + \\ & 16\mu(\alpha')^3(2\gamma' + \lambda') + 4\mu(2\lambda'(\alpha^{(3)} + 2\gamma^{(3)} + \lambda^{(3)}) + (\alpha'' + 2\gamma'' + \lambda'')^2 - \\ & 2(\lambda')^2(3\alpha'' + 6\gamma'' + 2\lambda'')) + 4\gamma'(4\mu(\alpha^{(3)} + 2\gamma^{(3)} + \lambda^{(3)}) - \\ & \lambda'(4\mu(8\alpha'' + 15\gamma'' + 7\lambda'') + e^{2(\alpha+2\gamma+\lambda)}) + 8\mu(\lambda')^3) + \\ & 4\mu(\alpha')^2(-2(2\alpha'' + 6\gamma'' + 3\lambda'') + 52\gamma'\lambda' + 48(\gamma')^2 + 11(\lambda')^2) + \\ & 2\alpha'(4\mu(\alpha^{(3)} - \lambda'(7\alpha'' + 16\gamma'' + 7\lambda'') + \\ & 2\gamma'(-7\alpha'' - 15\gamma'' - 8\lambda'' + 13(\lambda')^2) + 2\gamma^{(3)} + 59(\gamma')^2\lambda' + \\ & 30(\gamma')^3 + \lambda^{(3)} + 2(\lambda')^3) - e^{2(\alpha+2\gamma+\lambda)}(2\gamma' + \lambda')) + 240\mu(\gamma')^3\lambda' + \\ & 76\mu(\gamma')^4 + e^{4(\alpha+2\gamma+\lambda)}(-3\mu\rho + 2\mu P - 16\rho\pi) = 0, \end{aligned} \quad (23)$$

$$\begin{aligned} & -2(\gamma')^2(4\mu(34\alpha'' + 54\gamma'' + 35\lambda'' - 38(\lambda')^2) + e^{2(\alpha+2\gamma+\lambda)}) + \\ & 48\mu(\alpha')^3(2\gamma' + \lambda') + 4\gamma'(4\mu(5\alpha^{(3)} + 9\gamma^{(3)} + 5\lambda^{(3)}) - \\ & \lambda'(72\mu\alpha'' + e^{2(\alpha+2\gamma+\lambda)} + 136\mu\gamma'' + 60\mu\lambda'') + 16\mu(\lambda')^3) + \\ & 4\mu(\alpha')^2(3(4\gamma' + \lambda')(8\gamma' + 7\lambda') - 2(6\alpha'' + 11(2\gamma'' + \lambda''))) + \\ & 4\mu(-2(\alpha^{(4)} + 2\gamma^{(4)} + \lambda^{(4)}) + 2\lambda'(5\alpha^{(3)} + 10\gamma^{(3)} + 4\lambda^{(3)}) + \\ & 12\alpha''(2\gamma'' + \lambda'') - 8(\lambda')^2(2\alpha'' + 4\gamma'' + \lambda'') + 3(\alpha'')^2 + 28\gamma''\lambda'' + \\ & 24(\gamma'')^2 + 5(\lambda'')^2) + 2\alpha'(4\mu(5\alpha^{(3)} - \lambda'(17\alpha'' + 44\gamma'' + 17\lambda'') + \\ & \gamma'(-34\alpha'' - 78\gamma'' - 44\lambda'' + 46(\lambda')^2) + 12\gamma^{(3)} + 103(\gamma')^2\lambda' + \\ & 54(\gamma')^3 + 6\lambda^{(3)} + 4(\lambda')^3) - e^{2(\alpha+2\gamma+\lambda)}(2\gamma' + \lambda'))2e^{2(\alpha+2\gamma+\lambda)}(2\gamma'' + \lambda'') + + \\ & 400\mu(\gamma')^3\lambda' + 132\mu(\gamma')^4 + e^{4(\alpha+2\gamma+\lambda)}(\mu\rho - 4P(\mu + 4\pi)) = 0, \end{aligned} \quad (24)$$

$$\begin{aligned} & 2e^{2(\alpha+2\gamma+\lambda)}(\alpha'' + \gamma'' + \lambda'') - 2(\gamma')^2(4\mu(41\alpha'' + 63\gamma'' + 41\lambda'' - 42(\lambda')^2) + \\ & e^{2(\alpha+2\gamma+\lambda)}) + 32\mu(\alpha')^3(2\gamma' + \lambda') + 4\gamma'(2\mu(11\alpha^{(3)} + 20\gamma^{(3)} + 11\lambda^{(3)}) - \\ & \lambda'(82\mu\alpha'' + e^{2(\alpha+2\gamma+\lambda)} + 144\mu\gamma'' + 64\mu\lambda'') + 16\mu(\lambda')^3) + \\ & 4\mu(\alpha')^2(-8(\alpha'' + 4\gamma'' + 2\lambda'') + 88\gamma'\lambda' + 84(\gamma')^2 + 17(\lambda')^2) + \\ & 4\mu(-2(\alpha^{(4)} + 2\gamma^{(4)} + \lambda^{(4)}) + 2\lambda'(5\alpha^{(3)} + 10\gamma^{(3)} + 4\lambda^{(3)}) + \\ & 2\alpha''(13\gamma'' + 7\lambda'') - 8(\lambda')^2(2\alpha'' + 4\gamma'' + \lambda'') + 5(\alpha'')^2 + 26\gamma''\lambda'' + \\ & 20(\gamma'')^2 + 5(\lambda'')^2) + 2\alpha'(4\mu(4\alpha^{(3)} - \lambda'(15\alpha'' + 37\gamma'' + 15\lambda'') + \\ & \gamma'(-32\alpha'' - 72\gamma'' - 41\lambda'' + 44(\lambda')^2) + 10\gamma^{(3)} + 105(\gamma')^2\lambda' + \\ & 60(\gamma')^3 + 5\lambda^{(3)} + 4(\lambda')^3) - e^{2(\alpha+2\gamma+\lambda)}(2\gamma' + \lambda')) + \\ & 480\mu(\gamma')^3\lambda' + 164\mu(\gamma')^4 + e^{4(\alpha+2\gamma+\lambda)}(\mu\rho - 4P(\mu + 4\pi)) = 0, \end{aligned} \quad (25)$$

$$\begin{aligned}
& 2e^{2(\alpha+2\gamma+\lambda)}(\alpha'' + 2\gamma'') - 2(\gamma')^2(4\mu(35\alpha'' + 54\gamma'' + 34\lambda'' - 48(\lambda')^2) + e^{2(\alpha+2\gamma+\lambda)}) + \\
& 32\mu(\alpha')^3(2\gamma' + \lambda') + 4\gamma'(4\mu(5\alpha^{(3)} + 9\gamma^{(3)} + 5\lambda^{(3)}) - \\
& \lambda'(88\mu\alpha'' + e^{2(\alpha+2\gamma+\lambda)} + 156\mu\gamma'' + 68\mu\lambda'') + 24\mu(\lambda')^3) + \\
& 4\mu(\alpha')^2(-8(\alpha'' + 4\gamma'' + 2\lambda'') + 92\gamma'\lambda' + 76(\gamma')^2 + 21(\lambda')^2) + \\
& 4\mu(-2(\alpha^{(4)} + 2\gamma^{(4)} + \lambda^{(4)}) + 2\lambda'(6\alpha^{(3)} + 12\gamma^{(3)} + 5\lambda^{(3)}) + \\
& 4\alpha''(7\gamma'' + 3\lambda'') - 2(\lambda')^2(11\alpha'' + 22\gamma'' + 6\lambda'') + \\
& 5(\alpha'')^2 + 3(8\gamma''\lambda'' + 8(\gamma'')^2 + (\lambda'')^2)) + 2\alpha'(4\mu(4\alpha^{(3)} - \lambda'(17\alpha'' + 44\gamma'' + 17\lambda'') + \\
& \gamma'(-30\alpha'' - 68\gamma'' - 36\lambda'' + 54(\lambda')^2) + 10\gamma^{(3)} + 103(\gamma')^2\lambda' + \\
& 50(\gamma')^3 + 5\lambda^{(3)} + 6(\lambda')^3) - e^{2(\alpha+2\gamma+\lambda)}(2\gamma' + \lambda')) + \\
& 432\mu(\gamma')^3\lambda' + 132\mu(\gamma')^4 + e^{4(\alpha+2\gamma+\lambda)}(\mu\rho - 4P(\mu + 4\pi)) = 0 \quad (26)
\end{aligned}$$

where the powers (3) and (4) denote the third- and fourth-order derivative w.r.t. τ .

We note that, in this case, the field Equations (23)–(26) yield the same solution (17)–(19), where $k_1 = \pm k_3$ and $m = \pm 1$. However, the other two physical parameters become $P = \rho = 0$.

3.3. $f(R, T) = R\mu + RT\mu^2$

We consider here the third case, i.e., $f(R, T) = f_1(R) + f_2(R)f_3(T)$, where $f_1(R) = f_2(R) = \mu R$ and $f_3(T) = \mu T$.

Now, Equation (3) becomes

$$G_{ij} + \mu^2(g_{ij}\square - \nabla_i \nabla_j)T = \frac{\mu PR g_{ij}}{\mu T + 1} + \frac{T_{ij}(\mu^2 R + 8\pi)}{\mu(\mu T + 1)}. \quad (27)$$

Therefore, for Equation (2), the field equations in $f(R, T)$ theory using Equations (11) and (27) are

$$\begin{aligned}
& -e^{2(\alpha+2\gamma+\lambda)}(\mu + 16\rho\pi) - 2(\mu(2\mu(P + \rho) - 1) + 1)(\alpha'' + 2\gamma'' + \lambda'') + \\
& 2\alpha'(2\gamma' + \lambda')(\mu(\mu\rho + 6\mu P - 2) + 1) + 4\gamma'\lambda'(\mu(\mu\rho + 6\mu P - 2) + 1) + \\
& 2(\gamma')^2(\mu(\mu\rho + 6\mu P - 2) + 1) = 0, \quad (28)
\end{aligned}$$

$$\begin{aligned}
& 2(\alpha'' + 2\gamma'' + \lambda'' - \mu(\alpha'' + \mu(4P - \rho)(2\gamma'' + \lambda'')) + \\
& \alpha'(2\gamma' + \lambda')(\mu^2(4P - \rho) - 1) + 2\gamma'\lambda'(\mu^2(4P - \rho) - 1) + \\
& (\gamma')^2(\mu^2(4P - \rho) - 1)) + e^{2(\alpha+2\gamma+\lambda)}(\mu - 16P\pi) = 0, \quad (29)
\end{aligned}$$

$$\begin{aligned}
& 2(\alpha'' + 2\gamma'' + \lambda'' + \mu(-\gamma'' - \mu(4P - \rho)(\alpha'' + \gamma'' + \lambda'')) + \\
& \alpha'(2\gamma' + \lambda')(\mu^2(4P - \rho) - 1) + 2\gamma'\lambda'(\mu^2(4P - \rho) - 1) + \\
& (\gamma')^2(\mu^2(4P - \rho) - 1)) + e^{2(\alpha+2\gamma+\lambda)}(\mu - 16P\pi) = 0, \quad (30)
\end{aligned}$$

$$\begin{aligned}
& -2(\mu - 1)\lambda'' + 2(\mu^2(4P - \rho) - 1)(-\alpha'' + \alpha'(2\gamma' + \lambda') - 2\gamma'' + 2\gamma'\lambda' + (\gamma')^2) + \\
& e^{2(\alpha+2\gamma+\lambda)}(\mu - 16P\pi) = 0. \quad (31)
\end{aligned}$$

In this case also, the field Equations (28)–(31) admit the same solutions set (17)–(19) as obtained in Section 3.1; however, the rest of the parameters are

$$\rho = \frac{c_1 k_1 (2c_1 k_1 (4\mu - 5)\mu^2 + e^{2c_2 \tau} (5\mu^3 - 16\mu\pi + 8\pi)) - 4\mu\pi e^{4c_2 \tau}}{10c_1 k_1 \mu^2 (c_1 k_1 \mu^2 - 4\pi e^{2c_2 \tau}) + 64\pi^2 e^{4c_2 \tau}}, \quad (32)$$

$$P = \frac{c_1 k_1 (c_1 k_1 \mu^3 - 4\pi e^{2c_2 \tau}) + 2\mu\pi e^{4c_2 \tau}}{5c_1 k_1 \mu^2 (c_1 k_1 \mu^2 - 4\pi e^{2c_2 \tau}) + 32\pi^2 e^{4c_2 \tau}}. \quad (33)$$

The expression for the ratio of the pressure and density (i.e., the EOS parameter w) is

$$w = \frac{2(c_1 k_1 (c_1 k_1 \mu^3 - 4\pi e^{2c_2 \tau}) + 2\mu \pi e^{4c_2 \tau})}{c_1 k_1 (2c_1 k_1 (4\mu - 5)\mu^2 + e^{2c_2 \tau} (5\mu^3 - 16\mu\pi + 8\pi)) - 4\mu \pi e^{4c_2 \tau}}. \quad (34)$$

In the above-mentioned three subsections we have shown variations of different model parameters in Figures 1–4 which are satisfactory as far as physical features are concerned.




Figure 1. Variation of the pressure and density w.r.t. time (Case 3.1). Here, we considered the following parametric values: $k_1 = 0.13$, $k_3 = 0.1$, and $m = 0.5$, which will also be followed in all other plots.




Figure 2. Variation of (a) $\rho - P$, (b) $\rho + P$, and (c) $\rho + 3P$ w.r.t. time (Case 3.1).




Figure 3. Variation of the pressure and density w.r.t. time (Case 3.3).




Figure 4. Variation of (a) $\rho - P$, (b) $\rho + P$, and (c) $\rho + 3P$ w.r.t. time (Case 3.3).

Hence, the five-dimensional cosmological model in the $f(R, T)$ theory of gravity corresponding to the solutions of Sections 3.1–3.3 can be uniquely presented as

$$dS^2 = d\tau^2 - e^{2k_3\tau}dX^2 - e^{2k_1\tau}(dY^2 + dZ^2) - e^{2mk_1\tau}dN^2. \quad (35)$$

4. Some Physical and Geometrical Properties

In this section, we study some physical and geometrical properties of the models obtained in the preceding subsections under the five-dimensional cosmological model in the $f(R, T)$ theory of gravity.

4.1. Status of the Model

The spatial volume (V), scalar expansion (θ), Hubble parameter (H), shear scalar (σ^2), and redshift (z) for the model are given by

$$V = \sqrt{g} = AB^2F = e^{c_2\tau}, \quad (36)$$

$$\theta = \frac{A_t}{A} + \frac{2B_t}{B} + \frac{F_t}{F} = c_2e^{-c_2\tau}, \quad (37)$$

$$H = \frac{1}{4} \left(\frac{A_t}{A} + \frac{2B_t}{B} + \frac{F_t}{F} \right) = \frac{1}{4}c_2e^{-c_2\tau}, \quad (38)$$

$$\begin{aligned} \sigma^2 &= \frac{1}{2} \left(-\frac{1}{4} \left(\frac{A_t}{A} + \frac{2B_t}{B} + \frac{F_t}{F} \right)^2 + \frac{(A_t)^2}{A^2} + \frac{2(B_t)^2}{B^2} + \frac{(F_t)^2}{F^2} \right) \\ &= \frac{1}{8}e^{-2c_2\tau} (k_1^2(m(3m-4)+4) - 2k_1k_3(m+2) + 3k_3^2), \end{aligned} \quad (39)$$

$$z = \frac{1}{\sqrt[4]{AB^2F}} - 1 = \frac{1}{\sqrt[4]{e^{c_2\tau}}} - 1, \quad (40)$$

$$\frac{\sigma^2}{\theta^2} = \frac{k_1^2(m(3m-4)+4) - 2k_1k_3(m+2) + 3k_3^2}{8c_2}. \quad (41)$$

From the above solution set, we notice that at $\tau = 0$, $V = 1$ and, as $\tau \rightarrow \infty$, $V \rightarrow \infty$. Therefore, it can be inferred that our model is free from the initial singularity. We also note that the pressure and density are finite at $\tau = 0$, which decrease as τ increases and tend to zero when $\tau \rightarrow \infty$. This means at infinite time that our model leads to a vacuum model. Further, as $\frac{\sigma^2}{\theta^2} \neq 0$, so our model is anisotropic throughout the evolution. Equation (40)

exhibits the expansion of the spacetime in the universe, when $\tau \rightarrow \infty$; however, in the present model,

$$q = - \frac{4 \left(\frac{A_{tt}}{A} + \frac{1}{4} \left(\frac{A_t}{A} + \frac{2B_t}{B} + \frac{F_t}{F} \right)^2 - \frac{(A_t)^2}{A^2} + \frac{2B_{tt}}{B} - \frac{2(B_t)^2}{B^2} + \frac{F_{tt}}{F} - \frac{(F_t)^2}{F^2} \right)}{\left(\frac{A_t}{A} + \frac{2B_t}{B} + \frac{F_t}{F} \right)^2} = 3,$$

which means that the universe is in a decelerating phase.

In Case 3.2, the model degenerates to a pure vacuum model in the $f(R, T)$ theory of gravity. We have shown different features of the model in Table 1.

Table 1. Features of the model.

For Case 3.2	$k_1 = k_3, m = 1$	$k_1 = k_3, m = -1$	$k_1 = -k_3, m = 1$	$k_1 = -k_3, m = -1$
V	expanding	expanding	decreasing	constant
θ	decreasing	decreasing	negative	0
q	3	3	3	undefined
z	decreasing	decreasing	increasing	0
$\frac{\sigma^2}{\theta^2}$	0	$\frac{3}{8}$	$\frac{3}{8}$	undefined

4.2. Stability of the Model

The stability of the model is obtained by considering the ratio $\frac{dp}{d\rho}$, which can be shown as equivalent to C_s^2 . If C_s^2 is positive, then the model is stable, whereas if C_s^2 is negative, the model is unstable. In our case, $\frac{dp}{d\rho} = 1 - \frac{\mu}{3\mu+8\pi}$. From this relation, we notice that C_s^2 is positive for $\mu > -4\pi$ and, thus, provides a stable model under this restrictive condition for Case 3.1. Under the same condition, the model for Case 3.3 (Figure 5) exhibits a stable model throughout the evolution.




Figure 5. $\frac{dp}{d\rho}$ of Case 3.3.

4.3. EOS Parameter (w)

In the present model, the EOS parameter is governed by the parameter μ . In Figure 6, one can note that different values of the parameter lead to a different model in $f(R, T)$ gravity. Caldwell and coworkers [72,73] pointed out that $w < -1$ is a better fit for the observed astrophysical data.




Figure 6. w of (a) Case 3.1 and (b) Case 3.3.

5. Discussion and Conclusions

In the present work, our motivation was to obtain exact solutions to the Einstein field equations for $5d$ locally rotationally symmetric (LRS) Bianchi type-I spacetime in the $f(R, T)$ theory of gravity. We presented cosmological models under the following three specifications: (i) $f(R, T) = \mu(R + T)$, (ii) $f(R, T) = R\mu + RT\mu^2$, and (iii) $f(R, T) = R + \mu R^2 + \mu T$.

The solution set under these models, via the graphical plots (Figure 6), exhibits that the EOS parameter w is completely governed by μ . Fine-tuning of the parameter μ provides the effect of phantom cosmology. Moreover, we were able to obtain a stable model of the universe by imposing restrictions upon this parameter.

In this connection, we also would like to mention here that Figures 1–5 are self-explanatory, which in a similar way, depict various physical properties of the model, e.g., in Figure 1, we exhibit the variation of the pressure and density w.r.t. time for different parametric values (say $k_1 = 0.13$, $k_3 = 0.1$, and $m = 0.5$).

We can explore various energy conditions, i.e., the null energy condition (NEC), $\rho + P \geq 0$, dominant energy condition (DEC), $\rho - P \geq 0$, weak energy condition (WEC) $\rho \geq 0$, and strong energy condition (SEC) $\rho + 3P \geq 0$ from Figures 1–4. Here, the NEC represents the attractive nature of gravity, which is true in Case 3.1, whereas it is violated in Case 3.3. According to the DEC, in Cases 3.1 and 3.3, the mass–energy parameter cannot be seen moving faster than light. The WEC is true in Case 3.1 and Case 3.3, which show that that the matter density observed by the respective observers is always positive. In Case 3.1, the SEC is satisfied, while it is violated in Case 3.3.

Some other salient and characteristics features of the cosmological models are as follows:

- (1) We notice that the model is free from the initial singularity and, hence, physically viable. This feature is obvious, as for $\tau = 0$, we obtain $V = 1$, and for $\tau \rightarrow \infty$, one can obtain $V \rightarrow \infty$.
- (2) The cosmic distribution has a finite fluid pressure and matter density at $\tau = 0$. The physical quantities decrease as τ increases and tend to zero when $\tau \rightarrow \infty$. Thus, our presented model leads to a vacuum cosmological solution at infinite time.
- (3) As $\frac{\sigma^2}{\rho^2} \neq 0$, so the model is anisotropic throughout the evolution. Again, $\tau \rightarrow \infty$ exhibits the expanding universe. However, $q = 3$ dictates that the universe is decelerating.
- (4) The stability of the model is obtained by considering the ratio $\frac{dp}{d\rho}$, which is positive for $\mu > -4\pi$, to yield a stable model.
- (5) The EOS parameter is governed by the parameter μ , and its value can be found as $\mu < -3.2\pi$. This is related to $w < -1$, which behaves like a phantom-energy-inspired cosmology. This type of phantom cosmology allows us to account for the dynamics

and matter content of the universe, tracing back the evolution to the inflationary epoch [74]. In this connection, we would also like to point out that while the dependence of μ is explicit across all cases, this is not overall true, as this situation is solely visible in the results from Case 3.1. One can note that Case 3.3 shows a clear time dependence (and therefore, very dependent on the magnitude of c_2).

- (6) The anisotropic/isotropic behavior of the models for different choices of the parameters are given in Table 1 in connection with Case 3.2.

Thus, an obvious issue is here: how to incorporate an accelerating phase of the universe, which is the present cosmological scenario, along with the decelerating phase in our phantom-type dark energy model. However, following Capozziello et al. [75], one can make an endeavor to obtain a transition from the deceleration to the acceleration phase of the universe. Therefore, this issue can be addressed in a future project.

Author Contributions: Conceptualization, R.R.S.; writing—original draft preparation, K.L.M.; writing—review and editing, S.R. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Data Availability Statement: Not applicable.

Acknowledgments: The authors would like to express gratitude to the Referees for their constructive comments for the improvement of the paper. S.R. is thankful to the authority of IUCAA, Pune, for providing the Visiting Associateship, under which a part of this work has been performed. Furthermore, thanks are due to Ashutosh Singh for his technical assistance.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Minkowski, H. Die Grundgleichungen für die Elektromagnetischen Vorgänge in bewegten Körpern. *Wissenschaften zu Göttingen, Mathematisch-Physikalische* **1908**, *1908*, 53–111.
- Minkowski, H. Jahresbericht der Deutschen Mathematiker-Vereinigung. *Raum und Zeit* **1909**, *18*, 75–88.
- Landau, L.D.; Lifshitz, E.M. *The Classical Theory of Fields. Course of Theoretical Physics*, 4th ed.; Butterworth-Heinemann: Oxford, UK, 2002; Volume 2.
- Kaluza, T. Zum Unitätsproblem in der Physik. *Sitz. Preuss. Akad Wiss.* **1921**, *33*, 966–972.
- Klein, O. Quantentheorie und fünfdimensionale Relativitätstheorie. *Z. Phys.* **1926**, *37*, 895–906. [[CrossRef](#)]
- Chodos, A.; Detweiler, S. Where has the fifth dimension gone? *Phys. Rev. D* **1980**, *21*, 2167–2170. [[CrossRef](#)]
- Alvarez, E.; Gavela, M.B. Entropy from Extra Dimensions. *Phys. Rev. Lett.* **1983**, *51*, 931–934. [[CrossRef](#)]
- Marciano, W.J. Time Variation of the Fundamental “Constants” and Kaluza–Klein Theories. *Phys. Rev. Lett.* **1984**, *52*, 489–491. [[CrossRef](#)]
- Gegenberg, J.D.; Das, A. Five-dimensional cosmological models with massless scalar fields. *Phys. Lett.* **1985**, *112A*, 427–430. [[CrossRef](#)]
- Lorenz-Petzold, D. Higher-dimensional extensions of Bianchi type-I cosmologies. *Phys. Rev.* **1985**, *31*, 929–931. [[CrossRef](#)]
- Wesson, P.S. Astrophysical data and cosmological solutions of a Kaluza–Klein theory of gravity. *Astron. Astrophys.* **1986**, *166*, 1–3.
- Grøn, Ø. Inflationary cosmology according to Wesson’s gravitational theory. *Astron. Astrophys.* **1988**, *193*, 1–4.
- Wesson, P. *5-Dimensional Physics: Classical and Quantum Consequences of Kaluza Klein Cosmology*; World Scientific Publishing Company: Singapore, 2006.
- Abbott, R.B.; Barr, S.M.; Ellis, S.D. Kaluza–Klein cosmologies and inflation. *Phys. Rev. D* **1984**, *30*, 720–727. [[CrossRef](#)]
- Sahdev, D. Towards a realistic Kaluza–Klein cosmology. *Phys. Lett.* **1984**, *137B*, 155–159. [[CrossRef](#)]
- Chatterjee, S.; Bhui, B. Homogeneous cosmological model in higher dimension. *Mon. Not. R. Astron. Soc.* **1990**, *247*, 57–61.
- Bergamini, R.; Orzalesi, C. Towards a cosmology for multidimensional unified theories. *Phys. Lett. B* **1984**, *135*, 38–42. [[CrossRef](#)]
- Ishihara, H. Kaluza–Klein Inflation. *Prog. Theor. Phys.* **1984**, *72*, 376–378. [[CrossRef](#)]
- Banerjee, A.; Bhui, B.; Chatterjee, S. Bianchi type I cosmological models in higher dimensions. *Astrophys. J.* **1990**, *358*, 23–27. [[CrossRef](#)]
- Chatterjee, S.; Bhui, B.; Basu, M.B.; Banerjee, A. Inhomogeneous model with a cosmological constant. *Phys. Rev. D* **1994**, *50*, 2924–2927. [[CrossRef](#)]
- Banerjee, A.; Panigrahi, D.; Chatterjee, S. A class of inhomogeneous cosmological models in Kaluza–Klein spacetime. *Class. Quantum Gravit.* **1994**, *11*, 1405–1414. [[CrossRef](#)]
- Banerjee, A.; Das, A.; Panigrahi, D. Singularity-free cylindrically symmetric solutions in Kaluza–Klein space-time. *Phys. Rev. D* **1995**, *51*, 6816–6820. [[CrossRef](#)]

23. Ledesma, D.S.; Bellini, M. Single field inflationary models with non-compact Kaluza–Klein theory. *Phys. Lett. B* **2004**, *581*, 1–8. [\[CrossRef\]](#)
24. Aguilar, J.E.M.; Bellini, M. Noncompact Kaluza–Klein theory and inflationary cosmology: a complete formalism. *Phys. Lett. B* **2004**, *596*, 116–122. [\[CrossRef\]](#)
25. Bellini, M. Inflationary cosmology from STM theory of gravity. *Nucl. Phys. B* **2003**, *660*, 389–400. ; Erratum: *Nucl. Phys. B* **2003**, *671*, 501. [\[CrossRef\]](#)
26. Liko, T.; Wesson, P.S. An exact solution of the five-dimensional Einstein equations with four-dimensional de Sitter-like expansion. *J. Math. Phys.* **2005**, *46*, 062504. [\[CrossRef\]](#)
27. Mukhopadhyay, U.; Ghosh, P.P.; Ray, S. Higher dimensional dark energy investigation with variable Λ and G . *Int. J. Theor. Phys.* **2010**, *49*, 1622–1627. [\[CrossRef\]](#)
28. Sharif, M.; Khanum, F. Kaluza–Klein cosmology with varying G and Λ . *Astrophys. Space Sci.* **2011**, *334*, 209–214. [\[CrossRef\]](#)
29. Pahwa, I. Higher dimensional cosmological models: An alternative explanation for late time cosmic acceleration. *J. Phys. Conf. Ser.* **2014**, *484*, 012045. [\[CrossRef\]](#)
30. Perlmutter, S.; Aldering, G.; Della Valle, M.; Deustua, S.; Ellis, R.S.; Fabbro, S.; Fruchter, A.; Goldhaber, G.; Groom, D.E.; Hook, I.M.; et al. Discovery of a supernova explosion at half the age of the Universe. *Nature* **1998**, *391*, 51–54. [\[CrossRef\]](#)
31. Perlmutter, S.; Aldering, G.; Goldhaber, G.; Knop, R.A.; Nugent, P.; Castro, P.G.; Deustua, S.; Fabbro, S.; Goobar, A.; Groom, D.E.; et al. Measurements of Ω and Λ from 42 high-redshift supernovae. *Astrophys. J.* **1999**, *517*, 565–586. [\[CrossRef\]](#)
32. Knop, R.A.; Aldering, G.; Amanullah, R.; Astier, P.; Blanc, G.; Burns, M.S.; Conley, A.; Deustua, S.E.; Doi, M.; Ellis, R.; et al. New constraints on Ω_M , Ω_Λ , and w from an independent set of 11 high-redshift supernovae observed with the Hubble Space Telescope. *Astrophys. J.* **2003**, *598*, 102–137. [\[CrossRef\]](#)
33. Riess, A.G.; Filippenko, A.V.; Challis, P.; Clocchiatti, A.; Diercks, A.; Garnavich, P.M.; Gilliland, R.L.; Hogan, C.J.; Jha, S.; Kirshner, R.P.; et al. Observational evidence from supernovae for an accelerating Universe and a cosmological constant. *Astron. J.* **1998**, *116*, 1009–1038. [\[CrossRef\]](#)
34. Riess, A.G.; Strolger, L.-G.; Tonry, J.; Casertano, S.; Ferguson, H.C.; Mobasher, B.; Challis, P.; Filippenko, A.V.; Jha, S.; Li, W.; et al. Type Ia supernova discoveries at $z > 1$ from the Hubble Space Telescope: Evidence for past deceleration and constraints on dark energy evolution. *Astron. J.* **2004**, *607*, 665–687. [\[CrossRef\]](#)
35. Tonry, J.L.; Schmidt, B.P.; Barris, B.; Candia, P.; Challis, P.; Clocchiatti, A.; Coil, A.L.; Filippenko, A.V.; Garnavich, P.; Hogan, C.; et al. Cosmological results from high- z supernovae. *Astron. J.* **2003**, *594*, 1–24. [\[CrossRef\]](#)
36. Valentino, E.D.; Mena, O.; Pan, S.; Visinelli, L.; Yang, W.; Melchiorri, A.; Mota, D.F.; Riess, A.G.; Silk, J. In the realm of the Hubble tension—a review of solutions. *Class. Quantum Grav.* **2021**, *38*, 153001. [\[CrossRef\]](#)
37. Valentino, E.D. Challenges of the standard cosmological model. *Universe* **2022**, *8*, 399. [\[CrossRef\]](#)
38. Harko, T.; Lobo, F.S.N.; Nojiri, S.; Odintsov, S.D. $f(R, T)$ gravity. *Phys. Rev. D* **2011**, *84*, 024020. [\[CrossRef\]](#)
39. Capozziello, S.; Laurentis, M.D. Extended theories of gravity. *Phys. Rep.* **2011**, *509*, 167–321. [\[CrossRef\]](#)
40. Adhav, K.S. LRS Bianchi type-I cosmological model in $f(R, T)$ theory of gravity. *Astrophys. Space Sci.* **2012**, *339*, 365–369. [\[CrossRef\]](#)
41. Alvarenga, F.G.; Houndjo, M.J.S.; Monwanou, A.V.; Orou, J.B.C. Testing some $f(R, T)$ gravity models from energy conditions. *J. Mod. Phys.* **2013**, *4*, 130–139. [\[CrossRef\]](#)
42. Mahanta, K.L. Bulk viscous cosmological models in $f(R, T)$ theory of gravity. *Astrophys. Space Sci.* **2014**, *353*, 683–689. [\[CrossRef\]](#)
43. Mishra, B.; Sahoo, P.K. Bianchi type VI_h perfect fluid cosmological model in $f(R, T)$ theory. *Astrophys Space Sci.* **2014**, *352*, 331–336. [\[CrossRef\]](#)
44. Cai, Y.-F.; Capozziello, S.; Laurentis, M.D.; Saridakis, E.N. $f(T)$ teleparallel gravity and cosmology. *Rept. Prog. Phys.* **2016**, *79*, 106901. [\[CrossRef\]](#) [\[PubMed\]](#)
45. Das, A.; Rahaman, F.; Guha, B.K.; Ray, S. Compact stars in $f(R, T)$ gravity. *Eur. Phys. J. C* **2016**, *76*, 654. [\[CrossRef\]](#)
46. Das, A.; Ghosh, S.; Guha, B.K.; Das, S.; Rahaman, F.; Ray, S. Gravastars in $f(R, T)$ gravity. *Phys. Rev. D* **2017**, *95*, 124011. [\[CrossRef\]](#)
47. Moraes, P.H.R.S.; Sahoo, P.K. Modeling wormholes in $f(R, T)$ gravity. *Phys. Rev. D* **2018**, *97*, 024007. [\[CrossRef\]](#)
48. Mishra, B.; Tripathy, S.K.; Tarai, S. Cosmological models with a hybrid scale factor in an extended gravity theory. *Mod. Phys. Lett. A* **2018**, *33*, 1850052. [\[CrossRef\]](#)
49. Deb, D.; Guha, B.K.; Rahaman, F.; Ray, S. Anisotropic strange stars under simplest minimal matter-geometry coupling in the $f(R, T)$ gravity. *Phys. Rev. D* **2018**, *97*, 084026. [\[CrossRef\]](#)
50. Deb, D.; Rahaman, F.; Ray, S.; Guha, B.K. Strange stars in $f(R, T)$ gravity. *J. Cosmol. Astropart. Phys.* **2018**, *03*, 044. [\[CrossRef\]](#)
51. Biswas, S.; Ghosh, S.; Guha, B.K.; Rahaman, F.; Ray, S. Strange stars in Krori–Barua spacetime under $f(R, T)$ gravity. *Ann. Phys.* **2019**, *401*, 1–20. [\[CrossRef\]](#)
52. Deb, D.; Ketov, S.V.; Khlopov, M.; Ray, S. Study on charged strange stars in $f(R, T)$ gravity. *J. Cosmol. Astropart. Phys.* **2019**, *10*, 070. [\[CrossRef\]](#)
53. Deb, D.; Ketov, S.V.; Maurya, S.K.; Khlopov, M.; Moraes, P.H.R.S.; Ray, S. Exploring physical features of anisotropic strange stars beyond standard maximum mass limit in $f(R, T)$ gravity. *Mon. Not. R. Astron. Soc.* **2019**, *485*, 5652–5665. [\[CrossRef\]](#)
54. Fisher, S.B.; Carlson, E.D. Reexamining $f(R, T)$ gravity. *Phys. Rev. D* **2019**, *100*, 064059. [\[CrossRef\]](#)
55. Biswas, S.; Shee, D.; Ray, S.; Guha, B.K. Anisotropic strange star with Tolman–Kuchowicz metric under $f(R, T)$ gravity. *Eur. Phys. J. C* **2020**, *80*, 175. [\[CrossRef\]](#)

56. Ray, S.; Sengupta, R.; Nimesh, H. Gravastar: An alternative to black hole. *Int. J. Mod. Phys. D* **2020**, *29*, 2030004. [[CrossRef](#)]
57. Hulke, N.; Singh, G.P.; Bishi, B.K.; Singh, A. Variable Chaplygin gas cosmologies in $f(R, T)$ gravity with particle creation, *New Astron.* **2020**, *77*, 101357. [[CrossRef](#)]
58. Mishra, B.; Esmeili, F.M.; Ray, S. Cosmological models with variable anisotropic parameter in $f(R, T)$ gravity. *Ind. J. Phys.* **2021**, *95*, 2245–2254. [[CrossRef](#)]
59. Varshney, G.; Sharma, U.K.; Pradhan, A.; Kumar, N. Reconstruction of Tachyon, Dirac-Born-Infeld-essence and Phantom model for Tsallis holographic dark energy in $f(R, T)$ gravity. *Chin. J. Phys.* **2021**, *73*, 56–73. [[CrossRef](#)]
60. Pradhan, A.; Garg, P.; Dixit, A. FRW cosmological models with cosmological constant in $f(R, T)$ theory of gravity. *Can. J. Phys.* **2021**, *99*, 741–753. [[CrossRef](#)]
61. Bhardwaj, V.K.; Pradhan, A. Evaluation of cosmological models in gravity in different dark energy scenario. *New Astron.* **2022**, *91*, 101675. [[CrossRef](#)]
62. Tiwari, R.K.; Beesham, A.; Mishra, S.; Dubey, V. Bianchi type I cosmological model in $f(R, T)$ gravity. *Phys. Sci. Forum* **2021**, *2*, 38. [[CrossRef](#)]
63. Tiwari, R.K.; Beesham, A.; Mishra, S.; Dubey, V. Anisotropic cosmological model in a modified theory of gravitation. *Universe* **2021**, *7*, 226. [[CrossRef](#)]
64. Bertolami, O.; Lobo, F.S.N.; Paramos, J. Nonminimal coupling of perfect fluids to curvature. *Phys. Rev. D* **2008**, *78*, 064036. [[CrossRef](#)]
65. Schutz, B.F. Perfect fluids in general relativity: Velocity potentials and a variational principle. *Phys. Rev. D* **1970**, *2*, 2762. [[CrossRef](#)]
66. Brown, J.D. Action functionals for relativistic perfect fluids. *Class. Quantum Gravit.* **1993**, *10*, 1579. [[CrossRef](#)]
67. Baffou, E.H.; Houndjo, M.J.S.; Rodrigues, M.E.; Kpadonou, A.V.; Tossa, J. Cosmological evolution in $f(R, T)$ theory with collisional matter. *Phys. Rev. D* **2015**, *92*, 084043. [[CrossRef](#)]
68. Mohanty, G.; Sahoo, R.R. Mesonic stiff fluid distribution in five dimensional Bianchi type space time. *Astrophys. Space Sci.* **2008**, *314*, 1–6. [[CrossRef](#)]
69. Collins, C.B.; Glass, E.N.; Wilkinson, D.A. Exact spatially homogeneous cosmologies. *Gen. Relativ. Gravit.* **1980**, *12*, 805–823. [[CrossRef](#)]
70. Moraes, P.H.R.S. Cosmology from induced matter model applied to 5D $f(R, T)$ theory. *Astrophys. Space Sci.* **2014**, *352*, 273–279. [[CrossRef](#)]
71. Borkar, M.S.; Ameen, A. Evaluation of Bianchi type VI₀ magnetized anisotropic dark energy models with constant deceleration parameter in bimetric theory of gravitation. *Int. J. Mod. Phys. D* **2015**, *24*, 1550019. [[CrossRef](#)]
72. Caldwell, R.R. A phantom menace? Cosmological consequences of a dark energy component with super-negative equation of state. *Phys. Lett. B* **2002**, *545*, 23–29. [[CrossRef](#)]
73. Caldwell, R.R.; Kamionkowski, M.; Weinberg, N.N. Phantom energy: Dark energy with $w < -1$ causes a cosmic doomsday. *Phys. Rev. Lett.* **2003**, *91*, 071301. [[CrossRef](#)] [[PubMed](#)]
74. Capozziello, S.; Nojiri, S.; Odintsov, S.D. Unified phantom cosmology: Inflation, dark energy and dark matter under the same standard. *Phys. Lett. B* **2006**, *632*, 597–604. [[CrossRef](#)]
75. Astashenok, A.V.; Nojiri, S.; Odintsov, S.D.; Yurov, A.V. Phantom cosmology without Big Rip singularity. *Phys. Lett. B* **2012**, *709*, 396–403. [[CrossRef](#)]