

Improved Model of Primordial Black Hole Formation after Starobinsky Inflation

Sultan Saburov¹ and Sergei V. Ketov^{1,2,3,*} 

- ¹ Interdisciplinary Research Laboratory, Tomsk State University, Tomsk 634050, Russia; saburov@mail.tsu.ru
² Department of Physics, Tokyo Metropolitan University, 1-1 Minami-ohsawa, Hachioji 192-0397, Japan
³ Kavli Institute for the Physics and Mathematics of the Universe (WPI), The University of Tokyo Institutes for Advanced Study, Chiba 277-8583, Japan
* Correspondence: ketov@tmu.ac.jp

Abstract: A new (improved) model of inflation and primordial black hole (PBH) formation is proposed by combining the Starobinsky model of inflation, Appleby–Battye–Starobinsky (ABS) model of dark energy, and a quantum correction in the modified $F(R)$ gravity. The energy scale parameter in the ABS model is taken to be close to the inflationary scale, in order to describe double inflation instead of dark energy. The quantum correction is given by the term quartic in the spacetime scalar curvature R with a negative coefficient ($-\delta$) in the $F(R)$ function. It is demonstrated that very good agreement (within 1σ) with current measurements of the cosmic microwave background (CMB) radiation can be achieved by choosing the proper value of δ , thus solving the problem of low values of the tilt of CMB scalar perturbations in the earlier proposed model in arXiv:2205.00603. A large (by a factor of 10^7 against CMB) enhancement in the power spectrum of scalar perturbations is achieved by fine tuning the parameters of the model. It is found by numerical analysis that it can lead to formation of asteroid-size PBHs with masses up to 10^{20} g, which may form dark matter in the current universe.

Keywords: inflation; primordial black holes; dark matter



Citation: Saburov, S.; Ketov, S.V. Improved Model of Primordial Black Hole Formation after Starobinsky Inflation. *Universe* **2023**, *9*, 323. <https://doi.org/10.3390/universe9070323>

Academic Editors: Galina L. Klimchitskaya, Vladimir M. Mostepanenko and Sergey V. Sushkov

Received: 15 June 2023
Revised: 29 June 2023
Accepted: 5 July 2023
Published: 6 July 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

This year is the centennial anniversary of Friedmann’s prediction for expanding universe, which was based on a nonstationary solution to Einstein’s equations. In recent years, the expanding universe was extended by other important features, such as cosmological inflation, dark matter, and dark energy.

A paradigm of inflation in the early universe, proposed a long time ago [1,2] as a possible solution to internal problems in theoretical cosmology (flatness, horizon, initial conditions, and structure formation), is currently well supported by precision measurements of the cosmic microwave background (CMB) radiation by WMAP and Planck satellite missions combined with recent BICEP/Keck data [3–5]. The original (1980) Starobinsky model of inflation [6] gives a simple theoretical description of inflation by using only gravitational interactions, which is in very good agreement with the current CMB measurements. It is, therefore, natural to extend the Starobinsky model of inflation by including more features, such as production of primordial black holes (PBH). The PBH formation may explain the origin of black holes and dark matter in the current universe [7–12].

In modern terms, see, e.g., Refs. [13–15] for more details, the Starobinsky model is a special case among modified gravity theories with the action:

$$S = \frac{M_{\text{Pl}}^2}{2} \int d^4x \sqrt{-g} F(R) . \quad (1)$$

The Starobinsky F -function of spacetime scalar curvature R is given by:

$$F_S(R) = R + \frac{R^2}{6M^2}, \tag{2}$$

where $M_{\text{Pl}} = (8\pi G)^{-1/2} = 2.435 \times 10^{18}$ GeV is the reduced Planck mass and M is the only (mass) parameter. The known CMB amplitude determines $M \approx 1.3 \times 10^{-5} M_{\text{Pl}}$, so that the Starobinsky model has no free parameters.

An $F(R)$ -gravity model is well known to be equivalent to the quintessence (scalar-tensor) gravity model in terms of the inflaton scalar field ϕ with the scalar potential $V(R(\phi))$ in the parametric form, see, e.g., Refs. [16,17] for a derivation:

$$V(R) = M_{\text{Pl}}^2 \frac{F'R - F}{2(F')^2}, \quad \phi(R) = \sqrt{\frac{3}{2}} M_{\text{Pl}} \ln F', \tag{3}$$

where the primes denote the derivatives with respect to the given variable (R). It is usually impossible to analytically derive the inverse function $R(\phi)$ from a given function $\phi(R)$, with the notable exception being the Starobinsky case (2), where one obtains a simple and well-known answer:

$$V_S(\phi) = \frac{3}{4} M_{\text{Pl}}^2 M^2 \left[1 - \exp\left(-\sqrt{\frac{2}{3}} \phi / M_{\text{Pl}}\right) \right]^2. \tag{4}$$

To form PBH out of a gravitational collapse of large (curvature) perturbations in the early universe, one needs a large enhancement of the power spectrum of scalar perturbations by six or seven orders of magnitude against the CMB spectrum. In the context of single-field models of inflation, it can be achieved in the double inflation scenario via engineering a near-inflection point in the inflaton scalar potential at lower (than inflation) scales [10,18]. The potential (4) does not have an inflection point and, hence, does not lead to PBH production. However, one can modify the Starobinsky $F(R)$ -gravity function (2) by extra terms that lead to an inflection point. This should be done in agreement with CMB observables, leading to constraints on eligible F -functions. Moreover, there are other conditions, such as absence of ghosts and singularities, and the correct Newtonian limit. Once all the necessary conditions are satisfied, one has to achieve the required enhancement of the power spectrum and generate PBH with the masses beyond the Hawking radiation limit of 10^{15} g because, otherwise, all those PBH would evaporate before the present times.

These problems were partially solved in Ref. [19] by adding to the $F(R)$ -gravity function (2) the additional terms known in the literature as the Appleby–Battye–Starobinsky (ABS) model of dark energy [20], and described by the F -function:

$$F_{\text{ABS}}(R) = (1 - g)R + gE_{\text{AB}} \ln \left[\frac{\cosh\left(\frac{R}{E_{\text{AB}}} - b\right)}{\cosh(b)} \right], \tag{5}$$

where the Appleby–Battye parameter E_{AB} has been introduced as:

$$E_{\text{AB}} = \frac{R_0}{2g \ln(1 + e^{2b})} \tag{6}$$

with the new dimensional scale R_0 and the dimensionless positive parameters g and b . The function (5) was carefully chosen in Ref. [20] in order to meet the no-ghost (stability) conditions in modified $F(R)$ gravity, which are given by:

$$F'(R) > 0 \quad \text{and} \quad F''(R) > 0, \tag{7}$$

it avoids singularities, obeys the Newtonian limit, and mimics a positive cosmological constant representing the dark energy for $R \gg R_0$ with proper values of the parameters g and b defining the shape of the scalar potential. To describe the present dark energy in the universe, the parameter R_0 representing the vacuum value of the scalar curvature should be very small, $\sqrt{R_0} \sim 10^{-33}$ eV and, hence, the AB-parameter should be small too.

The shape of the F -function (5), thus, leads to a meta-stable de Sitter vacuum after inflation and, hence, a near-inflection point in the potential. In Ref. [19], the dark energy value of $\sqrt{R_0}$ was replaced by a much higher value of the order M below the inflationary scale $H_{\text{inf.}} \sim 10^{14}$ GeV.¹ The new model with the F -function:

$$F_{\text{modified}}(R) = F_{\text{ABS}}(R) + \frac{R^2}{6M^2} \tag{8}$$

leads to the desired enhancement of the power spectrum of scalar perturbations and the formation of PBH with the asteroid-size masses of an order of 10^{19} g, which may form the whole dark matter in the present universe. In order to obtain these results, the parameters g and b were fine tuned, but agreement with the observed CMB tilt n_s of scalar perturbations was not good enough (outside 1σ but within 3σ lower) in Ref. [19].

In this communication, we propose the improved (new) model of Starobinsky inflation and PBH formation, having very good agreement (within 1σ) to CMB measurements. The new model is defined in Section 2 by combining phenomenological and theoretical considerations. Inflation is studied in Section 3. The power spectrum is derived in Section 4 together with the PBH masses. Section 5 contains our Conclusion.

2. The New Model

Our improved modified gravity model is (phenomenologically) defined by the F -function:

$$F(R) = (1 - g_1)R + gE_{AB} \ln \left[\frac{\cosh\left(\frac{R}{E_{AB}} - b\right)}{\cosh(b)} \right] + \frac{R^2}{6M^2} - \delta \frac{R^4}{48M^6} , \tag{9}$$

where we have changed the coefficient at the first term with another parameter $g_1 \neq g$ and have added the new term quartic in R with the new parameter δ . The AB parameter E_{AB} is still defined by Equation (6), where $R_0 = \beta M^2$ with yet another parameter β . All the parameters (g_1, g, b, β, δ) are dimensionless by definition.

The significance of each term in Equation (9) can be explained as follows.

The second term in Equation (9) becomes approximately linear in R both for small and large values of R/E_{AB} , and thus, correlates with the first term linear in R . Hence, in the low-curvature regime for small values of R/E_{AB} , consistency with gravity measurements inside the Solar system requires the Einstein–Hilbert effective action, which implies the following relation:

$$g_1 = -g \tanh b . \tag{10}$$

Starobinsky inflation is essentially governed by the third term quadratic in R in Equation (9), which leads to inflaton slow-roll (SR) in the high-curvature regime for the values of R/M^2 between 220 and 10 [17]. The scalar potential (4) of the Starobinsky model has an infinite plateau, which allows arbitrary duration of inflation, measured by a number N_e of e-folds. However, the Starobinsky inflation is unstable against quantum gravity corrections of the higher order in the scalar curvature. As was demonstrated in Refs. [21–23], the leading superstring-inspired quantum correction should be quartic in R , while it eliminates the infinite plateau in the Starobinsky potential and, hence, restricts the maximal number of e-folds. In order to be consistent with CMB measurements, the value of the δ -coefficient should be small enough, for instance, $|\delta| < 3.9 \times 10^{-6}$ according to Ref. [21].

It was assumed in Ref. [21] that the coefficient in front of the R^4 term is positive, which led to the inflaton scalar potential going down and hilltop inflation. In this paper, this coefficient is taken to be negative ($-\delta < 0$), which leads to the potential going up before inflation, see the next Section. The quartic term with $\delta > 0$ in Equation (9) is also responsible for raising the CMB values of the scalar tilt n_s in the improved model (9) against those in Ref. [19], as is demonstrated below. A similar effect was noticed in the modified E-type inflationary models of alpha-attractors and PBH formation, proposed in Ref. [24].

PBH formation may happen at the energy scales below the inflationary scale, which are governed by the parameter $\sqrt{R_0}$ of the order M . Hence, the parameter β in Equation (9) should be of the order one. The remaining parameters g and b are also of the order one, while their values should be tuned in order to generate a large peak $\sim 10^{-2}$ in the power spectrum of scalar perturbations. This can only be done numerically by scanning the parameter space, as in Refs. [19,24].

We found that in the model (9), the parameters (β, g, b) must be fine tuned to very specific values, namely:

$$\beta \approx 3.00, \quad g \approx 2.25 \quad \text{and} \quad b \approx 2.89, \tag{11}$$

because, otherwise, a peak in the power spectrum is either absent, too small, or too high.

It is easy to verify that the first and second derivatives of the F -function (9) are positive:

$$F'(R) = 1 + g \tanh(b) + g \tanh\left(\frac{R}{E_{AB}} - b\right) + \frac{R}{3M^2} - \frac{\delta R^3}{12M^6} > 0 \tag{12}$$

and

$$F''(R) = \frac{g}{E_{AB}} \operatorname{sech}^2\left(\frac{R}{E_{AB}} - b\right) + \frac{1}{3M^2} - \frac{\delta R^2}{4M^6} > 0, \tag{13}$$

for the given values of the parameters and the relevant values of $R < 240M^2$.

3. Inflaton Potential and Slow-Roll

According to Equations (3) and (9), the scalar potential $V(\phi)$ of the inflaton field ϕ in the parametric form (with the parameter R) is given by:

$$\begin{aligned} \frac{V(R)}{M_{\text{Pl}}^4} &= \frac{1}{2} e^{-2\sqrt{\frac{2}{3}}\phi/M_{\text{Pl}}} \times \\ &\times \left\{ gR \tanh\left(\frac{R}{E_{AB}} - b\right) - gE_{AB} \ln\left[\frac{\cosh\left(\frac{R}{E_{AB}} - b\right)}{\cosh(b)}\right] + \frac{R^2}{6M^2} - \frac{\delta R^4}{16M^6} \right\}, \end{aligned} \tag{14}$$

where

$$\exp\left(\sqrt{\frac{2}{3}}\frac{\phi}{M_{\text{Pl}}}\right) = 1 + g \tanh(b) + g \tanh\left(\frac{R}{E_{AB}} - b\right) + \frac{R}{3M^2} - \frac{\delta R^3}{12M^6}. \tag{15}$$

The function $\phi(R)$ in Equation (15) cannot be inverted in a useful analytic form, so we employ numerical calculations with Mathematica in what follows. The profiles of the inflaton potential $V(\phi)$ for the selected values of R_0 and δ are given in Figure 1.

The plot on the left-hand side of Figure 1 demonstrates that the potential has two plateaus at different scales; it is going up for $\delta > 0$ before inflation, while the value of the parameter δ determines the location where the potential goes up at the beginning of SR inflation. The height of the higher (Starobinsky) plateau is determined by M , the height of the lower plateau is controlled by R_0 , the length of the lower plateau is controlled by g , and the flatness of the lower plateau is controlled by b . The plot on the right-hand side of

Figure 1 shows a shallow meta-stable de Sitter minimum (dip), a near-inflection point, and a small bump (local maximum).




Figure 1. The inflaton potential for selected values of R_0 and δ at fixed $g = 2.25$ and $b = 2.89$ with $V_0 = \frac{3}{4} M_{\text{Pl}}^2 M^2$ (on the left). The potential for low values of ϕ / M_{Pl} are shown on the right.

The inflaton potential $V(\phi)$ for selected values of g at fixed $R_0 = 4M^2$ and $b = 2.89$ is displayed in Figure 2. This figure shows that duration of the second inflation (length of the lower plateau) is sensitive to the value of the parameter g . The longest lower plateau corresponds to $g \approx 3.0$.




Figure 2. The inflaton potential for selected values of g at fixed $R_0 = 4M^2$ and $b = 2.89$.

The SR conditions are given by smallness of the standard SR parameters, $\epsilon_{\text{sr}} \ll 1$ and $|\eta_{\text{sr}}| \ll 1$, where:

$$\epsilon_{\text{sr}}(\phi) = \frac{M_{\text{Pl}}^2}{2} \left(\frac{V'(\phi)}{V(\phi)} \right)^2 \quad \text{and} \quad \eta_{\text{sr}}(\phi) = M_{\text{Pl}}^2 \frac{V''(\phi)}{V(\phi)}. \tag{16}$$

A duration of inflation is defined by the number N of e-folds:

$$N = \int_{t_{\text{in}}}^{t_{\text{end}}} H(t) dt \approx \frac{1}{M_{\text{Pl}}^2} \int_{\phi_{\text{end}}}^{\phi_{\text{in}}} \frac{V(\phi)}{V'(\phi)} d\phi, \tag{17}$$

where $H(t)$ is the Hubble function. The CMB observable (tilt) n_s of scalar perturbations and the tensor-to-scalar ratio r are related to the values of the SR parameters at the horizon exit with the standard pivot scale $k_* = 0.05 \text{ Mpc}^{-1}$, which (in our model) is close to the scale at the beginning of SR inflation by $M_{\text{Pl}} \delta t \approx 2$ or $\delta N \approx 1$ or $\delta \phi / M_{\text{Pl}} \sim 10^{-2}$. Hence, ϕ_{in} can be identified with ϕ_{exit} at the horizon exit in a derivation of the CMB tilts, leading to:

$$n_s = 1 + 2\eta_{\text{sr}}(\phi_{\text{in}}) - 6\epsilon_{\text{sr}}(\phi_{\text{in}}) \quad \text{and} \quad r = 16\epsilon_{\text{sr}}(\phi_{\text{in}}). \tag{18}$$

The running SR parameters for selected values of R_0 and δ are displayed in Figure 3.




Figure 3. The SR parameter $\epsilon_{sr}(\phi)$ (on the left) and the SR parameter $\eta_{sr}(\phi)$ (on the right) for selected values of R_0 and δ at fixed $g = 2.25$ and $b = 2.89$. The end of Starobinsky inflation is reached at $\phi_{end} \approx 2.98M_{Pl}$ when $\epsilon_{sr}(\phi_{end}) \approx 1$.

It follows from Figure 3 that the peak in $\epsilon_{sr}(\phi)$ is sensitive to the value of R_0 , while the tails of $\eta_{sr}(\phi)$ are sensitive to the value of δ . Therefore, the value of δ affects the value of n_s , and then the value of r becomes weakly dependent upon δ , in agreement with Ref. [21].

The standard equations of motion of inflaton field $\phi(t)$ read:

$$\ddot{\phi} + 3H\dot{\phi} + V'(\phi) = 0, \quad 3H^2 = \frac{1}{M_{Pl}^2} \left[\frac{1}{2}\dot{\phi}^2 + V(\phi) \right], \quad (19)$$

where the dots denote the derivatives with respect to time t . These equations can be numerically solved with given initial conditions ϕ_{in} and $\dot{\phi}_{in}$. Due to the attractor type of Starobinsky inflation the dependence upon initial conditions is weak [25]. A solution with the initial conditions $\phi_{in} = 7.01M_{Pl}$ and $\dot{\phi}_{in} = 0$ leading to $\phi_{exit}/M_{Pl} = 6.98$ is given in Figure 4, where the Hubble function has two plateaus.




Figure 4. The evolution of inflaton field $\phi(t)$ and Hubble function $H(t)$ with the initial conditions $\phi_{in} = 7.01 M_{Pl}$ and $\dot{\phi}_{in} = 0$, and the parameters $\delta = 2.7 \cdot 10^{-8}$ and $R_0 = 3.0 M^2$.

Inflaton slowly rolls along both plateaus; between them, there is a period of ultra-slow-roll (USR) where dynamics are different [26,27]. Namely, the acceleration term should be kept, but the potential term can be ignored in the first equation of motion (19). Because of that, it is more illuminating to use the Hubble flow parameters outside the CMB region, which are defined by:

$$\epsilon_H = -\frac{\dot{H}}{H^2}, \quad \eta_H = \epsilon_H - \frac{\dot{\epsilon}_H}{2\epsilon_H H}. \quad (20)$$

The evolution is given in Figure 5. In the USR regime, the parameter ϵ_H becomes very small. The parameter η_H drops below -6 , while the corresponding hole (dip) defines the duration of the USR regime.




Figure 5. The evolution of the Hubble flow parameters $\epsilon_H(t)$ and $\eta_H(t)$ with the initial conditions $\phi_{in} = 7.01 M_{Pl}$ and $\dot{\phi}_{in} = 0$, and the parameters $\delta = 2.7 \cdot 10^{-8}$ and $R_0 = 3.0 M^2$.

4. Power Spectrum of Scalar Perturbations and PBH Masses

The power spectrum $P_\zeta(k)$ of scalar (curvature) perturbations, as a function of scale k , is usually derived as a solution to the Mukhanov–Sasaki (MS) Equation [28,29], which is often possible only numerically. However, we found that the following, well-known simple analytic formula [12,18,30]:

$$P_\zeta(t) = \frac{H^2}{8M_{Pl}^2 \pi^2 \epsilon_H} \tag{21}$$

gives a good approximation [19,24]. Our new result is given by Figure 6, which shows a large enhancement (peak) in the power spectrum by a factor of 10^7 against the CMB level (on the left-hand side from the peak).




Figure 6. The primordial power spectrum $P_\zeta(t)$ of scalar (curvature) perturbations, derived from Equation (21).

The PBH masses can be estimated from the peak in the power spectrum as follows [31]:

$$M_{PBH} \simeq \frac{M_{Pl}^2}{H(t_{peak})} \exp \left[2(N_{total} - N_{peak}) + \int_{t_{peak}}^{t_{total}} \epsilon_H(t) H(t) dt \right], \tag{22}$$

where N_{peak} and t_{peak} stand for the peak event, while N_{total} and t_{total} denote the end of the whole inflation comprising three stages (SR/USR/SR).

According to Equation (22), the PBH masses are exponentially sensitive to the number of e-folds around the inflection point, $\Delta N = (N_{total} - N_{peak})$, while the integral in the exponential describes the subleading correction that is of the order one.

Our findings are summarized in Table 1 where the values of the key observables n_s , r , and M_{PBH} in our model are collected with the fine-tuned parameters $b = 2.89$ and $g = 2.25$. The height of the peak is sensitive to R_0 , whose value $R_0/M^2 = 3.001$ was chosen to obtain a height equal to 10^{-2} .

Table 1. CMB observables, PBH masses, and duration of inflation in our model.

ϕ_{in}/M_{Pl}	δ	n_s	r	M_{PBH}, g	N_{peak}	N_{total}	ΔN
6.36	2.55×10^{-7}	0.964959	0.0359	5.0×10^{19}	26	47	21
6.70	8.74×10^{-8}	0.964905	0.0182	2.0×10^{19}	34	54	20
7.01	2.70×10^{-8}	0.964944	0.0095	1.0×10^{20}	43	65	22
7.07	2.05×10^{-8}	0.964917	0.0083	2.6×10^{18}	45	64	19
7.12	1.60×10^{-8}	0.964925	0.0074	1.0×10^{17}	47	65	18
7.15	1.36×10^{-8}	0.964908	0.0070	5.0×10^{16}	49	66	17
7.20	1.02×10^{-8}	0.964961	0.0062	1.6×10^{15}	51	64	13

The tensor-to-scalar ratio r is inside the current observational bound, $r < 0.032$, except the first line in the table (given for comparison only). The tilt n_s of scalar perturbations agrees within 1σ with the current CMB measurements [3–5]:

$$n_s = 0.9649 \pm 0.0042 . \tag{23}$$

To obtain PBH masses beyond the Hawking (black hole) evaporation limit of $10^{15} g$, so that these PBH can survive in the present universe and form dark matter, the duration ΔN should be between 17 and 22 e-folds. It also follows from the table that the total duration of inflation should be between 54 and 66 e-folds.

Increasing the parameter δ allows us to compensate the decreasing scalar tilt n_s . When trying to increase the PBH masses by reducing the total duration of inflation, we find that the value of the tensor-to-scalar ratio r increases and reaches the maximal observationally allowed bound. Increasing ΔN even higher is also not possible, because it leads to the peak height beyond observational constraints. Therefore, it is not possible to increase the PBH masses beyond the asteroid-size with $10^{20} g$ or, equivalently, beyond $10^{-13} M_{Sun}$ in our model.

5. Conclusions

The main new results of this paper are summarised in the abstract. The modified gravity framework is entirely based on gravitational interactions, which implies the gravitational origin of both inflation and PBH formation in our approach. The good agreement (within 1σ) with CMB observations is achieved by fine tuning the parameters of the improved model. The PBH masses found are in the mass window that allows the whole dark matter composed of PBH of the asteroid size [11,12]. The stochastic gravitational waves induced by PBH production in the Starobinsky-like gravity were investigated in Ref. [32].

Fine tuning of the parameters in our model is necessary to obtain a significant enhancement of the power spectrum of scalar perturbations leading to the PBH with masses beyond the Hawking evaporation limit and, hence, the possible PBH dark matter, cf. Ref. [33].

A large peak in the primordial power spectrum may lead to large quantum corrections, which may rule out the near-inflection mechanism of PBH production in all single-field models of inflation [34–36]. However, validity of the mechanism based on a near-inflection point was recently defended in Refs. [37–41]. Modified gravity offers a different perspective on the issue of quantum corrections when assuming the gravitational origin of inflation (as in the Starobinsky model) and PBH production in the context of $F(R)$ gravity theories, as in our paper. Both inflation and PBH formation can be destroyed by adding to the F -function terms with the higher powers of R describing quantum gravity corrections, with the R^4 term being their representative. To avoid that, the coefficients in those terms should be small enough in order to keep validity of the gravitational effective action described by Equation (9). Naturally, there should also be a fundamental mechanism in quantum gravity that keeps these coefficients small; however, this issue is beyond the scope of this paper.

Author Contributions: All authors have contributed equally. All authors have read and agreed to the published version of the manuscript.

Funding: S.S. and S.V.K. were supported by Tomsk State University. S.V.K. was also supported by Tokyo Metropolitan University, the Japanese Society for Promotion of Science under the grant No. 22K03624, and the World Premier International Research Center Initiative (MEXT, Japan).

Data Availability Statement: No new data was created.

Acknowledgments: The authors are grateful to Antonio Iovino Junior, Theodoros Papanikolaou and Jacopo Fumagalli for correspondence.

Conflicts of Interest: The authors declare no conflict of interest.

Note

¹ Accordingly, we changed the notation for the AB-parameter denoted by ϵ_{AB} in Ref. [20] to E_{AB} .

References

- Guth, A.H. The Inflationary Universe: A Possible Solution to the Horizon and Flatness Problems. *Phys. Rev. D* **1981**, *23*, 347–356. [\[CrossRef\]](#)
- Linde, A.D. A New Inflationary Universe Scenario: A Possible Solution of the Horizon, Flatness, Homogeneity, Isotropy and Primordial Monopole Problems. *Phys. Lett. B* **1982**, *108*, 389–393. [\[CrossRef\]](#)
- Akrami, Y. et al. [Planck Collaboration]. Planck 2018 results. X. Constraints on inflation. *Astron. Astrophys.* **2020**, *641*, A10. [\[CrossRef\]](#)
- Ade, P.A.R. et al. [BICEP, Keck Collaboration]. Improved Constraints on Primordial Gravitational Waves using Planck, WMAP, and BICEP/Keck Observations through the 2018 Observing Season. *Phys. Rev. Lett.* **2021**, *127*, 151301. [\[CrossRef\]](#)
- Tristram, M.; Banday, A.J.; Górski, K.M.; Keskitalo, R.; Lawrence, C.R.; Andersen, K.J.; Wehus, I.K. Improved limits on the tensor-to-scalar ratio using BICEP and Planck data. *Phys. Rev. D* **2022**, *105*, 83524. [\[CrossRef\]](#)
- Starobinsky, A.A. A new type of isotropic cosmological models without singularity. *Phys. Lett. B* **1980**, *91*, 99–102. [\[CrossRef\]](#)
- Novikov, I.; Zeldovic, Y. Cosmology. *Ann. Rev. Astron. Astrophys.* **1967**, *5*, 627–649. [\[CrossRef\]](#)
- Hawking, S. Gravitationally collapsed objects of very low mass. *Mon. Not. R. Astron. Soc.* **1971**, *152*, 75. [\[CrossRef\]](#)
- Barrow, J.D.; Copeland, E.J.; Liddle, A.R. The Cosmology of black hole relics. *Phys. Rev. D* **1992**, *46*, 645–657. [\[CrossRef\]](#) [\[PubMed\]](#)
- Ivanov, P.; Naselsky, P.; Novikov, I. Inflation and primordial black holes as dark matter. *Phys. Rev. D* **1994**, *50*, 7173–7178. [\[CrossRef\]](#)
- Carr, B.; Kohri, K.; Sendouda, Y.; Yokoyama, J. Constraints on primordial black holes. *Rept. Prog. Phys.* **2021**, *84*, 116902. [\[CrossRef\]](#)
- Escrivà, A.; Kuhnel, F.; Tada, Y. Primordial Black Holes. *arXiv* **2022**, arXiv:2211.05767
- Ketov, S.V.; Starobinsky, A.A. Inflation and non-minimal scalar-curvature coupling in gravity and supergravity. *J. Cosmol. Astropart. Phys.* **2012**, *8*, 22. [\[CrossRef\]](#)
- Ketov, S.V. Multi-Field versus Single-Field in the Supergravity Models of Inflation and Primordial Black Holes. *Universe* **2021**, *7*, 115. [\[CrossRef\]](#)
- Ketov, S. On the large-field equivalence between Starobinsky and Higgs inflation in gravity and supergravity. In Proceedings of the 7th Symposium on Prospects in the Physics of Discrete Sources, Bergen, Norway, 29 November–3 December 2021; p. 14. [\[CrossRef\]](#)
- Maeda, K.-I. Towards the Einstein–Hilbert Action via Conformal Transformation. *Phys. Rev. D* **1989**, *39*, 3159. [\[CrossRef\]](#)
- Ivanov, V.R.; Ketov, S.V.; Pozdeeva, E.O.; Vernov, S.Y. Analytic extensions of Starobinsky model of inflation. *J. Cosmol. Astropart. Phys.* **2022**, *3*, 58. [\[CrossRef\]](#)
- Garcia-Bellido, J.; Morales, E.R. Primordial black holes from single field models of inflation. *Phys. Dark Univ.* **2017**, *18*, 47–54. [\[CrossRef\]](#)
- Frolovsky, D.; Ketov, S.V.; Saburov, S. Formation of primordial black holes after Starobinsky inflation. *Mod. Phys. Lett. A* **2022**, *37*, 2250135. [\[CrossRef\]](#)
- Appleby, S.A.; Battye, R.A.; Starobinsky, A.A. Curing singularities in cosmological evolution of F(R) gravity. *J. Cosmol. Astropart. Phys.* **2010**, *6*, 5. [\[CrossRef\]](#)
- Ketov, S.V.; Pozdeeva, E.O.; Vernov, S.Y. On the superstring-inspired quantum correction to the Starobinsky model of inflation. *J. Cosmol. Astropart. Phys.* **2022**, *12*, 32. [\[CrossRef\]](#)
- Ivanov, V.; Ketov, S.; Pozdeeva, E.; Vernov, S. On Extensions of the Starobinsky Model of Inflation. *Phys. Sci. Forum* **2023**, *7*, 6. [\[CrossRef\]](#)
- Pozdeeva, E.; Ketov, S.; Vernov, S. String-Inspired Correction to R^2 Inflation. *Phys. Sci. Forum* **2023**, *7*, 2.
- Frolovsky, D.; Ketov, S.V. Inflationary E-models revisited. *arXiv* **2023**, arXiv:2304.12558.
- Aoki, S.; Ishikawa, R.; Ketov, S.V. Pole inflation and primordial black holes formation in Starobinsky-like supergravity. *Class. Quant. Grav.* **2023**, *40*, 65002. [\[CrossRef\]](#)

26. Germani, C.; Prokopec, T. On primordial black holes from an inflection point. *Phys. Dark Univ.* **2017**, *18*, 6–10. [[CrossRef](#)]
27. Dimopoulos, K. Ultra slow-roll inflation demystified. *Phys. Lett. B* **2017**, *775*, 262–265. . [[CrossRef](#)]
28. Mukhanov, V.F. Gravitational Instability of the Universe Filled with a Scalar Field. *JETP Lett.* **1985**, *41*, 493–496.
29. Sasaki, M. Large Scale Quantum Fluctuations in the Inflationary Universe. *Prog. Theor. Phys.* **1986**, *76*, 1036. [[CrossRef](#)]
30. Karam, A.; Koivunen, N.; Tomberg, E.; Vaskonen, V.; Veermäe, H. Anatomy of single-field inflationary models for primordial black holes. *J. Cosmol. Astropart. Phys.* **2023**, *3*, 13. [[CrossRef](#)]
31. Pi, S.; Zhang, Y.-L.; Huang, Q.-G.; Sasaki, M. Scalon from R^2 -gravity as a heavy field. *J. Cosmol. Astropart. Phys.* **2018**, *5*, 42. [[CrossRef](#)]
32. Papanikolaou, T.; Tzerefos, C.; Basilakos, S.; Saridakis, E.N. Scalar induced gravitational waves from primordial black hole Poisson fluctuations in $f(R)$ gravity. *J. Cosmol. Astropart. Phys.* **2022**, *10*, 13. [[CrossRef](#)]
33. Cole, P.S.; Gow, A.D.; Byrnes, C.T.; Patil, S.P. Primordial black holes from single-field inflation: A fine-tuning audit. *arXiv* **2023**, arXiv:2304.01997.
34. Kristiano, J.; Yokoyama, J. Ruling Out Primordial Black Hole Formation From Single-Field Inflation. *arXiv* **2022**, arXiv:2211.03395
35. Kristiano, J.; Yokoyama, J. Response to criticism on “Ruling Out Primordial Black Hole Formation From Single-Field Inflation”: A note on bispectrum and one-loop correction in single-field inflation with primordial black hole formation. *arXiv* **2023**, arXiv:2303.00341.
36. Choudhury, S.; Panda, S.; Sami, M. Quantum loop effects on the power spectrum and constraints on primordial black holes. *arXiv* **2023**, arXiv:2303.06066.
37. Riotto, A. The Primordial Black Hole Formation from Single-Field Inflation is Not Ruled Out. *arXiv* **2023**, arXiv:2301.00599.
38. Riotto, A. The Primordial Black Hole Formation from Single-Field Inflation is Still Not Ruled Out. *arXiv* **2023**, arXiv:2303.01727.
39. Firouzjahi, H.; Riotto, A. Primordial Black Holes and Loops in Single-Field Inflation. *arXiv* **2023**, arXiv:2304.07801.
40. Franciolini, G.; Iovino, A., Jr.; Taoso, M.; Urbano, A. One loop to rule them all: Perturbativity in the presence of ultra slow-roll dynamics. *arXiv* **2023**, arXiv:2305.03491.
41. Fumagalli, J. Absence of one-loop effects on large scales from small scales in non-slow-roll dynamics. *arXiv* **2023**, arXiv:2305.19263.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.