

Article

Quantum Fisher Information of Three-Level Atom under the Influence of the Stark Effect and Intrinsic Decoherence

S. Jamal Anwar , M. Ramzan and M. Khalid Khan *

Department of Physics, Quaid-i-Azam University, Islamabad 15320, Pakistan

* Correspondence: kk@hotmail.com

Abstract: We study the dynamical evolution of quantum Fisher information (QFI) and von Neumann entropy (VNE) for a three-level atomic system interacting with the single-mode coherent field in the presence of the Stark effect and intrinsic decoherence (ID) with and without atomic motion. The effect of the ID is significant on the VNE and QFI for a three-level atom in the absence of atomic motion. It is observed that in the case of a three-level atomic system in the presence of ID, the decay of QFI and VNE is rapid and significant but no prominent effect of the Stark effect is observed. Hence, for a three-level atom, the decay of quantum entanglement (QE) with respect to time is very fast and rapid in the absence of atomic motion with an increasing value of ID. Moreover, ID is not suitable to maintain the QE for three-level atomic systems in the absence of atomic motion. The Stark effect has no significant effect on the QE. In the case of three-level atoms, ID and the Stark do not affect the periodic nature of QFI and VNE with time evolution in the presence of atomic motion. The periodic response of QFI and VNE is observed under the effect of the Stark effect and ID in the presence of a motion of a three-level atom. The QE sudden death and birth is observed in the presence of atomic motion. Therefore, the ID with the Stark effect is suitable to sustain and maintain the QE in the presence of atomic motion for three-level atomic systems. These results show the strong dependence of QFI and VNE on the Stark effect and ID.

Keywords: quantum entanglement (QE); quantum fisher information (QFI); the Stark effect; von Neumann entropy (VNE)



Citation: Anwar, S.J.; Ramzan, M.; Khan, M.K. Quantum Fisher Information of Three-Level Atom under the Influence of the Stark Effect and Intrinsic Decoherence.

Atoms **2023**, *11*, 42. <https://doi.org/10.3390/atoms11030042>

Academic Editor: Yew Kam Ho

Received: 15 November 2022

Revised: 21 February 2023

Accepted: 22 February 2023

Published: 28 February 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Quantum entanglement (QE) plays a crucial character in quantum information [1,2]. Therefore, more efforts were made to investigate the formation of the QE in various physical atom-field models and to initiate top QE quantifiers as entropy [3,4] and negativity [5,6]. Entropy is very useful in explaining field and atom correlations and the quantum effects in their connection. In particular, entropy is a much better measure of the QE [7–10]. The binomial states were used to show states intermediate between the coherent and the number states. They are considered in the free-electron laser [11] and have essential quantum characteristics such as anti-bunching [12] and squeezing [13,14]. Recently, a great deal of work is performed on QE in the case of non-inertial frames. Entanglement measures of W-state in non-inertial frames are studied by Ariadna J and co-workers [15]. Qiang, WC., Dong, Q., Mercado Sanchez, M.A. et al. studied QE property of the Werner state in accelerated frames [16]. Qian Dong et al. studied tripartite QE measures of generalized GHZ State in uniform acceleration [17].

Fisher information (FI) was first introduced by Fisher as the main quantity in estimation theory [18]. The members of probability distributions can be detected by FI. In this regard, FI can perform optimal tasks in detecting the optimal measurement of the considered quantum systems from the point of view of the theory of quantum estimation. The parameter estimation problem being the more significant is the main support of various science and technology branches. The phase estimation plays the essential role of state

generation, decoherence and loss [19–24]. The lower bound of Quantum Fisher information (QFI) can be obtained from the quantum version of the Cramér–Rao (CR) inequality [25]. Interestingly, the mathematical treatments of the lower bound in physical models have been clarified [26,27], in this way the optimal resource for the phase estimation has been discussed [28]. Fisher information is a metric for figuring out how much information an observable variable x has about an unknown parameter of a distribution that represents X in mathematics. For example, it is the variance of the score or the expected value of the observed data [14,29]. As a result, a greater FI suggests that we can more precisely anticipate the parameter [30]. Quantum Fisher information (QFI) is FI’s quantum extension, which is stated in terms of quantum Cramér–Rao bound [31–33]. According to the quantum Cramér–Rao inequality [34], the value of QFI provides an upper bound on the parameter precision. Therefore, the larger (QFI) indicates higher precision. QFI has been widely researched from a variety of perspectives. It has been discussed how to increase estimation precision [35,36], measurement effectiveness [37,38], and QFI function self-utilization [39,40]. It is known that the atomic parameter estimation, nonclassical properties of the field, and its atomic version of the QFI are essential tools in some applications of quantum optics and information processing. QFI is a useful quantifier to estimate different information resources, such as the entanglement, quantum speedup [41], and quantum phase transition [42]. It also has practical applications in quantum metrology [43], involves high precision.

Quantum von Neumann is considered the main entanglement measure for pure states. The correlation between the quantum entanglement measured by the von Neumann entropy and Wehrl entropy has been taken into account [44,45]. On the other hand, the correlation between QE and Fisher information (FI), as we know about a certain parameter in a quantum state, has not been studied widely. However, there are some studies to quantify pure state entanglement by using FI. In this regard, the QE evaluation with atomic classical Fisher information has been investigated [27,46,47]. It has been shown that QE of a two-level atom can be quantified by atomic FI and their marginal distribution. Furthermore, the correlation between the FI and QE during the time evolution for a trapped ion in the laser field. It is found that FI is an important tool to study single qubit dynamics as an indicator of QE under certain conditions [48]. Additionally, the time evolution of the QFI of a system whose dynamics are described by the phase-damped model has been studied [49]. It observed that there is an interesting monotonic relation between the QFI and non-local correlation behavior measured by the negativity depending on choosing the estimator parameter during the time evolution.

The interactions between atoms and the electromagnetic field are the roots of quantum optics and fundamental resources of quantum information and there are plenty of various models that describe these types of interactions. One of the most important types is a system consisting of a three-level atom that interacts with single or multi-mode fields [50–52]. As an example, Obada et al. [53] discussed the behaviour of non-classical correlation for a three-level atom interacting with non-linear two-cavity modes. The Kerr Medium’s effect on the QE that is generated between a Cascade-type of a three-level atom and a bimodal cavity field is discussed by Teng et al. [54]. Mortezaapour et al. [55] investigated the QE phenomena generated between an atomic system consisting of three levels prepared initially on the configuration with two non-linear cavity modes in the presence of a classical driving field. The phenomena of the sudden death/birth of different dimensions of atomic systems interacting with a classical driven classical field are investigated by Metwally et al. [56]. However, these models have been expanded to hold: the atomic motion [57], damping field via Caldirola and Kanai (CK) Hamiltonian [58], classical homogeneous gravitational field [59] and correlated two-mode field [60]. The dynamics of the QE in these systems are investigated for different considerations as two-cavity modes with parametric down-conversion [57] and time-dependent field [61].

In open quantum systems where the dissipative effects, a type of decoherence destroys the QE and mixture of the system and also it weakens the useful quantum properties of the system and its sub-systems, such as the quantum correlations. There are several methods

to study quantum decoherence that are responsible for the transition between quantum and classical systems. Intrinsic decoherence is one of them that is based on the modified Schrodinger equation which reduces to the Milburn equation that describes the intrinsic decoherence [62,63]. In this modification, the quantum coherence is automatically destroyed as the system evolves. Intrinsic decoherence (ID) is a mechanism by which the system loses its coherence because of its intrinsic degrees of freedom. Several models of ID have been proposed [64–66], and one of them is the Milburn decoherence model. Milburn proposed a simple modification to the Schrodinger equation based on the assumption that the change in the state of the system, i.e., $\rho(t) \rightarrow \rho(t + \tau)$ is uncertain and occurs with a probability of $0 < p(\tau) < 1$ on a sufficiently small time scale [62]. The change of state is always certain in standard quantum mechanics and $p(\tau) = 1$. In the last decade, the ID of quantum coupled discrete systems has been widely studied [67–70], but the majority of those works dealt with diffusion approximation, i.e., first order-correction of the von Neumann equation. A.B.A. Mohamed observed non-local correlation and quantum discord in two atoms in the non-degenerate model [71]. Mohamed, A.B.A., Eleuch, H. described optical tomography dynamics induced by qubit-resonator interaction under intrinsic decoherence [72]. A.B.A. Mohamed et al. studied quantum coherence induced by a flux qubit coupled by a resonator coherent field through a two-photon interaction [73]. A.B.A. Mohamed and H.A. Hessian discussed non-classicality in an open two-mode parametric amplifier cavity containing a qutrit system [74].

In atomic physics, the Stark effect is the split and shift of a spectral line into several components in the presence of an electric field. The amount of splitting is called the Stark shift [75]. It is analogous to the Zeeman effect where a spectral line is split into several components in the presence of a magnetic field. Its effect has been studied on some phenomena in different quantum systems. The interaction between a two-level atom and coherent field has been studied [76], where the effects of Kerr-like medium and Stark shift parameters on Wehrl entropy and field purity are discussed. The effect of Stark shift terms on the interaction between a radiation field and two two-level atoms is studied [77]. The dynamic of quantum Fisher information and entanglement for three- and four-level atomic systems interacting with a coherent field under the effect of Stark shift and Kerr medium is considered in [78]. Additionally, the entanglement for two three-level atomic systems influenced by Stark shift and Kerr-like medium has been studied [79]. The dynamics of quantum entanglement for an N-level atomic system interacting with a coherent field in the presence of Stark effects and Kerr-like medium has been studied [80].

The QFI and VNE quantum systems like three-level atoms with ID in the presence of the Stark effect have not been investigated so far on a large scale. This motivated us to investigate a three-level atomic system with ID in the presence of the Stark effect. It is very important to study the behaviour of QE for three-level atomic systems in the presence of environmental effects such as the Stark effect and ID. Mahmoud Abdel-Aty, S. Furuichi and A-S. F. Obada studied the influences of non-linearity, Stark shifts and detuning on the degree of entanglement [81]. Baghshahi H. R. et al. studied entropy squeezing and atomic inversion in the k-photon Jaynes Cummings model in the presence of the Stark shift and a Kerr medium [82]. The Dynamics and maintenance of bipartite entanglement via the Stark shift effect inside the dissipative reservoirs is studied by S. Golkar1 and M. K. Tavassoly [83], A-S. F. Obada1, S. Abdel-Khalek, E. M. Khalil, and S. I. Ali studied effects of Stark shift and decoherence terms on the dynamics of phase-space entropy of the multiphoton Jaynes Cummings model [84].

The present paper aims to find the QE dynamics of the three-level atomic system with atomic motion in the presence of the Stark effect and ID. We investigate the dynamics of VNE and QFI for moving three-level atomic systems under the influence of the ID and Stark effect. It is seen that the ID and Stark play a dominant role during the time evolution of the quantum system. The dynamics of QFI are heavily influenced by the Stark effect in the presence of ID. Moreover, the Stark effect is more prominent on the VNE in the presence

of the ID without atomic motion. Finally, the three-level atomic system is found highly sensitive to these environmental effects.

The paper is organized as follows. In Section 2, we present the model Hamiltonian and interaction dynamics of the coherent field of moving-level atomic systems influenced by the ID and the Stark effect. In Section 3, numerical results and discussions are presented. In Section 4, we present a brief conclusion.

2. Hamiltonian Model

We consider moving three-level atoms entering the cavity at time $t = 0$ in the superposition state in the presence of the Stark effect influenced by the ID. We consider the cascade configuration of moving a three-level atomic system in the presence of the Stark effect with ID.

The total Hamiltonian of the system \hat{H}_T under the RWA for a particular system can be described as [85]

$$\hat{H}_T = \hat{H}_{Atom-Field} + \hat{H}_I. \tag{1}$$

where $\hat{H}_{Atom-Field}$ is representing the Hamiltonian for the non-interacting atom and field, and the interaction part is given by \hat{H}_I . We will write $\hat{H}_{Atom-Field}$ as

$$\hat{H}_{Atom-Field} = \sum_j \omega_j \hat{\sigma}_{j,j} + \nu \hat{a}^\dagger \hat{a}, \tag{2}$$

where $\hat{\sigma}_{j,j} = |j\rangle\langle j|$ are described as population operators for the j th level ν is the frequency of oscillation of field mode. The interaction Hamiltonian of a three-level atomic system for the non-resonant case can be written as [85]

$$\hat{H}_I = \Omega(t) [\hat{a}(|2\rangle\langle 3|e^{i\Delta t} + \hat{a}|1\rangle\langle 2|e^{-i\Delta t}) + h.c]. \tag{3}$$

where \hat{a} and \hat{a}^\dagger is the annihilation (creation) operator of the field mode. Two is the middle energy level, one is the upper energy level and two is the lower energy level. Δ is the detuning of the field mode.

In the case of the Stark effect, the interaction Hamiltonian can be written as

$$\hat{H}_I = \Omega(t) [\hat{a}(|2\rangle\langle 3|e^{i\Delta t} + \hat{a}|1\rangle\langle 2|e^{-i\Delta t}) + h.c] + \beta \hat{a}^{\dagger 2} \hat{a}^2, \tag{4}$$

β is the parameter of the Stark effect and the detuning parameter is described as

$$\Delta = \omega - (\omega_2 - \omega_1) = \omega - (\omega_2 - \omega_1). \tag{5}$$

The coupling constant for the atom and field is g , $\Omega(t)$ represents the shape function of the cavity-field mode [86] and the motion of the atom is along the z -axis. For particular interest

$$\begin{aligned} \Omega(t) &= g \sin(p\pi vt/L) \text{ in the presence of atomic motion, } p \neq 0 \\ \Omega(t) &= g \text{ in the absence of atomic motion } p = 0 \end{aligned} \tag{6}$$

where the velocity of motion of the atom is v and p denotes half of the number of wavelengths of the mode in the cavity and L describes the length of the cavity along the z -direction. Now take the velocity of an atom as $v = gL / \pi$ which gives us

$$\Omega_1(t) = \int_0^t \Omega(\tau) d\tau = \frac{1}{p} (1 - \cos(p\pi vt/L)) \text{ for } p \neq 0 \tag{7}$$

$$= gt \text{ for } p = 0. \tag{8}$$

To find the phase shift parameter as precisely as possible, we consider the optimal input state as

$$|\Psi(0)\rangle_{\text{Opt}} = \frac{1}{\sqrt{2}}(|1\rangle + |2\rangle) \otimes |\alpha\rangle \tag{9}$$

where $|1\rangle$ and $|2\rangle$ describe the states of the atom and α is the coherent state of the input field given as

$$|\alpha\rangle = \sum_{n=0}^{\infty} \alpha^n \sqrt{e^{-|\alpha|^2}/n!} |n\rangle. \tag{10}$$

We consider a single-atom phase gate that introduces the phase shift as

$$\hat{U}_\phi = |1\rangle\langle 1| + e^{i\phi}|2\rangle\langle 2|, \tag{11}$$

$|\Psi(0)\rangle$ is obtained from the operation of the single-atom phase gate on $|\Psi(0)\rangle_{\text{Opt}}$

$$\hat{U}_\phi |\Psi(0)\rangle_{\text{Opt}} = |\Psi(0)\rangle \tag{12}$$

$$= \frac{1}{\sqrt{2}}(|1\rangle + e^{i\phi}|2\rangle) \otimes |\alpha\rangle \tag{13}$$

After the operation of the phase gate, the system will interact with a field. For the time-dependent case, the wave function can be characterized by the transformation matrix $\hat{U}(t)$ as given by

$$|\Psi(t)\rangle = \hat{U}(t)|\Psi(0)\rangle, \tag{14}$$

One can write the explicit expression of the density matrix as

$$\hat{\rho}(t) = \sum_{m,n}^N |\Psi_n(t)\rangle\langle \Psi_n(t)| \hat{\rho}(0) |\Psi_m(t)\rangle\langle \Psi_m(t)| \tag{15}$$

and in terms of ID, the density matrix can be as [87]

$$\hat{\rho}(t) = \sum \exp\left(\frac{\gamma t}{2}(E_m - E_n)^2 - i(E_m - E_n)t\right) \times \langle \Psi_m | \hat{\rho}(0) | \Psi_n \rangle | \Psi_m \rangle \langle \Psi_n |, \tag{16}$$

where $E_{m,n}$ and $\Psi_{m,n}$ are the eigenvalues and the corresponding eigenvectors of H_I , respectively, and γ is the parameter of ID.

In this fashion, we can define the QFI of a bipartite density operator ρ_{AB} , in terms of ϕ as [88]

$$I_{QF}(t) = I(\phi, t) = \text{Tr}[\rho_{AB}(\phi, t) \{L^2(\phi, t)\}], \tag{17}$$

where $L(\phi, t)$ is the quantum score [89] (the symmetric logarithmic derivative) which can be found as

$$\frac{\partial \rho(\phi, t)}{\partial \phi} = \frac{1}{2} [L(\phi, t) \rho_{AB}(\phi, t) + \rho_{AB}(\phi, t) L(\phi, t)]. \tag{18}$$

Similarly, the VNE can be defined as

$$S_A = -\text{Tr}(\rho_A \ln \rho_A) = -\sum_i r_i \ln r_i, \tag{19}$$

where r_i are the eigenvalues of the atomic density matrix $\rho_A = \text{Tr}_B(\rho_{AB})$.

Now, the influence of different environmental parameters γ , ϕ and p on the evolution of the QFI and VNE is presented in the next section.

3. Numerical Results and Discussion

In this section, we present the results for the dynamical evolution of QFI and VNE for moving three-level atomic system interacting with a coherent field with the Stark effect

in the presence of the ID. We study the time evolution of QFI and VNE of the three-level atomic system in the presence of the Stark effect and ID with and without atomic motion.

In Figures 1–3, we plot QFI and VNE vs time for two- and three-level atomic system interacting with a coherent field for $|\alpha|^2 = 6$ and the phase shift parameter $\phi = 0$ (left panel) and $\pi/4$ (right panel) for $\beta = 0.3, 1$ and 3 and ID parameter $\gamma = 0.00001, 0.0001$ and 0.001 without atomic motion, i.e., $p = 0$. The effect of the Stark shift is not seen as significant on the VNE and QFI of the three-level atom in the absence of atomic motion. It is observed that ID influenced the QFI and VNE and on the increasing value of ID parameter γ , the decay is observed in both QFI and VNE with the evolution of time. At the value $\gamma = 0.001$, the QFI and VNE decrease more rapidly with the evolution of time. It means QE is decreasing rapidly with the increasing value of the ID parameter. However, at $\beta = 0.3, 1$ and 3 , the Stark shift does not influence the QE significantly, but the ID influences the QE strongly.

The periodic behaviour of QFI and VNE is suppressed under the effect of the Stark effect with ID in the absence of atomic motion. It is observed the ID affects the QFI and VNE at larger values but the increasing value of the Stark effect does not affect QFI and VNE significantly. It means the Stark effect does not affect the QE of the three-level atomic system.




Figure 1. (Colour online) The QFI (upper panel) and VNE (lower panel) as a function of time for three-level atom interacting with a coherent field for $\alpha = 6$ in the presence of the Stark effect ($\beta = 0.3$), $\gamma = 0.00001, 0.0001$ and 0.001 and the phase shift estimator parameter $\phi = 0$ (left panel) and $\pi/4$ (right panel). The atomic motion parameter p is neglected.




Figure 2. (Colour online) The QFI (upper panel) and VNE (lower panel) as a function of time for three-level atom interacting with a coherent field for $\alpha = 6$ in the presence of the Stark effect ($\beta = 1$), $\gamma = 0.00001, 0.0001$ and 0.001 and the phase shift estimator parameter $\phi = 0$ (left panel) and $\pi/4$ (right panel). The atomic motion parameter p is neglected.




Figure 3. (Colour online) The QFI (upper panel) and VNE (lower panel) as a function of time for four-level atom interacting with a coherent field for $\alpha = 6$ in the presence of the Stark effect ($\beta = 3$), $\gamma = 0.00001, 0.0001$ and 0.001 and the phase shift estimator parameter $\phi = 0$ (left panel) and $\pi/4$ (right panel). The atomic motion parameter p is neglected.

In Figures 4–6, we plot QFI and VNE for three-level atomic systems interacting with the coherent field in the presence of the ID for $\beta = 0.3, 1$ and 3 and $p = 1$. It is observed that in the case of atomic motion ($p = 1$), both QFI and VNE show periodic behaviour with respect to time for three-level atoms. The Stark effect and ID have no significant effect on the periodicity of the VNE and QFI for three-level atom. For three-level atom the periodic behaviour is the same for ID parameter $\gamma = 0.00001, 0.0001$ and 0.001 . So it means that in the case of three level-atom, ID and the Stark effect do not affect the periodic nature of QFI and VNE with time evolution. Hence it is observed that sudden death and birth of QE is not affected by ID and the Stark effect for three-level atom in the presence of the atomic motion ($p = 1$). Therefore, it is concluded that the ID with the Stark effect is suitable to sustain and maintain the QE in the presence of atomic motion for the N-level atomic system.




Figure 4. (Colour online) The QFI (upper panel) and VNE (lower panel) as a function of time for two-level atom interacting with a coherent field for $\alpha = 6$ in the presence of the Stark effect ($\beta = 0.3$), $\gamma = 0.00001, 0.0001$ and 0.001 and the phase shift estimator parameter $\phi = 0$ (left panel) and $\pi/4$ (right panel). The atomic motion parameter $p = 1$.




Figure 5. Cont.




Figure 5. (Colour online) The QFI (upper panel) and VNE (lower panel) as a function of time for three-level atom interacting with a coherent field for $\alpha = 6$ in the presence of the Stark effect ($\beta = 1$), $\gamma = 0.00001, 0.0001$ and 0.001 and the phase shift estimator parameter $\phi = 0$ (left panel) and $\pi/4$ (right panel). The atomic motion parameter $p = 1$.




Figure 6. (Colour online) The QFI (upper panel) and VNE (lower panel) as a function of time for three-level atom interacting with a coherent field for $\alpha = 6$ in the presence of the Stark effect ($\beta = 3$), $\gamma = 0.00001, 0.0001$ and 0.001 and the phase shift estimator parameter $\phi = 0$ (left panel) and $\pi/4$ (right panel). The atomic motion parameter $p = 1$.

4. Conclusions

We have investigated the dynamics of VNE and QFI for the three-level atomic system affected by the Stark effect and ID. The time evolution of QFI and VNE for three-level atomic systems influenced by ID and the Stark was examined with and without atomic motion. It is observed that the ID has a prominent role during the time evolution of the quantum system but the Stark effect does not influence the three-level atomic system significantly. The effect of ID is seen prominent on the VNE as compared to the QFI in the absence of the atomic motion but the Stark effect shows no prominent effect. The influence of the Stark effect is not found significant on the VNE and QFI of the three-level atom in the absence of atomic motion. It is observed that in the case of larger values of the ID parameter γ especially at $\gamma = 0.001$, the VNE decreases more rapidly as compared to the QFI. It means QE is influenced by a larger value ID parameter and it reduces rapidly. So

it is concluded that for the three-level atomic system, the QE does not maintain itself at increasing values of ID but increasing values of the Stark parameter β does not affect the QE. Hence, the Stark effect does not influence the QE of the three-level atom, so the Stark effect is useful to maintain QE but ID is not suitable to sustain QE. For a three-level atom, ID and the Stark effect do not affect the periodic nature of QFI and VNE with time evolution in the presence of atomic motion. It is observed that sudden death and birth of QE is not affected by ID and the Stark effect for the three-level atom in the presence of the atomic motion ($p = 1$). The periodic response of QFI and VNE is not suppressed under the effect of the Stark effect and ID in the presence of motion of atoms with larger values of ID and the Stark effect parameter. The QE sudden death and birth is not suppressed at smaller or larger values of the ID and the Stark parameter as compared to the larger values. Therefore, it is concluded that the ID with the Stark effect is suitable to sustain and maintain the QE in the presence of atomic motion for a three-level atomic system. These results show the strong dependence of QFI and VNE on the Stark effect and ID.

Author Contributions: S.J.A., makes the calculations and write the paper. M.R. reviews the paper and make corrections. M.K.K. corrects the english. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Nielsen, M.A.; Chuang, I.L. *Quantum Computation and Quantum Information*; Cambridge University Press: Cambridge, UK, 2000.
- Englert, B.-G.; Metwally, N. Kinematics of qubit pairs. In *Mathematics of Quantum Computation*; Brylinski, R., Chen, G., Eds.; CRC Press, Taylor & Francis Group: Boca Raton, FL, USA, 2002; pp. 25–75.
- Phoenix, S.J.D.; Knight, P.L. Fluctuations and entropy in models of quantum optical resonance. *Ann. Phys.* **1988**, *186*, 381–407. [[CrossRef](#)]
- Phoenix, S.J.D.; Knight, P.L. Establishment of an entangled atom-field state in the Jaynes–Cummings model. *Phys. Rev. A* **1991**, *44*, 6023–6029. [[CrossRef](#)]
- Vedral, V.; Plenio, M.B.; Rippin, M.A.; Knight, P.L. Quantifying entanglement. *Phys. Rev. Lett.* **1997**, *78*, 2275–2279. [[CrossRef](#)]
- Metwally, N. Quantum dense coding and dynamics of information over Bloch channels. *J. Phys. A* **2011**, *44*, 055305. [[CrossRef](#)]
- Obada, A.S.; Ahmed, M.M.A.; Faramawy, F.K.; Khalil, E.M. Entropy and entanglement of the nonlinear Jaynes–Cummings model. *Chin. J. Phys.* **2004**, *42*, 79–91
- Hines, A.P.; Dawson, C.M.; McKenzie, R.H.; Milburn, G.J. Entanglement and bifurcations in Jahn–Teller models. *Phys. Rev. A* **2004**, *70*, 022303. [[CrossRef](#)]
- Jia, X.; Subramaniam, A.R.; Gruzberg, I.A.; Chakravarty, S. Entanglement entropy and multifractality at localization transitions. *Phys. Rev. B* **2008**, *77*, 014208. [[CrossRef](#)]
- Abdelghany, R.A.; Mohamed, A.B.; Tammam, M.; Kuo, W.; Eleuch, H. Tripartite entropic uncertainty relation under phase decoherence. *Sci. Rep.* **2021**, *11*, 11830. [[CrossRef](#)] [[PubMed](#)]
- Dattoli, G.; Gallardo, J.; Torre, A. Binomial states of the quantized radiation field: Comment. *J. Opt. Soc. Am. B* **1987**, *4*, 185–187. [[CrossRef](#)]
- Verma, A.; Sharma, N.K.; Pathak, A. Higher order antibunching in intermediate states. *Phys. Lett. A* **2008**, *372*, 5542–5551. [[CrossRef](#)]
- Agarwal, G.S.; Tara, K. Nonclassical properties of states generated by the excitations on a coherent state. *Phys. Rev. A* **1991**, *43*, 492–497. [[CrossRef](#)]
- Tavassoly, M.K.; Hekmatara, H. Entanglement and other nonclassical properties of two two-level atoms interacting with a two-mode binomial field: Constant and intensity-dependent coupling regimes. *Commun. Theor. Phys.* **2015**, *64*, 439–446. [[CrossRef](#)]
- Torres-Arenas, A.J.; Dong, Q.; Sun, G.H.; Qiang, W.C.; Dong, S.H. Entanglement measures of W-state in noninertial frames. *Phys. Lett. B* **2019**, *789*, 93–105. [[CrossRef](#)]
- Qiang, W.C.; Dong, Q.; Mercado Sanchez, M.A.; Sun, G.H.; Dong, S.H. Entanglement property of the Werner state in accelerated frames. *Quantum Inf. Process.* **2019**, *18*, 314. [[CrossRef](#)]
- Dong, Q.; Sanchez, M.M.; Sun, G.H.; Toutounji, M.; Dong, S.H. Tripartite Entanglement Measures of Generalized GHZ State in Uniform Acceleration. *Chin. Phys. Lett.* **2019**, *36*, 100301. [[CrossRef](#)]
- Fisher, R.A. Theory of Statistical Estimation. *Proc. Camb. Philos. Soc.* **1929**, *22*, 700; reprinted in *Collected Papers of R.A. Fisher*; Bennett, J.H., Ed.; Univ. of Adelaide Press: Adelaide, South Australia, 1972; pp. 15–40. [[CrossRef](#)]

19. Giovannetti, V.; Lloyd, S.; Maccone, L. Quantum-Enhanced Measurements: Beating the Standard Quantum Limit. *Science* **2004**, *306*, 1330–1336. [[CrossRef](#)] [[PubMed](#)]
20. Dowling, J. Quantum Optical Metrology—The Lowdown On High-N00N States. *Contemp. Phys.* **2008**, *49*, 125–143. [[CrossRef](#)]
21. Jones, J.A.; Karlen, S.D.; Fitzsimons, J.; Ardavan, A.; Benjamin, S.C.; Briggs, G.A.D.; Morton, J.J.L. Magnetic Field Sensing Beyond the Standard Quantum Limit Using 10-SpinNOON States. *Science* **2009**, *324*, 1166–1168. [[CrossRef](#)] [[PubMed](#)]
22. Simmons, S.; Jones, J.A.; Karlen, S.D.; Ardavan, A.; Morton, J.J.L. Magnetic field sensors using 13-spin cat states. *Phys. Rev. A* **2010**, *82*, 022330. [[CrossRef](#)]
23. Higgins, B.L.; Berry, D.W.; Bartlett, S.D.; Wiseman, H.M.; Pryde, G.J. Entanglement-free Heisenberg-limited phase estimation. *Nature* **2007**, *450*, 393–396. [[CrossRef](#)] [[PubMed](#)]
24. Dorner, U.; Smith, B.J.; Lundeen, J.S.; Wasilewski, W.; Banaszek, K.; Walmsley, I.A. Quantum phase estimation with lossy interferometers. *Phys. Rev. A* **2009**, *80*, 013825.
25. Cramer, H. *Mathematical Methods of Statistics*; Princeton University Press: Princeton, NJ, USA, 1946.
26. Dorner, U.; Demkowicz-Dobrzanski, R.; Smith, B.J.; Lundeen, J.S.; Wasilewski, W.; Banaszek, K.; Walmsley, I.A. Optimal Quantum Phase Estimation. *Phys. Rev. Lett.* **2009**, *102*, 040403. [[CrossRef](#)] [[PubMed](#)]
27. Abdel-Khalek, S. Dynamics of Fisher information in Kerr medium. *Int. J. Quantum Inf.* **2009**, *7*, 1541–1548. [[CrossRef](#)]
28. Joo, J.; Munro, W.J.; Spiller, T.P. Quantum Metrology with Entangled Coherent States. *Phys. Rev. Lett.* **2011**, *107*, 083601. [[CrossRef](#)]
29. Rissanen, J.J. Fisher information and stochastic complexity. *IEEE Trans. Inf. Theory* **1996**, *42*, 40–47. [[CrossRef](#)]
30. Braunstein, S.L.; Caves, C.M. Statistical distance and the geometry of quantum states. *Phys. Rev. Lett.* **1994**, *72*, 3439. [[CrossRef](#)]
31. Alipour, S.; Mehboudi, M.; Rezakhani, A.T. Quantum metrology in open systems: Dissipative Cramér-Rao bound. *Phys. Rev. Lett.* **2014**, *112*, 120405. [[CrossRef](#)]
32. Liu, Y.; Yu, M.; Yang, P.; Gong, M.; Cao, Q.; Zhang, S.; Cai, J. Saturating the quantum Cramér-Rao bound and measuring the related quantum Fisher information in a nitrogen-vacancy center in diamond. *arXiv* **2020**, arXiv:2003.08373.
33. Liu, J.; Yuan, H.; Lu, X.M.; Wang, X. Quantum Fisher information matrix and multiparameter estimation. *J. Phys. A Math. Theor.* **2019**, *53*, 023001. [[CrossRef](#)]
34. Helstrom, C.W. *Quantum Detection and Estimation Theory*; Academic Press: New York, NY, USA, 1967.
35. Roy, S.M.; Braunstein, S.L. Exponentially enhanced quantum metrology. *Phys. Rev. Lett.* **2008**, *100*, 220501. [[CrossRef](#)] [[PubMed](#)]
36. Boixo, S.; Datta, A.; Davis, M.J.; Flammia, S.T.; Shaji, A.; Caves, C.M. Quantum metrology: Dynamics versus entanglement. *Phys. Rev. Lett.* **2008**, *101*, 040403. [[CrossRef](#)] [[PubMed](#)]
37. Ma, J.; Huang, Y.X.; Wang, X.; Sun, C.P. Quantum Fisher information of the Greenberger-Horne-Zeilinger state in decoherence channels. *Phys. Rev. A* **2011**, *84*, 022302. [[CrossRef](#)]
38. Monras, A.; Paris, M.G. Optimal quantum estimation of loss in bosonic channels. *Phys. Rev. Lett.* **2007**, *98*, 160401. [[CrossRef](#)]
39. Luo, S. Wigner-yanase skew information vs. quantum Fisher information. *Proc. Am. Math. Soc.* **2004**, *132*, 885–890. [[CrossRef](#)]
40. Luo, S. Quantum Fisher information and uncertainty relations. *Lett. Math. Phys.* **2000**, *53*, 243–251. [[CrossRef](#)]
41. Taddei, M.M.; Escher, B.M.; Davidovich, L.; de Matos Filho, R.L. Quantum Speed Limit for Physical Processes. *Phys. Rev. Lett.* **2013**, *110*, 050402. [[CrossRef](#)]
42. Sun, Z.; Ma, J.; Lu, X.-M.; Wang, X. Fisher information in a quantum-critical environment. *Phys. Rev. A* **2010**, *82*, 022306. [[CrossRef](#)]
43. Chapeau-Blondeau, F. Entanglement-assisted quantum parameter estimation from a noisy qubit pair: A Fisher information analysis. *Phys. Lett. A* **2017**, *381*, 1369–1378. [[CrossRef](#)]
44. Abdel-Khalek, S.; Khalil, E.M.; Ali, S.I. Entanglement of a two-level atom papered in a finite Trio Coherent state. *Laser Phys.* **2008**, *18*, 135–143. [[CrossRef](#)]
45. Abdel-Khalek, S.; Obada, A.S.F. New features of Wehrl entropy and Wehrl PD of a single-Cooper pair box placed inside a dissipative cavity. *Ann. Phys.* **2010**, *325*, 2542–2549. [[CrossRef](#)]
46. Obada, A.-S.F.; Abdel-Khalek, S. Entanglement evaluation with atomic Fisher information. *Physica A* **2010**, *389*, 891–898. [[CrossRef](#)]
47. Obada, A.-S.F.; Abdel-Khalek, S.; Plastino, A. Information quantifiers’s description of weak field vs. strong field dynamics for a trapped ion in a laser field. *Physica A* **2011**, *390*, 525–533. [[CrossRef](#)]
48. Abdel-Khalek, S.; Berrada, K.; Obada, A.S.F. An investigation of quantum Fisher information of a single qubit system. *Eur. Phys. J. D* **2012**, *66*, 69. [[CrossRef](#)]
49. Berrada, K.; Abdel-Khalek, S.; Obada, A.S.F. Quantum Fisher information for a qubit system placed inside a dissipative cavity. *Phys. Lett. A* **2012**, *376*, 1412–1416. [[CrossRef](#)]
50. Li, X.-S.; Lin, D.L.; Gong, C.-D. Nonresonant interaction of a three-level atom with cavity fields. I. General formalism and level occupation probabilities. *Phys. Rev. A* **1987**, *36*, 5209–5219. [[CrossRef](#)]
51. Abdel-Wahab, N.H. The general formalism for a three-level atom interacting with a two-mode cavity field. *Phys. Scr.* **2007**, *76*, 233. [[CrossRef](#)]
52. Abdel-Wahab, N.H. A three-level atom interacting with a single mode cavity field: Different configurations. *Phys. Scr.* **2007**, *76*, 244. [[CrossRef](#)]
53. Obada, A.-S.F.; Eied, A.A.; Abd Al-Kader, G.M. Entanglement of a general formalism-type three-level atom interacting with a non-correlated two-mode cavity field in the presence of nonlinearities. *J. Phys. B* **2008**, *41*, 195503. [[CrossRef](#)]

54. Teng, J.-H.; Wang, H.-F.; Qi, L.N.; Zhang, S. Influence of Kerr medium on entanglement of Cascade-type three-level atoms and a bimodal cavity field. *Int. J. Theor. Phys.* **2009**, *48*, 2818–2825. [[CrossRef](#)]
55. Mortezapoura, A.; Mahmoudib, M.; Khajehpour, M.R.H. Atom–photon, two-mode entanglement and two-mode squeezing in the presence of cross-Kerr nonlinearity. *Opt. Quantum. Electron.* **2015**, *47*, 2311–2329. [[CrossRef](#)]
56. Metwally, N.; Eleuch, H.; Obada, A.-S. Sudden death and rebirth of entanglement for different dimensional systems driven by a classical random external field. *Laser Phys. Lett.* **2016**, *13*, 105206. [[CrossRef](#)]
57. Faghihnia, M.J.; Tavassolya, M.K.; Hatamid, M. Dynamics of entanglement of a three-level atom in motion interacting with two coupled modes including parametric down conversion. *Physica A* **2014**, *407*, 100–109. [[CrossRef](#)]
58. Ismail, M.K.; El-Shahat, T.M. The damped interaction between a single-mode cavity field with Caldirola–Kanai Hamiltonian and a three-level atom. *Chin. J. Phys.* **2019**, *59*, 273–280. [[CrossRef](#)]
59. El-Wahab, N.A.; Rady, A.A.; Osman, A.N.A.; Salah, A. Influence of the gravitational field on the statistics of a three-level atom interacting with a one-mode cavity field. *J. Russ. Laser Res.* **2015**, *36*, 423–429. [[CrossRef](#)]
60. Abdel-Aty, M.; Obada, A.-S.F. Engineering entanglement of a general three-level system interacting with a correlated two-mode nonlinear coherent state. *Eur. Phys. J. D* **2003**, *23*, 155–165. [[CrossRef](#)]
61. Abdel-Wahab, N.H.; Salah, A. On the interaction between a time-dependent field and a two-level atom. *Mod. Phys. Lett. A* **2019**, *34*, 1950081. [[CrossRef](#)]
62. Milburn, G.J. Intrinsic decoherence in quantum mechanics. *Phys. Rev. A* **1991**, *44*, 5401. [[CrossRef](#)]
63. Gardiner, C.W. *Quantum Noise*; Springer: Berlin, Germany, 1991.
64. Ghirardi, G.C.; Rimini, A.; Weber, T. Unified dynamics for microscopic and macroscopic systems. *Phys. Rev. D* **1986**, *34*, 470. [[CrossRef](#)]
65. Diosi, L. Models for universal reduction of macroscopic quantum fluctuations. *Phys. Rev. A* **1989**, *40*, 1165. [[CrossRef](#)]
66. Ellis, J.; Mohanty, S.; Nanopoulos, D.V. Wormholes violate quantum mechanics in SQUIDs. *Phys. Lett. B* **1990**, *235*, 305. [[CrossRef](#)]
67. Anwar, S.J.; Ramzan, M.; Khan, M.K. Dynamics of entanglement and quantum Fisher information for N-level atomic system under intrinsic decoherence. *Quantum Inf. Process.* **2017**, *16*, 142. [[CrossRef](#)]
68. Zheng, L.; Zhang, G.F. Intrinsic decoherence in Jaynes-Cummings model with Heisenberg exchange interaction. *Eur. Phys. J. D* **2017**, *71*, 288. [[CrossRef](#)]
69. Liao, Y.Y.; Jian, S.R.; Lee, J.R. Effect of intrinsic decoherence on entanglement of three polar molecules with two-dimensional rotation. *Eur. Phys. J. D* **2019**, *73*, 47. [[CrossRef](#)]
70. Obada, A.S.F.; Mohamed, A.B.A. Quantum correlations of two non-interacting ion’s internal electronic states with intrinsic decoherence. *Opt. Commun.* **2013**, *309*, 236–241. [[CrossRef](#)]
71. Mohamed, A.-B. Non-local correlation and quantum discord in two atoms in the non-degenerate model. *Ann. Phys.* **2012**, *327*, 3130–3137. [[CrossRef](#)]
72. Mohamed, A.B.A.; Eleuch, H. Optical tomography dynamics induced by qubit-resonator interaction under intrinsic decoherence. *Sci. Rep.* **2022**, *12*, 17162. [[CrossRef](#)]
73. Mohamed, A.A.; Hessian, H.A.; Al-Duais, F.S.; Eleuch, H. Quantum coherence induced by a flux qubit coupled by a resonator coherent field through a two-photon interaction. *Phys. Scr.* **2021**, *96*, 125120. [[CrossRef](#)]
74. Mohamed, A.-B.A.; Hessian, H.A. Non-classicality in an open two-mode parametric amplifier cavity containing a qutrit system. *Phys. Scr.* **2021**, *96*, 055102. [[CrossRef](#)]
75. Alliluev, S.P.; Malkin, I.A. Calculations of the Stark effect in hydrogen atoms by using the dynamical symmetry $O(2, 2) \times 278$. *Zh. Eksp. Teor. Fiz.* **1974**, *66*, 1283–1294.
76. Al Naim, A.F.; Khan, J.Y.; Khalil, E.M.; Addel-Khalek, S. Effects of Kerr medium and Stark shift parameter on Wehrl entropy and the field purity for two-photon Jaynes-Cummings model under dispersive approximation. *J. Russ. Laser Res.* **2019**, *40*, 20–29. [[CrossRef](#)]
77. Hilal, E.M.A.; Khalil, E.M. Quantum statistical aspects of interactions between the radiation field and two entangled two-level atoms in the presence of Stark shift terms. *J. Russ. Laser Res.* **2018**, *39*, 207–215. [[CrossRef](#)]
78. Anwar, S.J.; Ramzan, M.; Usman, M.; Khan, M.K. Entanglement dynamics of three and four level atomic system under Stark effect and Kerr-like medium. *Quantum Rep.* **2019**, *1*, 23–36. [[CrossRef](#)]
79. Anwar, S.J.; Ramzan, M.; Khan, M.K. Effect of Stark and Kerr-like medium on the entanglement dynamics of two three-level atomic system. *Quantum Inf. Process.* **2019**, *18*, 192. [[CrossRef](#)]
80. Anwar, S.J.; Ramzan, M.; Usman, M.; Khan, M.K. Stark and Kerr effects on the dynamics of moving N-level atomic system. *J. Quantum Inf.* **2019**, *9*, 22. [[CrossRef](#)]
81. Abdel-Aty, M.; Furuichi, S.; Obada, A.F. Entanglement degree of a nonlinear multiphoton Jaynes-Cummings model. *J. Opt. B Quantum Semiclass. Opt.* **2002**, *4*, 37. [[CrossRef](#)]
82. Baghshahi, H.R.; Tavassoly, M.K.; Behjat, A. Entropy squeezing and atomic inversion in the k-photon Jaynes Cummings model in the presence of the Stark shift and a Kerr medium: A full nonlinear approach. *Chin. Phys. B* **2014**, *23*, 074203. [[CrossRef](#)]
83. Golkar, S.; Tavassoly, M. Dynamics and maintenance of bipartite entanglement via the Stark shift effect inside dissipative reservoirs. *Laser Phys. Lett.* **2018**, *15*, 035205. [[CrossRef](#)]
84. Obada, A.F.; Abdel-Khalek, S.; Khalil, E.M.; Ali, S.I. Effects of Stark shift and decoherence terms on the dynamics of phase-space entropy of the multiphoton Jaynes Cummings model. *Phys. Scr.* **2012**, *86*, 055009. [[CrossRef](#)]

85. Abdel-Khalek, S. Quantum Fisher information for moving three-level atom. *Quantum Inf. Process.* **2013**, *12*, 3761–3769. [[CrossRef](#)]
86. Enaki, N.A.; Ciobanu, N.J. Quantum trapping conditions for three-level atom flying through bimodal cavity field. *Mod. Opt.* **2008**, *55*, 589–598. [[CrossRef](#)]
87. Guo, J.L.; Song, H.S. Entanglement between two Tavis–Cummings atoms with phase decoherence. *J. Mod. Opt.* **2009**, *56*, 496–501. [[CrossRef](#)]
88. Lu, X.; Wang, X.; Sun, C.P. Quantum Fisher information flow and non-Markovian processes of open systems. *Phys. Rev. A* **2010**, *82*, 042103. [[CrossRef](#)]
89. Barndorff-Nielsen, O.E.; Gill, R.D.; Jupp, P.E. On quantum statistical inference. *J. R. Stat. Soc. B* **2003**, *65*, 775–816. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.