

Article

Shannon, Rényi, Tsallis Entropies and Onicescu Information Energy for Low-Lying Singly Excited States of Helium

Jen-Hao Ou ^{1,2}*^D and Yew Kam Ho¹

- ¹ Institute of Atomic and Molecular Sciences, Academia Sinica, Taipei 10617, Taiwan
- ² Department of Chemistry, National Taiwan University, Taipei 10617, Taiwan
- * Correspondence: b02203025@ntu.edu.tw

Received: 10 June 2019 ; Accepted: 11 July 2019; Published: 18 July 2019



Abstract: Knowledge of the electronic structures of atomic and molecular systems deepens our understanding of the desired system. In particular, several information-theoretic quantities, such as Shannon entropy, have been applied to quantify the extent of electron delocalization for the ground state of various systems. To explore excited states, we calculated Shannon entropy and two of its one-parameter generalizations, Rényi entropy of order α and Tsallis entropy of order α , and Onicescu Information Energy of order α for four low-lying singly excited states ($1s2s \, {}^{1}S^{e}$, $1s2s \, {}^{3}S^{e}$, $1s3s \, {}^{1}S^{e}$, and $1s3s \, {}^{3}S^{e}$ states) of helium. This paper compares the behavior of these three quantities of order 0.5 to 9 for the ground and four excited states. We found that, generally, a higher excited state had a *larger* Rényi entropy, *larger* Tsallis entropy, and *smaller* Onicescu information energy. However, this trend was not definite and the singlet–triplet reversal occurred for Rényi entropy, Tsallis entropy at a certain range of order α .

Keywords: Shannon entropy; Rényi entropy; Tsallis entropy; Onicescu information energy; singly excited states; helium; Hylleraas wave functions; singlet–triplet reversal

1. Introduction

The electronic structure, a unique feature of an atomic or molecular system, is fundamental for further understanding of other properties. To characterize an electronic structure, one critical aspect is whether the electron distribution is localized or delocalized [1,2]. Originated from the information theory and later extended to the position space, the *information-theoretic (IT) quantities* provide a possible quantitative account of the extent of electron localization. For instance, the more delocalized is the electron distribution, the larger is the Shannon entropy (*S*) [3–14]. Moreover, Rényi entropy of order α (R_{α}) [15–17] and Tsallis entropy of order α (T_{α}) [18–21] are the one-parameter generalizations of Shannon entropy and offer extended descriptions. In addition, Onicescu Information Energy of order α (E_{α}^{O}) [22,23] is related to the frequency moment of the electron density [24–26]. These IT quantities have been applied to various atomic and molecular systems [4–14,27–80] and potentials [81–91]. Further background details of these IT quantities were summarized by Ilić and Stanković [92] and by us [80].

However, a large part of the reports focus on the ground state, to the best of our knowledge. Even though some results for Rydberg states [10,40,46,62–64,81] are available, behaviors of these IT quantities for excited states remain to be explored. In particular, for the helium atom, Shannon entropy generally increases for higher excited states [10–12] but several exceptions are also found [10]. To examine the trend in IT quantities for increasing excitations, we calculated Shannon, Rényi, and Tsallis entropies and Onicescu Information Energy for four low-lying singly excited states (1s2s ${}^{1}S^{e}$, 1s2s ${}^{3}S^{e}$, 1s3s ${}^{1}S^{e}$, and 1s3s ${}^{3}S^{e}$ states) of helium, subsequent to our previous benchmark report for the



ground state helium [80]. This paper compares the behavior of these quantities of order 0.5 to 9 for the ground and four excited states of helium.

2. Theoretical Methods

2.1. Determination of Ground and Singly Excited States of Helium via Hylleraas Wave Functions

The non-relativistic Hamiltonian of helium atom (in atomic units) is

$$H = -\frac{1}{2}\nabla_1^2 - \frac{1}{2}\nabla_2^2 - \frac{2}{r_1} - \frac{2}{r_2} + \frac{1}{r_{12}},$$

where subscripts 1 and 2 are indexes of electrons, and r_{12} is the distance between the two electrons. Featuring the explicit r_{12} term, Hylleraas wave functions [93–95] take the electron correlation into account [96,97] and are written as

$$\psi_{mnk}(r_1, r_2, r_{12}) = \sum_{mnk} C_{mnk}(e^{-\beta_1 r_1} e^{-\beta_2 r_2} r_1^m r_2^n r_{12}^k + e^{-\beta_1 r_2} e^{-\beta_2 r_1} r_2^m r_1^n r_{12}^k)$$

for singlet states, and

$$\psi_{mnk}(r_1, r_2, r_{12}) = \sum_{mnk} C_{mnk}(e^{-\beta_1 r_1} e^{-\beta_2 r_2} r_1^m r_2^n r_{12}^k - e^{-\beta_1 r_2} e^{-\beta_2 r_1} r_2^m r_1^n r_{12}^k)$$

for triplet states. *m*, *n*, *k*, and ω are zero or positive integers and $m + n + k \leq \omega$ so ω determines the size of basis set. Besides, β_1 and β_2 are variational parameters. For singlet states with arbitrary β_1 and β_2 , and also for triplet states with $\beta_1 \neq \beta_2$, $m \geq n$ and ω starts from zero. On the other hand, for triplet states with $\beta_1 = \beta_2$, m > n and ω counts from one. The restrictions above are summarized in Table 1.

Spin Multiplicity Of The State	Relation between eta_1 and eta_2	Restriction on <i>m</i> and <i>n</i>	ω Starts from
singlet	$eta_1 eq eta_2$		
singlet	$eta_1=eta_2$	$m \ge n$	$\omega \geq 0$
triplet	$eta_1 eq eta_2$		
triplet	$\beta_1 = \beta_2$	m > n	$\omega \ge 1$

Table 1. Restrictions on *m*, *n* and ω .

2.2. Definitions for Electron Density

In the position space, Shannon, Rényi, and Tsallis entropies and Onicescu informational energy are defined in terms of the electron density. Nonetheless, different definitions of electron density exist in the literature and are described as follows. For a *N*-electron wave function $\Psi(\vec{x}_1, \vec{x}_2, \dots, \vec{x}_N)$, the *spinless one-electron density* [98], or *N*-normalized one-electron density, $\rho(\vec{r})$ is defined as

$$\rho(\vec{r}_1) \equiv N \int |\Psi(\vec{x}_1, \vec{x}_2, \cdots, \vec{x}_N)|^2 ds_1 d\vec{x}_2 \cdots d\vec{x}_N,$$

where \vec{x}_i is the spin s_i and space $\vec{r}_i = (r_i, \theta_i, \phi_i)$ coordinates of the *i*th electron. Integration of this electron density equals to the number of electrons *N* in the system as

$$\int \rho(\vec{r}) d\vec{r} = N$$

On the other hand, the *one-normalized one-electron density* $\sigma(\vec{r})$ is defined by dividing $\rho(\vec{r})$ by *N* as [99]

$$\sigma(\vec{r}) \equiv \frac{\rho(\vec{r})}{N}.$$

Therefore, in contrast to the former choice of normalizing the electron density $\rho(\vec{r})$ to N, $\sigma(\vec{r})$ is normalized to one as

$$\int \sigma(\vec{r}) d\vec{r} = 1.$$

This $\sigma(\vec{r})$ is also called *shape factor* [99,100], *shape function* [99,101], *unit-normalized density* [27], 1-*normalized density* [29], or *density per particle* [102,103]. In this work, we computed the radial part of one-normalized one-electron density $\sigma(r_1)$ according to Calais and Lowdin as [104]

$$\sigma(r_1) \equiv \frac{\rho(r_1)}{N} = \int |\psi(r_1, r_2, r_{12})|^2 d\vec{r_2}$$

= $\frac{2\pi}{r_1} \int_0^\infty r_2 dr_2 \int_{|r_1 - r_2|}^{r_1 + r_2} r_{12} dr_{12} |\psi(r_1, r_2, r_{12})|^2.$

2.3. Information-Theoretic Quantities in the Position Space

In this work, we examined four information-theoretic quantities, including Shannon entropy (*S*), Rényi entropy of order α (R_{α}), Tsallis entropy of order α (T_{α}), and Onicescu informational energy of order α (E_{α}^{O}). These quantities originated from probability and information theory, and were later extended to the position space by taking the electron density as the argument. Further background details of these IT quantities are summarized in our previous work [80]. The information-theoretic quantities for one-normalized one-electron density $\sigma(r)$ are defined as

$$\begin{split} S[\sigma(r)] &\equiv -\int_0^\infty \sigma(r) \ln \sigma(r) 4\pi r^2 \, dr, \\ R_\alpha[\sigma(r)] &\equiv \frac{1}{1-\alpha} \ln \left[\int_0^\infty \sigma(r)^\alpha 4\pi r^2 \, dr \right], \\ T_\alpha[\sigma(r)] &\equiv \frac{1}{\alpha-1} \left[1 - \int_0^\infty \sigma(r)^\alpha 4\pi r^2 \, dr \right], \\ E^{\rm O}_\alpha[\sigma(r)] &\equiv \frac{1}{\alpha-1} \int_0^\infty \sigma(r)^\alpha 4\pi r^2 \, dr. \end{split}$$

The integrals were solved by 600-point Gauss-Laguerre quadrature.

Although not calculated in this work, the information-theoretic quantities for *N*-normalized one-electron density $\rho(r)$ are defined similarly as

$$\begin{split} S[\rho(r)] &\equiv -\int_0^\infty \rho(r) \ln \rho(r) 4\pi r^2 \, dr, \\ R_\alpha[\rho(r)] &\equiv \frac{1}{1-\alpha} \ln \left[\int_0^\infty \rho(r)^\alpha 4\pi r^2 \, dr \right], \\ T_\alpha[\rho(r)] &\equiv \frac{1}{\alpha-1} \left[1 - \int_0^\infty \rho(r)^\alpha 4\pi r^2 \, dr \right], \\ E_\alpha^{\rm O}[\rho(r)] &\equiv \frac{1}{\alpha-1} \int_0^\infty \rho(r)^\alpha 4\pi r^2 \, dr, \end{split}$$

Formulas for conversion between IT quantities using the two different densities are [27,29,32,55–57,59,72]

$$\begin{split} S[\rho(r)] &= S[\sigma(r)]N - N\ln N, \\ R_{\alpha}[\rho(r)] &= R_{\alpha}[\sigma(r)] + \frac{\alpha}{1-\alpha}\ln N, \\ T_{\alpha}[\rho(r)] &= T_{\alpha}[\sigma(r)]N^{\alpha} + \frac{1-N^{\alpha}}{\alpha-1}, \\ E^{O}_{\alpha}[\rho(r)] &= E^{O}_{\alpha}[\sigma(r)]N^{\alpha}. \end{split}$$

3. Results and Discussion

All values are in atomic units. We first examined the accuracy of calculated energy to confirm the quality of wave functions used for calculating other quantities. Moreover, the self consistency of our information-theoretic quantities were examined. Then, we compared and analyzed the behavior of the IT quantities for the ground and singly excited states.

3.1. Accuracy of Presented Energy and Information-Theoretic Quantities

3.1.1. Energy *E*

In our previous benchmark work for the ground state helium [80], we let $\beta_1 = \beta_2$ and changed only one parameter. In contrast, here for singly excited states, we varied β_1 and β_2 independently (Table 2) with size of basis set $\omega = 15$ (444 terms). Our energies satisfactorily agree with other high precision results [105] (Table 3), demonstrating the reliability of our calculations.

Table 2. Variational parameters β_1 and β_2 for the ground and low-lying singly excited states of helium with size of basis set $\omega = 15$.

State	β_1	β_2	State	β_1	β_2
1s1s ¹ S ^e	2.00	2.00			
1s2s ¹ S ^e	1.19	2.80	1s2s ³ <i>S</i> ^e	2.09	0.55
1s3s ¹ S ^e	0.74	1.94	1s3s ³ S ^e	0.78	1.90

Table 3. Energy *E* and Shannon entropy *S* for the ground and low-lying singly excited states of helium with size of basis set $\omega = 15$.

State	Ε	S
$1s^{2} S^{2}$	-2.903724371	2.705 102 85
Drake [105]	-2.9037243770341195	
Lin and Ho [52]	-2.9037243768	2.7051028
Ou and Ho [11]	-2.903724371	2.70510285
Restrepo Cuartas and Sanz-Vicario [10]	-2.903605	2.705
$1s2s^1S^e$	-2.145974042	5.491 968 78
Drake [105]	-2.145974046054419(6)	
Lin and Ho [106]	-2.145974042	
Ou and Ho [11]	-2.145974012	5.491 968 37
Katriel et al. [107]	-2.1459740	
Restrepo Cuartas and Sanz-Vicario [10]	-2.145967	5.492
$1s2s^3S^e$	-2.175229377	5.23597814
Drake [105]	-2.17522937823679130	
Lin and Ho [106]	-2.17522937822	
Katriel et al. [107]	-2.1752294	
Restrepo Cuartas and Sanz-Vicario [10]	-2.175229	5.236
1s3s ¹ S ^e	-2.061 271 958	6.769 321 78
Drake [105]	-2.061271989740911(5)	
Lin and Ho [106]	-2.06127196	
Restrepo Cuartas and Sanz-Vicario [10]	-2.061270	6.769
1s3s ³ S ^e	-2.068 689 066	6.605 067 28
Drake [105]	-2.06868906747245719	
Lin and Ho [106]	-2.06868906	
Restrepo Cuartas and Sanz-Vicario [10]	-2.068689	6.605

3.1.2. Shannon Entropy *S*, Rényi Entropy R_{α} , Tsallis Entropy T_{α} and Onicescu Information Energy E_{α}^{O}

Our Shannon entropy well agreed with the literature [10,11,52,80] (Table 3). Moreover, we tested the self consistency by seeing if our values satisfy the analytic relations between them as follows. Let us take the 1s2s ${}^{1}S^{e}$ state for example (Figures 1 and 2). First, by definition, both Rényi entropy R_{α} and Tsallis entropy T_{α} recover Shannon entropy when the order α goes to one as

$$\lim_{\alpha \to 1} R_{\alpha} = S,$$

$$\lim_{\alpha \to 1} T_{\alpha} = S.$$
(1)

Our numerical values obey this requirement because the curves of Rényi entropy and Tsallis entropy intersect each other at the value of Shannon entropy when order goes to one (Figure 1).



Figure 1. Rényi entropy R_{α} and Tsallis entropy T_{α} of order 0.5 to 9 of $1s2s {}^{1}S^{e}$ state. Both recover the value of Shannon entropy as order α approaches one. Selected values are in Tables 4 and 5 and full data are in Tables S1 and S2.

Second, the sum of Tsallis entropy T_{α} and Onicescu informational energy E_{α}^{O} equals to $1/(\alpha - 1)$ as

$$T_{\alpha} + E_{\alpha}^{\rm O} = \frac{1}{\alpha - 1} \tag{2}$$

This phenomenon is best seen in Figure 2. When the order α decreases toward 0.5, T_{α} increases, E_{α}^{O} decreases, and their sum $1/(\alpha - 1)$ tends to -2. Furthermore, when the order is close α to one, Onicescu informational energy diverges, but the sum of Tsallis entropy and Onicescu informational energy still equals $1/(\alpha - 1)$.

Third, Rényi entropy R_{α} , Tsallis entropy T_{α} and Onicescu informational energy E_{α}^{O} are related by the following relations: [18,80]

$$R_{\alpha} = \frac{\ln[1 + T_{\alpha}(1 - \alpha)]}{1 - \alpha},$$
$$R_{\alpha} = \frac{\ln[E_{\alpha}^{O}(\alpha - 1)]}{1 - \alpha}.$$

Our numerical values fulfill these analytic relations between them.



Figure 2. Tsallis entropy T_{α} and Onicescu informational energy E_{α}^{O} of order 0.50 to 1.50 of 1s2s ${}^{1}S^{e}$ state. The sum of Tsallis entropy and Onicescu Information Energy equals $1/(\alpha - 1)$. Selected values are in Tables 5 and 6 and full data are in Tables S2 and S3.

3.2. Comparison of Shannon Entropy S of the Ground and Four Singly Excited States

We first analyzed the behavior of Shannon entropy, from which Rényi entropy and Tsallis entropy were generalized. As shown in Table 3, the higher is the energy, the larger is the Shannon entropy [10–12]. However, although this trend was observed for the presented *S*-type states of helium, several exceptions have been reported if the states to be compared have different total orbital angular momentum *L* [10].

On the other hand, Shannon entropy reflects the extent of electron delocalization [10–12]. In particular, for a given configuration, the electron density of the triplet state is shift toward and more concentrated to the origin than that of the singlet (Figure 3). As a result, the triplet state has a lower Shannon entropy than the singlet. Restrepo Cuartas and Sanz-Vicario called this phenomenon the *Entropic Hund's first rule* [10].



Figure 3. Radial probability density $\sigma(r)4\pi r^2$ for the ground and low-lying singly excited states of helium.

3.3. Comparison of Rényi Entropy R_{α} , Tsallis Entropy T_{α} and Onicescu Information Energy E_{α}^{O} of the Ground and Four Singly Excited States

Now, we move on to Rényi entropy R_{α} , Tsallis entropy T_{α} , and Onicescu Information Energy E_{α}^{O} . For simplicity, we consider the singlet states or the triplet states individually, and then compare the singlet with the triplet for a given configuration in the next subsection.

Within either the series of singlets $(1s^{2} {}^{1}S^{e}, 1s2s {}^{1}S^{e}$ and $1s3s {}^{1}S^{e}$ states) or the series of triplets $(1s2s {}^{3}S^{e} \text{ and } 1s3s {}^{3}S^{e} \text{ states})$, higher excited states had *larger* Rényi entropy R_{α} (Figure 4a) and Tsallis entropy T_{α} (Figure 4b), and *smaller* Onicescu Information Energy (Figure 4c) throughout order α from 0.5 to 9. Furthermore, for a given IT quantity, the curves of different configurations gradually separate when the order α decreases toward 0.5. One possible explanation is that the electron distribution of higher excited states oscillated more (Figure 3), and this behavior may be highlighted at a certain range of order α .



Figure 4. (a) Rényi entropy; (b) Tsallis entropy; and (c) Onicescu Information Energy of order 0.5 to 9 for the ground and low-lying singly excited states of helium with size of basis set $\omega = 15$. Selected values are in Tables 4–6. Full data are in Tables S1–S3.

3.4. Singlet–Triplet Reversal of Rényi entropy R_{α} , Tsallis entropy T_{α} and Onicescu Information Energy E_{α}^{O}

Here we compare the IT quantities of the singlet state with those of the corresponding triplet state. At fist glance, for a given configuration, the singlet state generally had a *larger* Rényi entropy, *larger* Tsallis entropy and *smaller* Onicescu Information Energy than the triplet state (Figure 4). However, this ordering was not definite and may even be reversed. To be more specific, the difference between an IT quantity of the singlet state and that of the corresponding triplet state are in Figure 5a for the 1s2s and in Figure 5b for the 1s3s configuration. The singlet–triplet reversal occurred at $1.71 \le \alpha \le 2.2$ for the 1s2s and at $1.52 \le \alpha \le 2.8$ for the 1s3s configuration.



Figure 5. Singlet–triplet reversal of Rényi entropy R_{α} , Tsallis entropy T_{α} and Onicescu Information Energy E_{α}^{O} for: (a) 1s2s; and (b) 1s3s configuration.

Moreover, the difference between Onicescu Information Energy of the singlet state and that of the triplet state has opposite sign to the difference for Tsallis entropy (Figure 5). This behavior can be readily verified by

$$E_{\alpha}^{O}(\text{singlet}) - E_{\alpha}^{O}(\text{triplet})$$

= $[\frac{1}{\alpha - 1} - T_{\alpha}(\text{singlet})] - [\frac{1}{\alpha - 1} - T_{\alpha}(\text{triplet})]$
= $-[T_{\alpha}(\text{singlet}) - T_{\alpha}(\text{triplet})].$

Selected values for Rényi, Tsallis entropy and Onicescu Information Energy for the low-lying singly excited states of helium are in Tables 4–6, respectively, and the full data are in Tables S1–S3.

Additionally, singlet–triplet reversal has also been reported for the interacting repulsive potential energy between two electrons in singly-excited states in quantum dots under certain confinement conditions [107,108], and in singly-excited states of helium-like ions under screened Coulomb interacting potentials with certain screening strengths and atomic numbers Z [109].

Table 4. Selected values for Rényi entropy of order α (R_{α}) for the low-lying singly excited states of helium with size of basis set ω = 15. Values of R_1 are set to Shannon entropy. Full data are in Table S1.

	1s2s		1s3s	
α -	$^{1}S^{e}$	³ S ^e	$^{1}S^{e}$	³ S ^e
0.5	7.827 360 64	7.375 384 22	1.00515457×10^{1}	9.750 363 90
1	5.491 968 78	5.235 978 14	6.769 321 78	6.605 067 28
2	2.46576126	2.46773641	2.513 498 59	2.51571175
3	1.707 786 50	1.702 788 37	1.740 513 92	1.740 268 03
4	1.33674386	1.329 418 93	1.365 026 03	1.364 034 09
9	6.37201665×10^{-1}	6.28474643×10^{-1}	6.60262552×10^{-1}	6.58729039×10^{-1}

Table 5. Selected values for Tsallis entropy of order α (T_{α}) for low-lying singly excited states of helium with size of basis set $\omega = 15$. Values of T_1 are set to Shannon entropy. Full data are in Table S2.

α -	1s2s		1s3s	
	${}^{1}S^{e}$	³ S ^e	$^{1}S^{e}$	³ S ^e
0.5	9.81658692×10^{1}	7.79050690×10^{1}	3.02575822×10^2	$2.59995972 imes 10^2$
1	5.491 968 78	5.235 978 14	6.769 321 78	6.605 067 28
2	$9.15055847 imes10^{-1}$	$9.15223459 imes10^{-1}$	$9.19015588 imes10^{-1}$	$9.19194622 imes 10^{-1}$
3	$4.83571213 imes10^{-1}$	$4.83406163 imes10^{-1}$	$4.84612118 imes10^{-1}$	4.84604549×10^{-1}
4	$3.27290267 imes 10^{-1}$	$3.27156002 imes 10^{-1}$	$3.27781851 imes10^{-1}$	$3.27765306 imes10^{-1}$
9	$1.24236085 imes10^{-1}$	$1.24180846 imes 10^{-1}$	$1.24364781 imes10^{-1}$	$1.24356940 imes 10^{-1}$

Table 6. Selected values for Onicescu Information Energy of order α (E_{α}^{O}) for low-lying singly excited states of helium with size of basis set $\omega = 15$. E_{α}^{O} is undefined for $\alpha = 1$. Full data are in Table S3.

α -	1s2s		1s3s	
	$^{1}S^{e}$	³ S ^e	$^{1}S^{e}$	³ S ^e
0.5	-1.00165869×10^2	-7.99050690×10^{1}	-3.04575822×10^2	-2.61995972×10^2
2	8.49441529×10^{-2}	$8.47765409 imes10^{-2}$	$8.09844111 imes10^{-2}$	$8.08053779 imes10^{-2}$
3	$1.64287869 imes 10^{-2}$	$1.65938368 imes 10^{-2}$	$1.53878810 imes 10^{-2}$	$1.53954502 imes 10^{-2}$
4	$6.04306538 imes10^{-3}$	$6.17733033 imes 10^{-3}$	$5.55148184 imes 10^{-3}$	$5.56802665 imes10^{-3}$
9	7.63914364×10^{-4}	8.19153809×10^{-4}	6.35218223×10^{-4}	6.43059145×10^{-4}

4. Conclusions

Generally, a higher excited state of helium had a *larger* Shannon entropy, *larger* Rényi entropy of order α , *larger* Tsallis entropy of order α , and *smaller* Onicescu information energy of order α . This behavior possibly reflects the more delocalized nature of electron distribution of higher excited states. However, for Rényi entropy, Tsallis entropy and Onicescu information energy, this trend was not definite and singlet–triplet reversal occurred at a certain range of order α . In the future, the knowledge of bounds and asymptotic analysis of these quantities may provide additional understanding of this behavior [110–116].

On the other hand, these IT quantities not only reflect the extent of electron delocalization, as discussed in this article, but also have numerous applications. For instance, Rényi entropy has been widely used to examine quantum entanglement [117–123]. Furthermore, Tsallis statistics is relevant to certain systems of ultracold atoms [124–127]. Moreover, these IT quantities are applied to investigate electron correlation [30,31,33,39,72,128–136] and various molecular properties as

well [28,36,45,48,49,59–61,72,73]. Calculated with the highly correlated Hylleraas wave functions, the results in this work then serve as a useful and reliable reference for the various applications mentioned above.

As for a theoretical application, our present calculations would also play a role to test the Collins Conjecture [137]—correlation energy (the difference between the exact energy and Hartree–Fock energy) is directly proportional to entropy—when one could extend the calculations to other helium-like ions. Thus, such a conjecture was tested by Dehesa and coworkers [138,139] to be valid for von Neumann entropy in helium iso-electronic sequence. It would be of interest to test the Collins Conjecture for Shannon, Renyi, or Tsallis entropy to be valid for helium sequence using highly correlated wave functions.

Finally, it should be mentioned that, to the best of our knowledge, there is no formula relating entropy to the quantum number of any given excited state, in a manner similar to the formula relating quantum defect to the effective quantum number for such an excited state. Nevertheless, this is an interesting issue and it is worthwhile to explore further in future investigations.

Supplementary Materials: Full data for Rényi, Tsallis entropies and Onicescu Information Energy of order 0.5 to 9 for the ground and low-lying singly excited states of helium are in Table S1 to S3 and are available online at https://www.mdpi.com/2218-2004/7/3/70/s1.

Author Contributions: Both authors made equal contributions.

Acknowledgments: J.-H.O. thanks Yen-Chang Lin and Chien-Hao Lin for helpful discussions, and Roy Yeh for suggestions about artwork. Our work was supported by the Ministry of Science and Technology, Taiwan.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Chalvet, O.; Daudel, R.; Diner, S.; Malrieu, J.P. (Eds.) Localization and Delocalization in Quantum Chemistry Volume I Atoms and Molecules in the Ground State; Springer: Dordrecht, The Netherlands, 1975; Volume 1. [CrossRef]
- 2. Chalvet, O.; Daudel, R.; Diner, S.; Malrieu, J.P. (Eds.) *Localization and Delocalization in Quantum Chemistry Volume II Ionized and Excited States*; Springer: Dordrecht, The Netherlands, 1976; Volume 2. [CrossRef]
- 3. Shannon, C.E. A Mathematical Theory of Communication. Bell Syst. Tech. J. 1948, 27, 379–423. [CrossRef]
- 4. Aslangul, C.; Constanciel, R.; Daudel, R.; Kottis, P. Aspects of the Localizability of Electrons in Atoms and Molecules: Loge Theory and Related Methods. *Adv. Quantum Chem.* **1972**, *6*, 93–141. [CrossRef]
- 5. Sen, K.D. Characteristic features of Shannon information entropy of confined atoms. *J. Chem. Phys.* 2005, 123, 074110. [CrossRef] [PubMed]
- Corzo, H.H.; Laguna, H.G.; Sagar, R.P. Localization-delocalization phenomena in a cyclic box. *J. Math. Chem.* 2012, 50, 233–248. [CrossRef]
- 7. Cedillo, A. Comment on "Localization-delocalization phenomena in a cyclic box" by H. H. Corzo, H. G. Laguna, and R. P. Sagar. *J. Math. Chem.* **2017**, *55*, 1889–1892. [CrossRef]
- 8. Corzo, H.H.; Castaño, E.; Laguna, H.G.; Sagar, R.P. Measuring localization-delocalization phenomena in a quantum corral. *J. Math. Chem.* **2013**, *51*, 179–193. [CrossRef]
- 9. Laguna, H.G.; Sagar, R.P. Quantum uncertainties of the confined Harmonic Oscillator in position, momentum and phase-space. *Ann. Phys.* **2014**, *526*, 555–566. [CrossRef]
- 10. Restrepo Cuartas, J.P.; Sanz-Vicario, J.L. Information and entanglement measures applied to the analysis of complexity in doubly excited states of helium. *Phys. Rev. A* **2015**, *91*, 052301. [CrossRef]
- 11. Ou, J.H.; Ho, Y.K. Shannon Information Entropy in Position Space for the Ground and Singly Excited States of Helium with Finite Confinements. *Atoms* **2017**, *5*, 15. [CrossRef]
- 12. Ou, J.H.; Ho, Y.K. Shannon information entropy in position space for doubly excited states of helium with finite confinements. *Chem. Phys. Lett.* **2017**, *689*, 116–120. [CrossRef]
- Rodriguez-Bautista, M.; Vargas, R.; Aquino, N.; Garza, J. Electron-density delocalization in many-electron atoms confined by penetrable walls: A Hartree-Fock study. *Int. J. Quantum Chem.* 2018, 118, e25571. [CrossRef]

- 14. Majumdar, S.; Mukherjee, N.; Roy, A.K. Information entropy and complexity measure in generalized Kratzer potential. *Chem. Phys. Lett.* **2019**, *716*, 257–264. [CrossRef]
- Rényi, A. On measures of entropy and information. In *Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability*; Neyman, J., Ed.; University of California Press: Berkeley, CA, USA, 1961; Volume 1, pp. 547–561.
- 16. Renyi, A. Probability Theory; North-Holland Publishing Company: Amsterdam, The Netherlands, 1970.
- Beadle, E.; Schroeder, J.; Moran, B.; Suvorova, S. An overview of Renyi Entropy and some potential applications. In Proceedings of the 2008 42nd Asilomar Conference on Signals, Systems and Computers, Pacific Grove, CA, USA, 26–29 October 2008; pp. 1698–1704. [CrossRef]
- 18. Tsallis, C. Possible generalization of Boltzmann-Gibbs statistics. J. Stat. Phys. 1988, 52, 479–487. [CrossRef]
- 19. Tsallis, C. Introduction to Nonextensive Statistical Mechanics; Springer: New York, NY, USA, 2009. [CrossRef]
- 20. Tsallis, C. The Nonadditive Entropy *S*_{*q*} and Its Applications in Physics and Elsewhere: Some Remarks. *Entropy* **2011**, *13*, 1765–1804. [CrossRef]
- 21. Tsallis, C. On the foundations of statistical mechanics. Eur. Phys. J. Spec. Top. 2017, 226, 1433–1443. [CrossRef]
- 22. Onicescu, O. Energie Informationnelle. *Comptes Rendus Hebd. Seances L'Academie Sci. Ser. A* 1966, 263, 841–842.
- 23. Iosifescu, M. Octav Onicescu. In *Statisticians of the Centuries*; Heyde, C.C., Seneta, E., Crépel, P., Fienberg, S.E., Gani, J., Eds.; Springer: New York, NY, USA, 2001; pp. 415–419. [CrossRef]
- 24. Angulo, J.C.; Romera, E.; Dehesa, J.S. Inverse atomic densities and inequalities among density functionals. *J. Math. Phys.* **2000**, *41*, 7906–7917. [CrossRef]
- 25. Romera, E.; Angulo, J.C.; Dehesa, J.S. The Hausdorff entropic moment problem. *J. Math. Phys.* 2001, 42, 2309. [CrossRef]
- 26. Romera, E.; Angulo, J.C.; Dehesa, J.S. Erratum: The Hausdorff entropic moment problem [J. Math. Phys. 42, 2309 (2001)]. J. Math. Phys. 2003, 44, 2354–2354. [CrossRef]
- 27. Hô, M.; Sagar, R.P.; Pérez-Jordá, J.M.; Smith, V.H.; Esquivel, R.O. A numerical study of molecular information entropies. *Chem. Phys. Lett.* **1994**, *219*, 15–20. [CrossRef]
- 28. Hô, M.; Smith, V.H.; Weaver, D.F.; Gatti, C.; Sagar, R.P.; Esquivel, R.O. Molecular similarity based on information entropies and distances. *J. Chem. Phys.* **1998**, *108*, 5469–5475. [CrossRef]
- 29. Fazal, S.P.; Sen, K.D.; Gutierrez, G.; Fuentealba, P. Shannon entropy of 1-normalized electron density. *Indian J. Chem., Sect. A* 2000, *39*, 48–49.
- 30. Massen, S.E. Application of information entropy to nuclei. Phys. Rev. C 2003, 67, 014314. [CrossRef]
- 31. Guevara, N.L.; Sagar, R.P.; Esquivel, R.O. Shannon-information entropy sum as a correlation measure in atomic systems. *Phys. Rev. A* **2003**, *67*, 012507. [CrossRef]
- 32. Sen, K.D.; De Proft, F.; Borgoo, A.; Geerlings, P. N-derivative of Shannon entropy of shape function for atoms. *Chem. Phys. Lett.* 2005, 410, 70–76. [CrossRef]
- 33. Guevara, N.L.; Sagar, R.P.; Esquivel, R.O. Local correlation measures in atomic systems. *J. Chem. Phys.* 2005, 122, 084101. [CrossRef]
- 34. Moustakidis, C.; Chatzisavvas, K.; Panos, C. Theoretical Quantum-information Properties of Nuclei and Trapped Bose Gases. *Int. J. Mod. Phys. E* 2005, *14*, 1087–1104. [CrossRef]
- 35. Chatzisavvas, K.C.; Moustakidis, C.C.; Panos, C.P. Information entropy, information distances, and complexity in atoms. *J. Chem. Phys.* **2005**, *123*, 174111. [CrossRef]
- 36. Gregori-Puigjané, E.; Mestres, J. SHED: Shannon Entropy Descriptors from Topological Feature Distributions. *J. Chem. Inf. Model.* **2006**, *46*, 1615–1622. [CrossRef]
- Mohajeri, A.; Dasmeh, P. Evaluating the Nature of Chemical Bonds Based on Probabilistic Models. *Int. J. Mod. Phys. C* 2007, *18*, 1795. [CrossRef]
- 38. Romera, E.; Nagy, Á. Rényi information of atoms. Phys. Lett. A 2008, 372, 4918–4922. [CrossRef]
- 39. Mohajeri, A.; Alipour, M. Information Energy As an Electron Correlation Measure in Atomic and Molecular Systems. *Int. J. Quantum Inf.* **2009**, *07*, 801–809. [CrossRef]
- Dehesa, J.S.; López-Rosa, S.; Martínez-Finkelshtein, A.; Yáñez, R.J. Information theory of D-dimensional hydrogenic systems: Application to circular and Rydberg states. *Int. J. Quantum Chem.* 2009, 110, 1529–1548. [CrossRef]
- 41. Nagy, Á.; Romera, E. Relative Rényi entropy for atoms. *Int. J. Quantum Chem.* **2009**, 109, 2490–2494. [CrossRef]

- 42. Nagy, Á.; Romera, E. Maximum Rényi entropy principle and the generalized Thomas-Fermi model. *Phys. Lett. A* **2009**, *373*, 844–846. [CrossRef]
- 43. Antolín, J.; Angulo, J.C. Complexity analysis of ionization processes and isoelectronic series. *Int. J. Quantum Chem.* **2009**, *109*, 586–593. [CrossRef]
- 44. Antolín, J.; López-Rosa, S.; Angulo, J. Renyi complexities and information planes: Atomic structure in conjugated spaces. *Chem. Phys. Lett.* **2009**, 474, 233–237. [CrossRef]
- 45. Delgado-Soler, L.; Toral, R.; Tomás, M.S.; Rubio-Martinez, J. RED: A set of molecular descriptors based on Rényi entropy. *J. Chem. Inf. Model.* **2009**, *49*, 2457–2468. [CrossRef] [PubMed]
- 46. López-Rosa, S.; Toranzo, I.V.; Sánchez-Moreno, P.; Dehesa, J.S. Entropy and complexity analysis of hydrogenic Rydberg atoms. *J. Math. Phys.* **2013**, *54*, 052109. [CrossRef]
- 47. Nagy, Á. Shannon entropy density as a descriptor of Coulomb systems. *Chem. Phys. Lett.* **2013**, *556*, 355–358. [CrossRef]
- 48. Molina-Espíritu, M.; Esquivel, R.O.; Kohout, M.; Angulo, J.C.; Dobado, J.A.; Dehesa, J.S.; LópezRosa, S.; Soriano-Correa, C. Insight into the informational-structure behavior of the Diels-Alder reaction of cyclopentadiene and maleic anhydride. *J. Mol. Model.* **2014**, *20*, 2631. [CrossRef] [PubMed]
- 49. Welearegay, M.A.; Balawender, R.; Holas, A. Information and complexity measures in molecular reactivity studies. *Phys. Chem. Chem. Phys.* **2014**, *16*, 14928–14946. [CrossRef] [PubMed]
- 50. Nagy, Á.; Romera, E. Relative Rényi entropy and fidelity susceptibility. *Europhys. Lett.* **2015**, *109*, 60002. [CrossRef]
- 51. Esquivel, R.O.; Molina-Espíritu, M.; López-Rosa, S.; Soriano-Correa, C.; Barrientos-Salcedo, C.; Kohout, M.; Dehesa, J.S. Predominant Information Quality Scheme for the Essential Amino Acids: An Information-Theoretical Analysis. *ChemPhysChem* **2015**, *16*, 2571–2581. [CrossRef] [PubMed]
- 52. Lin, C.H.; Ho, Y.K. Shannon information entropy in position space for two-electron atomic systems. *Chem. Phys. Lett.* **2015**, *633*, 261–264. [CrossRef]
- 53. Lin, C.H.; Ho, Y.K. Quantum Entanglement and Shannon Information Entropy for the Doubly Excited Resonance State in Positronium Negative Ion. *Atoms* **2015**, *3*, 422–432. [CrossRef]
- 54. Peng, H.T.; Ho, Y.K. Statistical Correlations of the N-particle Moshinsky Model. *Entropy* **2015**, *17*, 1882–1895. [CrossRef]
- 55. Rong, C.; Lu, T.; Ayers, P.W.; Chattaraj, P.K.; Liu, S. Scaling properties of information-theoretic quantities in density functional reactivity theory. *Phys. Chem. Chem. Phys.* **2015**, *17*, 4977–4988. [CrossRef]
- 56. Rong, C.; Lu, T.; Ayers, P.W.; Chattaraj, P.K.; Liu, S. Correction: Scaling properties of information-theoretic quantities in density functional reactivity theory. *Phys. Chem. Chem. Phys.* **2015**, *17*, 11110–11111. [CrossRef]
- 57. Liu, S.B.; Rong, C.Y.; Wu, Z.M.; Lu, T. Rényi entropy, Tsallis entropy and Onicescu information energy in density functional reactivity theory. *Acta Phys.-Chim. Sin.* **2015**, *31*, 2057–2063. [CrossRef]
- Esquivel, R.O.; López-Rosa, S.; Molina-Espíritu, M.; Angulo, J.C.; Dehesa, J.S. Information-theoretic space from simple atomic and molecular systems to biological and pharmacological molecules. *Theor. Chem. Acc.* 2016, 135, 253. [CrossRef]
- Liu, S.B. Information-Theoretic Approach in Density Functional Reactivity Theory. *Acta Phys.-Chim. Sin.* 2016, 32, 98–118. [CrossRef]
- Zhou, X.Y.; Rong, C.; Lu, T.; Zhou, P.; Liu, S. Information Functional Theory: Electronic Properties as Functionals of Information for Atoms and Molecules. *J. Phys. Chem. A* 2016, 120, 3634–3642. [CrossRef] [PubMed]
- 61. Alipour, M.; Safari, Z. From information theory to quantitative description of steric effects. *Phys. Chem. Chem. Phys.* **2016**, *18*, 17917–17929. [CrossRef] [PubMed]
- 62. Toranzo, I.V.; Dehesa, J.S. Rényi, Shannon and Tsallis entropies of Rydberg hydrogenic systems. *Europhys. Lett.* **2016**, *113*, 48003. [CrossRef]
- 63. Toranzo, I.V.; Puertas-Centeno, D.; Dehesa, J.S. Entropic properties of D-dimensional Rydberg systems. *Phys. A* **2016**, 462, 1197–1206. [CrossRef]
- 64. Dehesa, J.S.; Toranzo, I.V.; Puertas-Centeno, D. Entropic measures of Rydberg-like harmonic states. *Int. J. Quantum Chem.* **2017**, *117*, 48–56. [CrossRef]
- 65. Huang, Y.; Rong, C.; Zhang, R.; Liu, S. Evaluating frontier orbital energy and HOMO/LUMO gap with descriptors from density functional reactivity theory. *J. Mol. Model.* **2017**, *23*, 3. [CrossRef]

- 66. Farid, M.; Abdel-Hady, A.; Nasser, I. Comparative study of the scaling behavior of the Rényi entropy for He-like atoms. *J. Phys. Conf. Ser.* **2017**, *869*, 012011. [CrossRef]
- 67. Nasser, I.; Zeama, M.; Abdel-Hady, A. The Rényi entropy, a comparative study for He-like atoms using the exponential-cosine screened Coulomb potential. *Results Phys.* **2017**, *7*, 3892–3900. [CrossRef]
- 68. Zan, L.R.; Jiao, L.G.; Ma, J.; Ho, Y.K. Information-theoretic measures of hydrogen-like ions in weakly coupled Debye plasmas. *Phys. Plasmas* **2017**, *24*, 122101. [CrossRef]
- 69. Cao, X.; Rong, C.; Zhong, A.; Lu, T.; Liu, S. Molecular acidity: An accurate description with information-theoretic approach in density functional reactivity theory. *J. Comput. Chem.* **2018**, *39*, 117–129. [CrossRef] [PubMed]
- 70. Jiao, L.G.; Zan, L.R.; Zhang, Y.Z.; Ho, Y.K. Benchmark values of Shannon entropy for spherically confined hydrogen atom. *Int. J. Quantum Chem.* **2017**, *117*, e25375. [CrossRef]
- 71. Laguna, H.G.; Sagar, R.P. Information theoretical measures from cumulative and survival densities in quantum systems. *Int. J. Quantum Chem.* **2017**, *117*, e25387. [CrossRef]
- 72. Alipour, M.; Badooei, Z. Toward Electron Correlation and Electronic Properties from the Perspective of Information Functional Theory. *J. Phys. Chem. A* **2018**, *122*, 6424–6437. [CrossRef]
- 73. Alipour, M. Which information theoretic quantity should we choose for steric analysis of water nanoclusters $(H_2O)_n$ (n = 6, 32, 64)? *Acta Phys.-Chim. Sin.* **2018**, *34*, 407–413. [CrossRef]
- 74. Mukherjee, N.; Roy, A.K. Information-entropic measures in free and confined hydrogen atom. *Int. J. Quantum Chem.* **2018**, *118*, e25596. [CrossRef]
- 75. Mukherjee, N.; Roy, A.K. Information-entropic measures for non-zero *l* states of confined hydrogen-like ions. *Eur. Phys. J. D* **2018**, 72, 118. [CrossRef]
- 76. Mukherjee, N.; Roy, A.K. Information-Entropic Measures in Confined Isotropic Harmonic Oscillator. *Adv. Theory Simul.* **2018**, *1*, 1800090. [CrossRef]
- 77. Nasser, I.; Zeama, M.; Abdel-Hady, A. The nonadditive entropy for the ground state of helium-like ions using Hellmann potential. *Mol. Phys.* **2019**, 1612105. [CrossRef]
- 78. Zeama, M.; Nasser, I. Tsallis entropy calculation for non-Coulombic helium. *Phys. A* 2019, 528, 121468. [CrossRef]
- 79. López-Rosa, S.; Martín, A.L.; Antolín, J.; Angulo, J.C. Electron-pair entropic and complexity measures in atomic systems. *Int. J. Quantum Chem.* **2019**, *119*, e25861. [CrossRef]
- Ou, J.H.; Ho, Y.K. Benchmark calculations of Rényi, Tsallis entropies, and Onicescu information energy for ground state helium using correlated Hylleraas wave functions. *Int. J. Quantum Chem.* 2019, 119, e25928. [CrossRef]
- 81. Sánchez-Moreno, P.; Omiste, J.J.; Dehesa, J.S. Entropic functionals of Laguerre polynomials and complexity properties of the half-line Coulomb potential. *Int. J. Quantum Chem.* **2011**, *111*, 2283–2294. [CrossRef]
- 82. Sun, G.H.; Dong, S.H.; Saad, N. Quantum information entropies for an asymmetric trigonometric Rosen-Morse potential. *Ann. Phys.* **2013**, *525*, 934–943. [CrossRef]
- 83. Dong, S.; Sun, G.H.; Dong, S.H.; Draayer, J. Quantum information entropies for a squared tangent potential well. *Phys. Lett. A* **2014**, *378*, 124–130. [CrossRef]
- 84. Yahya, W.A.; Oyewumi, K.J.; Sen, K.D. Position and momentum information-theoretic measures of the pseudoharmonic potential. *Int. J. Quantum Chem.* **2015**, *115*, 1543–1552. [CrossRef]
- 85. Mukherjee, N.; Roy, A.; Roy, A.K. Information entropy as a measure of tunneling and quantum confinement in a symmetric double-well potential. *Ann. Phys.* **2015**, *527*, 825–845. [CrossRef]
- 86. Mukherjee, N.; Roy, A.K. Quantum confinement in an asymmetric double-well potential through energy analysis and information entropic measure. *Ann. Phys.* **2016**, *528*, 412–433. [CrossRef]
- Ghosal, A.; Mukherjee, N.; Roy, A.K. Information entropic measures of a quantum harmonic oscillator in symmetric and asymmetric confinement within an impenetrable box. *Ann. Phys.* 2016, 528, 796–818. [CrossRef]
- 88. Isonguyo, C.N.; Oyewumi, K.J.; Oyun, O.S. Quantum information-theoretic measures for the static screened Coulomb potential. *Int. J. Quantum Chem.* **2018**, *118*, e25620. [CrossRef]
- 89. Onate, C.A.; Ikot, A.N.; Onyeaju, M.C.; Ebomwonyi, O.; Idiodi, J.O.A. Effect of dissociation energy on Shannon and Rényi entropies. *Karbala Int. J. Mod. Sci.* **2018**, *4*, 134–142. [CrossRef]
- Olendski, O. Quantum Information Measures of the One-Dimensional Robin Quantum Well. Ann. Phys. 2018, 530, 1700324. [CrossRef]

- 91. Olendski, O. Rényi and Tsallis entropies: three analytic examples. Eur. J. Phys. 2019, 40, 025402. [CrossRef]
- 92. Ilić, V.M.; Stanković, M.S. A unified characterization of generalized information and certainty measures. *Phys. A* **2014**, *415*, 229–239. [CrossRef]
- 93. Hylleraas, E.A. Über den Grundzustand des Heliumatoms. Z. Phys. 1928, 48, 469–494. [CrossRef]
- 94. Hylleraas, E.A. Neue Berechnung der Energie des Heliums im Grundzustande, sowie des tiefsten Terms von Ortho-Helium. *Z. Phys.* **1929**, *54*, 347–366. [CrossRef]
- 95. Hylleraas, E.A. Über den Grundterm der Zweielektronenprobleme von H⁻, He, Li⁺, Be⁺⁺ usw. Z. Phys. 1930, 65, 209–225. [CrossRef]
- Löwdin, P.O. Correlation Problem in Many-Electron Quantum Mechanics I. Review of Different Approaches and Discussion of Some Current Ideas. In *Advances in Chemical Physics*; Prigogine, I., Ed.; John Wiley & Sons, Inc.: Hoboken, NJ, USA, 1958; Volume 2, Chapter 7, pp. 207–322. [CrossRef]
- Yoshizumi, H. Correlation Problem in Many-Electron Quantum Mechanics. II. Bibliographical Survey of the Historical Development with Comments. In *Advances in Chemical Physics*; Prigogine, I., Ed.; John Wiley & Sons, Inc.: Hoboken, NJ, USA, 1958; Volume 2, Chapter 8, pp. 323–366. [CrossRef]
- 98. Parr, R.G.; Yang, W. Density-Functional Theory of Atoms and Molecules; Oxford University Press: Oxford, UK, 1994.
- 99. Parr, R.G.; Bartolotti, L.J. Some remarks on the density functional theory of few-electron systems. *J. Phys. Chem.* **1983**, *87*, 2810–2815. [CrossRef]
- 100. De Proft, F.; Geerlings, P. Contribution of the shape factor $\sigma(r)$ to atomic and molecular electronegativities. *J. Phys. Chem. A* **1997**, *101*, 5344–5346. [CrossRef]
- 101. Ayers, P.W.; De Proft, F.; Geerlings, P. Comparison of the utility of the shape function and electron density for predicting periodic properties: Atomic ionization potentials. *Phys. Rev. A* 2007, 75, 012508. [CrossRef]
- Ayers, P.W. Density per particle as a descriptor of Coulombic systems. *Proc. Natl. Acad. Sci. USA* 2000, 97, 1959–1964. [CrossRef] [PubMed]
- 103. De Proft, F.; Ayers, P.W.; Sen, K.D.; Geerlings, P. On the importance of the "density per particle" (shape function) in the density functional theory. *J. Chem. Phys.* **2004**, *120*, 9969–9973. [CrossRef] [PubMed]
- Calais, J.L.; Löwdin, P.O. A simple method of treating atomic integrals containing functions of r₁₂. J. Mol. Spectrosc. 1962, 8, 203–211. [CrossRef]
- 105. Drake, G. High Precision Calculations for Helium. In *Springer Handbook of Atomic, Molecular, and Optical Physics*; Drake, G., Ed.; Springer: New York, NY, USA, 2006; Chapter 11, pp. 199–219. [CrossRef]
- Lin, C.H.; Ho, Y.K. Quantification of Entanglement Entropy in Helium by the Schmidt-Slater Decomposition Method. *Few-Body Syst.* 2014, 55, 1141–1149. [CrossRef]
- 107. Katriel, J.; Montgomery, H.E.; Sen, K.D. Hund's rule in the $(1s2s)^{1,3}S$ states of the two-electron Debye atom. *Phys. Plasmas* **2018**, *25*, 092111. [CrossRef]
- 108. Katriel, J.; Montgomery, H.E. Atomic vs. quantum dot open shell spectra. *J. Chem. Phys.* **2017**, 146, 064104. [CrossRef]
- 109. Sarsa, A.; Buendía, E.; Gálvez, F.; Katriel, J. Singlet vs. triplet interelectronic repulsion in confined atoms. *Chem. Phys. Lett.* **2018**, *702*, 106–110. [CrossRef]
- 110. Angulo, J.C.; Dehesa, J.S. Tight rigorous bounds to atomic information entropies. *J. Chem. Phys.* **1992**, 97, 6485–6495. [CrossRef]
- Angulo, J.C.; Dehesa, J.S. Erratum: Tight rigorous bounds to atomic information entropies [J. Chem. Phys. 97, 6485 (1992)]. J. Chem. Phys. 1993, 98, 9223–9223. [CrossRef]
- 112. Tao, J.; Li, G.; Li, J. Bounds to information entropies for atomic systems. J. Chem. Phys. **1997**, 107, 1227–1231. [CrossRef]
- 113. Tao, J.; Li, G.; Li, J. Rigorous bounds to information entropies for atomic systems. *Phys. Scr.* **1997**, *56*, 284–288. [CrossRef]
- 114. Sánchez-Moreno, P.; Zozor, S.; Dehesa, J.S. Upper bounds on Shannon and Rényi entropies for central potentials. *J. Math. Phys.* **2011**, *52*, 022105. [CrossRef]
- 115. Sánchez-Moreno, P.; Zozor, S.; Dehesa, J.S. Rigorous bounds for Rényi entropies of spherically symmetric potentials. *AIP Conf. Proc.* **2011**, *1305*, 192–199. [CrossRef]
- 116. Hornyák, I.; Nagy, Á. Inequalities for phase-space Rényi entropies. Int. J. Quantum Chem. 2012, 112, 1285–1290. [CrossRef]

- 117. Nahum, A.; Ruhman, J.; Huse, D.A. Dynamics of entanglement and transport in one-dimensional systems with quenched randomness. *Phys. Rev. B* 2018, *98*, 035118. [CrossRef]
- 118. Barghathi, H.; Herdman, C.; Maestro, A.D. Rényi Generalization of the Accessible Entanglement Entropy. *Phys. Rev. Lett.* **2018**, *121*, 150501. [CrossRef]
- Herdman, C.M.; Roy, P.N.; Melko, R.G.; Maestro, A.D. Entanglement area law in superfluid ⁴He. *Nat. Phys.* 2017, 13, 556–558. [CrossRef]
- Kaufman, A.M.; Tai, M.E.; Lukin, A.; Rispoli, M.; Schittko, R.; Preiss, P.M.; Greiner, M. Quantum thermalization through entanglement in an isolated many-body system. *Science* 2016, 353, 794–800. [CrossRef]
- 121. Islam, R.; Ma, R.; Preiss, P.M.; Tai, M.E.; Lukin, A.; Rispoli, M.; Greiner, M. Measuring entanglement entropy in a quantum many-body system. *Nature* **2015**, *528*, 77–83. [CrossRef]
- 122. McMinis, J.; Tubman, N.M. Renyi entropy of the interacting Fermi liquid. *Phys. Rev. B* 2013, *87*, 081108(R). [CrossRef]
- 123. Lévay, P.; Nagy, S.; Pipek, J. Elementary formula for entanglement entropies of fermionic systems. *Phys. Rev. A* 2005, 72, 022302. [CrossRef]
- 124. Rouse, I.; Willitsch, S. Superstatistical Energy Distributions of an Ion in an Ultracold Buffer Gas. *Phys. Rev. Lett.* **2017**, *118*, 143401. [CrossRef] [PubMed]
- 125. Dechant, A.; Kessler, D.A.; Barkai, E. Deviations from Boltzmann-Gibbs Statistics in Confined Optical Lattices. *Phys. Rev. Lett.* **2015**, *115*, 173006. [CrossRef] [PubMed]
- 126. Douglas, P.; Bergamini, S.; Renzoni, F. Tunable Tsallis Distributions in Dissipative Optical Lattices. *Phys. Rev. Lett.* **2006**, *96*, 110601. [CrossRef] [PubMed]
- Lutz, E. Anomalous diffusion and Tsallis statistics in an optical lattice. *Phys. Rev. A* 2003, 67, 051402(R).
 [CrossRef]
- 128. Romera, E.; Dehesa, J.S. The Fisher-Shannon information plane, an electron correlation tool. *J. Chem. Phys.* **2004**, *120*, 8906–8912. [CrossRef] [PubMed]
- Sagar, R.P.; Guevara, N.L. Mutual information and correlation measures in atomic systems. *J. Chem. Phys.* 2005, 123, 044108. [CrossRef] [PubMed]
- 130. Grassi, A. A relationship between atomic correlation energy and Tsallis entropy. *Int. J. Quantum Chem.* 2008, 108, 774–778. [CrossRef]
- 131. Delle Site, L. On the scaling properties of the correlation term of the electron kinetic functional and its relation to the Shannon measure. *Europhys. Lett.* **2009**, *86*, 40004. [CrossRef]
- 132. Delle Site, L. Erratum: On the scaling properties of the correlation term of the electron kinetic functional and its relation to the Shannon measure. *Europhys. Lett.* **2009**, *88*, 19901. [CrossRef]
- 133. Grassi, A. A relationship between atomic correlation energy of neutral atoms and generalized entropy. *Int. J. Quantum Chem.* **2011**, *111*, 2390–2397. [CrossRef]
- 134. Delle Site, L. Shannon entropy and many-electron correlations: Theoretical concepts, numerical results, and Collins conjecture. *Int. J. Quantum Chem.* **2014**, *115*, 1396–1404. [CrossRef]
- 135. Flores-Gallegos, N. Informational energy as a measure of electron correlation. *Chem. Phys. Lett.* **2016**, 666, 62–67. [CrossRef]
- 136. Alipour, M.; Badooei, Z. Information theoretic approach provides a reliable description for kinetic component of correlation energy density functional. *Int. J. Quantum Chem.* **2018**, *118*, e25791. [CrossRef]
- 137. Collins, D.M. Entropy Maximizations on Electron Density. Z. Naturforsch. A 1993, 48, 68–74. [CrossRef]
- 138. López-Rosa, S.; Esquivel, R.O.; Plastino, A.R.; Dehesa, J.S. Quantum entanglement of helium-like systems with varying-Z: compact state-of-the-art CI wave functions. *J. Phys. B* **2015**, *48*, 175002. [CrossRef]
- 139. Esquivel, R.O.; López-Rosa, S.; Dehesa, J.S. Correlation energy as a measure of non-locality: Quantum entanglement of helium-like systems. *Europhys. Lett.* **2015**, *111*, 40009. [CrossRef]



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).