

SUPPLEMENTARY MATERIAL FOR

Polyketide-derived secondary metabolites from a Dothideomycetes fungus, *Pseudopalawania siamense* gen. et sp. nov., (Muyocoprionales) with antimicrobial and cytotoxic activities

**Ausana Mapook,¹ Allan Patrick G. Macabeo,^{2,3} Benjarong Thongbai,²
Kevin D. Hyde,¹ and Marc Stadler^{2,*}**

¹Center of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai, 57100 Thailand

²Department of Microbial Drugs, Helmholtz Centre for Infection Research and German Centre for Infection Research (DZIF), partner site Hannover/Braunschweig, Inhoffenstrasse 7, 38124, Braunschweig, Germany

³Laboratory for Organic Reactivity, Discovery and Synthesis (LORDS), Research Center for the Natural and Applied Sciences, University of Santo Tomas, 1015 Manila, Philippines

¹These authors contributed equally on this work.

LIST OF SUPPORTING INFORMATION

	Page
S1. ^1H NMR spectrum (CDCl_3 , 700 MHz) of pseudopalawanone (1)	3
S2. ^{13}C NMR spectrum (CDCl_3 , 175 MHz) of pseudopalawanone (1)	4
S3. HSQC-DEPT spectrum of pseudopalawanone A (1)	5
S4. COSY spectrum of pseudopalawanone (1)	6
S5. HMBC spectrum of pseudopalawanone (1)	7
S6. ROESY spectrum of pseudopalawanone (1)	8
S7. NOESY spectrum of pseudopalawanone (1)	9
S8. LC-HR-ESIMS spectrum of pseudopalawanone (1)	10
S9. CD spectrum (MeOH) of pseudopalawanone (1)	10




Figure S1. ¹H NMR spectrum (CDCl₃, 700 MHz) of pseudopalawanone A (1).




Figure S3. HSQC spectrum of pseudopalawanone A (**1**).




Figure S4. COSY spectrum of pseudopalawanone A (**1**).




Figure S5. HMBC spectrum of pseudopalawanone A (1).




Figure S6. ROESY spectrum of pseudopalawanone A (1).






Figure S7. NOESY spectrum of pseudopalawanone A (**1**).



S8. HR-ESIMS spectrum of pseudopalawanone (**1**).



S9. CD spectrum (MeOH) of pseudopalawanone (**1**).