

Supplementary materials

Anti-inflammatory functions of alverine via targeting Src in the NF-κB pathway

Chae Young Lee¹, Han Gyung Kim¹, Sang Hee Park², Seok Gu Jang³, Kyung Ja Park³, Dong Sam Kim^{3,*}, Ji Hye Kim^{1,*} and Jae Youl Cho^{1,2,*}

¹ Department of Integrative Biotechnology, Biomedical Institute for Convergence at SKKU, Sungkyunkwan University, Suwon 16419, Republic of Korea; chaeyoung2@skku.edu (C.Y.L.), hanks523@skku.edu (H.G.K.), kjhmlkjhml@hanmail.net (J.H.K.), and jaecho@skku.edu (J.Y.C.)

² Department of Biocosmetics, Sungkyunkwan University, Suwon 16419, Republic of Korea; 84701@naver.com (S.H.P.)

³ Samcheok Prasiola Japonica Research Center, Samcheok City Hall, Samcheok 25914, Republic of Korea; prasiolra@korea.kr (D.S.K.); jangsg69@korea.kr (S.J.); kyu5132@korea.kr (K.J.P.)

* Correspondence: jaecho@skku.edu (J.Y.C.); +82-31-290-7868 (J.Y.C.), kjhmlkjhml@hanmail.net (J.H.K.); +82-82-31-290-7878 (J.H.K.), and prasiolra@korea.kr (D.S.K.); (+82) 33-570-3337 (D.S.K.)

Supplementary Figure S1. Anti-inflammatory effects of alverine in an *in vivo* acute gastritis model. Mice were pretreated with alverine (0–200 mg/kg) orally twice per day for 3 days then injected orally with EtOH/HCl (150 mM). (A) mRNA expression levels of iNOS, TNF-α and GAPDH in stomach tissues were examined using RT-PCR. (B) Total and phosphorylated protein levels of IkBα, p65, p50, iNOS, and β-actin were analyzed using a Western blot analysis.



