

Review

Immunoaffinity Capillary Electrophoresis in the Era of Proteoforms, Liquid Biopsy and Preventive Medicine: A Potential Impact in the Diagnosis and Monitoring of Disease Progression

Norberto A. Guzman 1,* and Daniel E. Guzman 1,2,†

- ¹ Princeton Biochemicals, Inc., Princeton, NJ 08816, USA
- ² Division of Hospital Medicine, Department of Medicine, University of California at San Francisco, San Francisco, CA 94143, USA; daniel.guzman@ucsf.edu
 - Correspondence: guzman@affinityce.com; Tel.: +1-908-510-5258
 - + Affiliation 2 is present address.

Citation: Guzman, N.A.; Guzman, D.E. Immunoaffinity Capillary Electrophoresis in the Era of Proteoforms, Liquid Biopsy and Preventive Medicine: A Potential Impact in the Diagnosis and Monitoring of Disease Progression. *Biomolecules* 2021, *11*, 1443. https://doi.org/10.3390/ biom11101443

Academic Editor: Alessandro Alaimo

Received: 24 August 2021 Accepted: 29 September 2021 Published: 1 October 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/). Abstract: Over the years, multiple biomarkers have been used to aid in disease screening, diagnosis, prognosis, and response to therapy. As of late, protein biomarkers are gaining strength in their role for early disease diagnosis and prognosis in part due to the advancements in identification and characterization of a distinct functional pool of proteins known as proteoforms. Proteoforms are defined as all of the different molecular forms of a protein derived from a single gene caused by genetic variations, alternative spliced RNA transcripts and post-translational modifications. Monitoring the structural changes of each proteoform of a particular protein is essential to elucidate the complex molecular mechanisms that guide the course of disease. Clinical proteomics therefore holds the potential to offer further insight into disease pathology, progression, and prevention. Nevertheless, more technologically advanced diagnostic methods are needed to improve the reliability and clinical applicability of proteomics in preventive medicine. In this manuscript, we review the use of immunoaffinity capillary electrophoresis (IACE) as an emerging powerful diagnostic tool to isolate, separate, detect and characterize proteoform biomarkers obtained from liquid biopsy. IACE is an affinity capture-separation technology capable of isolating, concentrating and analyzing a wide range of biomarkers present in biological fluids. Isolation and concentration of target analytes is accomplished through binding to one or more biorecognition affinity ligands immobilized to a solid support, while separation and analysis are achieved by high-resolution capillary electrophoresis (CE) coupled to one or more detectors. IACE has the potential to generate rapid results with significant accuracy, leading to reliability and reproducibility in diagnosing and monitoring disease. Additionally, IACE has the capability of monitoring the efficacy of therapeutic agents by quantifying companion and complementary protein biomarkers. With advancements in telemedicine and artificial intelligence, the implementation of proteoform biomarker detection and analysis may significantly improve our capacity to identify medical conditions early and intervene in ways that improve health outcomes for individuals and populations.

Keywords: proteoforms; isoforms; low abundance biomarkers; liquid biopsy; immunoaffinity capillary electrophoresis; molecular biorecognition; point-of-care instrument; proteomics; precision medicine; preventive medicine; cancer; telemedicine; exosomes; circulating tumor cells; circulating immunological cells

1. Introduction

Clinical medicine is currently practiced based on the evaluation of symptoms and signs of disease with diagnostic tests and the lack thereof with a handful of screening tests;



the results of these tests help to determine what illness a patient has. Ultimately, identifying a disease and determining its severity leads to targeted therapies and lifestyle interventions, with the hope that a patient's symptoms and signs of disease will improve [1– 4]. The implication of a clinical structure that primarily functions on disease presentation is that most of healthcare spending is focused on treating rather than preventing chronic disease [5,6]. Based on the Centers for Medicare and Medicaid Services (CMS), the United States spends approximately 75% of annual healthcare spending on treating chronic disease alone [5,6]. Despite this, chronic diseases, such as heart disease, stroke, cancer, chronic kidney disease, and diabetes continue to be the leading causes of death, disability, and reduced quality of life in the United States [7,8]. Such a focus on treatment of illness once it clinically manifests has brought into question how improvements in and implementation of preventive medicine could reduce the healthcare burden on patients and the healthcare system as a whole.

For decades, we have known that disease manifests before it is clinically apparent or, in other words, before it manifests with symptoms and signs recognizable to patients and their health providers [9–11]. Many chronic diseases are "silent" and may take several years to manifest [12–18]. In 1999, for example, a large prospective study on Type 2 Diabetes Mellitus published in the United Kingdom demonstrated that pancreatic beta-cell dysfunction starts almost a decade before patients are diagnosed with Type 2 Diabetes Mellitus [19,20]. The consequence of this based on current testing is that patients will have already lost up to 50% of their beta cell function by the time practitioners are able to diagnose them with this disease, making treatment and cure of the disease more difficult [19– 23]. To give another example, in a study of the evolutionary history of 2658 cancers, researchers determined that there are genetic mutations that occur far before the clinical manifestation of cancers that can be detected, if tested for [24]. For instance, the rate of cytosine guanine (CpG) to thymine guanine (TpG) mutations can be a surrogate to predict latency of presentation of disease, with ovarian adenocarcinoma presenting up to 30 years after the occurrence of these mutations [24]. It has therefore become apparent that a stronger emphasis on screening for disease and disease monitoring, coupled with early targeted therapeutic interventions can ultimately prevent the manifestation and burden of symptoms and signs of disease.

While diagnostic testing methods have advanced in many ways over the past several decades, detection of biomarkers prior to or in the early stages of disease manifestation remains limited [25,26]. A handful of screening tests have been approved for diseases such as colon cancer, breast cancer, prostate cancer, and lung cancer, however most of these tests utilize imaging modalities and direct visualization with only colon cancer and prostate cancer offering laboratory diagnostic detection. The majority of laboratory diagnostic testing is now carried out via immunoassays [27], which have proved reliable if the clinical suspicion for disease is moderate to high. Nevertheless, current immunoassays remain limited when applied to early detection of disease due to polyreactivity of their antibodies, the presence of interfering substances in a sample, and concentrations of biomarkers below levels of detection [27,28]. For example, use of existing immunoassays to detect prostate specific antigen (PSA) for prostate cancer screening is controversial due to varying accuracy of results (see Figure 1).



Figure 1. Depicts a schematic published from the USPSTF in 2018, detailing the use of PSA for prostate cancer screening. Figure modified from [29].

In summary, the above schematic demonstrates that for every 1000 men screened for prostate cancer, only 1 will avoid death from prostate cancer and 3 will avoid metastatic spread of the cancer, with the remaining men suffering psychological harm of a potential diagnosis of cancer, risks of prostate biopsy, and side effects of erectile dysfunction and urinary incontinence from treatment of prostate cancer [29]. While there is no clear evidence to recommend for or against prostate cancer screening, it is clear that the potential harm from screening may outweigh the potential benefit for many individuals due to false positives and a lack of understanding of the severity of different prostate cancers.

In theory, elimination of false-positive and false-negative results from immunoassays could result in greater reliance on laboratory testing, particularly in the use of screening and monitoring for a variety of diseases without the need for clinical interpretation [30,31]. Ultimately, such a change could revolutionize the way clinical medicine is practiced and create a stronger emphasis and investment in prevention of disease through accurate screening tests. The goal of information gathering in the diagnostic process is to reduce the diagnostic uncertainty enough to make optimal decisions for subsequent care [32]. Profiling of the genome and transcriptome are now established tools for the discovery of novel biomarkers, but alterations in proteome expression are more likely to reflect changes in disease pathophysiology [33]. We will therefore discuss advancements in the field of proteomics and how a useful technology known as immunoaffinity capillary electrophoresis can potentially overcome limitations in sensitivity, specificity, and predictive value seen in current immunoassays.

2. Proteoforms as Diagnostic Biomarkers

Proteins are the true machines of life and are considered to be the "workhorses" of the cell [34]. They provide functional context to cells, tissues, and organisms in the form of receptors, signaling cascades, channels, and transporters. Proteins are also major structural components of the cytoskeleton and extracellular matrices [35]. They are uniquely identified by their proteoforms, defined as different forms of a protein derived from a single gene including all forms of genetic variations (e.g., amino acid variation), alternative splicing, and post-translational modifications (PTM) [36–39]. This means that one transcribed gene can lead to a variety of protein structures, and that the biological function of each proteoform, as well as the cellular location, binding partners and kinetics can vary greatly [40] (Figure 2). As a consequence of the vast number of proteoforms that can be formed, the number of proteins in an individual's proteome far exceeds the number of protein-coding genes in the human genome [41,42]. The identification of more proteoforms associated with different diseases will undoubtedly be an essential landmark discovery necessary for early diagnosis, prevention and treatment of disease [43,44].



Figure 2. Schematic representation of (**A**) a single gene and three isoforms; the isoform variations combine with site-specific changes to generate proteoforms via post-translational modifications (PTMs). (**B**) Schematic depicting various types of isoforms, including individual examples of an isoform, a proteoform, and a multiproteoform complex. Figures modified from [39,43].

While proteomics, or the study of proteins, has become an increasingly important topic of study amongst researchers, proteins and enzymes biomarkers have been used as diagnostic tools for many years [45–47]. This is because disruptions of the highly orchestrated actions of proteins is the cause of practically all diseases. Examples of proteoforms that have already had a large clinical impact are apolipoprotein proteoforms, β -type natriuretic peptide (β NP), proteins found in disorders of glycosylation, detection of structural changes in transthyretin, hemoglobin proteoforms, cystatin C-truncated proteoforms, C-reactive protein, vitamin D-binding protein, transferrin, and immunoglobulin G (NISTmAb) (See Table 1) [48–101]. Ultimately, analysis of the human proteome, can give further insight into the modifications and dynamic intermolecular collisions that proteins undergo in disease states [102,103].

Proteoform	Function	Clinical Significance	Reference
Cystatin C	Endogenous cysteine protease inhibitor.	Assessment of progression of kidney function. Increased levels of Cystatin C can be found in acute and chronic kidney disease.	[48]
Chemokine RANTES	Key role in inflammation, cell recruitment, and T cell activation.	Associated with autoimmune diseases, arthritis, diabetes, obesity, cardiovascular diseases, breast and cervical cancers, and other diseases.	[49]
Amyloid proteoforms	Complication of normally soluble proteins that can join together to form an insoluble protein.	Associated with infiltrative proteinopathies, including Alzheimer's disease, restrictive cardiomyopathy, infiltrative kidney disease, and other diseases.	[50,51]
Amyloid-beta (Aβ) proteoforms	Amyloid-beta $(A\beta)$ plays a key role in the pathogenesis of Alzheimer's disease. Post- mortem identification of 26 proteoforms in the brain of Alzheimer's disease patients has provided significant information in patients suffering from Alzheimer's disease (AD) and dementia.	The heterogeneity of $A\beta$ proteoforms deepens our understanding of Alzheimer's disease and offers new avenues for investigation into pathological mechanisms of the disease, with implications for therapeutic development.	[52]
Lacritin proteoforms	Prevent tear film collapse and maintain epithelial homeostasis. Proteoforms include an active monomer, inactive polymers, and a splice variant termed lacritrin-c.	Associated with dry eye disease. Important in visual acuity and promotes basal tearing. Quantitation of the different proteoforms of tear lacritin may provide a diagnostic tool for ocular diseases.	[53,54]
Tumor suppressor PTEN	Control various aspects of cellular function, grouping them into three categories: intrinsic, function-induced, and inducible proteoforms.	PTEN proteoforms offer novel therapeutic opportunities in the treatment of various cancers and other diseases.	[55]
Cardiac troponin	Critical regulator of cardiac muscle contraction and relaxation.	Assessment of cardiac troponin proteoforms in serum of patients with acute myocardial	[56,57]

		infarction and other forms of	
		myocardial injury.	
MM-9 proteoforms	Matrix metalloproteinases (MMPs) are a class of secreted or cell bound endopeptidases, implicated in every step of the process of inflammatory cell migration.	Specific inhibition of MMPs has been suggested to be an interesting approach to control inflammation.	[58]
Apolipoprotein(a)	Lipoprotein(a) (Lp(a)) is an LDL-like particle, that contains a single copy of the apolipoprotein(a) covalently linked by a disulfide bridge to apolipoprotein(b). Its function is in wound healing, where it promotes tissue repair and vascular remodeling.	Lipoprotein(a) plays a role as an independent risk factor in the development of atherosclerotic cardiovascular diseases and calcified aortic valve disease. New research indicates that Lp(a) should be evaluated in terms of its apo(a) component and no longer in terms of Lp(a) mass.	[59]
Apolipoprotein A-1	Apolipoprotein A1 (Apo A-1) is the major constituent of human high-density lipoproteins, which plays a key role in reverse cholesterol transport and lipid homeostasis.	Apo A-1 exhibits antioxidant and anti-inflammatory properties and inhibits the aggregation and neurotoxicity of amyloid-beta peptide in Alzheimer's disease. Apo A-1 may possibly provide protection against neurological disorders.	[60]
DJ-1 proteoforms	DJ-1 is a cancer associated protein that protects cells from oxidative stress, functioning as a deglycase enzyme. DJ-1 acts as a redox-sensitive chaperone and as an oxidative stress sensor.	Associated with breast cancer. The modulation of specific DJ-1 function might produce substantial anticancer effects.	[61]
Prostate specific antigen proteoforms	Prostate specific antigen (PSA) is a glycoprotein with protease activity. This androgen- regulated serine-protease participates in the dissolution of the seminal fluid coagulum and plays an important role in fertility. PSA is produced by both prostate epithelial cells and prostate cancer cells.	Commonly used as serum biomarker for prostate cancer. Analysis of PSA proteoforms in urine and assessment of intact protein and glycopeptide analysis might be useful in improving prostate cancer screening.	[62]
Haptoglobin proteoforms	Serum haptoglobin (Hp) is a glycoprotein that scavenges freely circulating hemoglobin leaked into the blood stream when erythrocytes are damaged or die. Hp possesses four N- glycosylation sites on the beta- chain.	Studies have shown differences in the glycosylation pattern among patients with liver cirrhosis and hepatocellular carcinoma.	[63]
Erythropoietin	Erythropoietin (EPO) is a glycoprotein hormone of significant importance in the formation of red blood cells, as	EPO is used as a therapeutic protein for the treatment of anemia in chronic kidney disease and cancer.	[64–66]

	well as in other physiological		
	functions		
	Protoing identified in peripheral	Non invasivo difforential	
Kidney allograft	blood mononuclear cells	diagnostics of dysfunction of a	
	(PBMCs) as molecular	transplanted kidney, or	[67]
proteoforms	(FBWCS) as molecular	history of the hidrory of the	[07]
	signatures of kidney allograft	biomarkers of the kidney graft	
	pathology.	rejection.	
	EgAg B proteoforms are parasite	Biomarkers for early detection	
	antigen proteins present in	and monitoring of the	
	cystic echinococcocis disease	progression of the cystic	
EgAg B proteoforms	caused by the Echininococcus	echinococcosis disease. Specific	[68]
	granulosus metacestode	immunodominant epitopes	
	parasite. These antigens are	change as the disease	
	immunopotent.	progresses.	
		Monitoring of the proteoform	
		pattern changes in a growth	
	Human growth hormone (hGH)	hormone-secreting pituitary	
Human growth hormone	is synthesized by, stored in, and	adenoma, when compared to	
proteoforms	secreted by the pituitary gland.	control pituitary tissues, can be	[69]
proteoronnis	It promotes human growth and	of significant value for the	
	metabolism.	predictive diagnosis, targeted	
		prevention, and treatment of	
		pituitary adenoma.	
	The p53 family of proteins,		
	including p53, p63 and p73,	Δ Np73 proteoforms shows a	
	have a role in tumor	specific seroreactivity different	
	suppression. The $\Delta Np63$ and	from that of p73 with a higher	
	ΔNp73 proteoforms are	diagnostic ability to discriminate	
	frequently overexpressed in a	between colorectal cancer	
$\Delta Np73$ proteoforms	wide range of tumors, where	patients and controls, and	[70]
1 1	they are associated with poorer	especially premalignant	
	prognosis. Furthermore, it has	individuals and controls which	
	been demonstrated that the	may have an important impact	
	presence of autoantibodies to	on cancer prevention to predict	
	p53, p63 and p73 proteins exists	premalignant tumors.	
	in the serum of cancer patients.	1 0	
	Surfactant protein B (SP-B) is a		
	protein vital for normal lung		
	function. Higher levels of		
	circulating SP-B have been	Circulating immature	
	found in heavy smokers.	proteoform of surfactant protein	
	Immature SP-B (proSP-B) flow	B (proSP-B) has been proposed	
Surfactant protein B immature	into the bloodstream where it	as the most reliable lung-specific	[71]
proteoform (proSP-B)	hinds high-density lipoprotein	marker for alveolar-capillary	[, 1]
	(HDL) modifying its function	membrane dysfunction and	
	Impairing the alveolar cell SP-B	overall clinical status of heart	
	metabolism is likely the trigger	failure.	
	of the smoke-induced pro-		
	atherosclerotic cascade		
	The nost-translational	Obtaining a comprehensive	
Sarcomeric proteoforms	modifications (PTMs) of	view of the changes in the	
	sarcomeric protoine are known	sarcomeric protoome is an	
	to be important modistors of	important first stop toward	[72]
	cordiac signaling and over	understanding the molecular	[/ 2]
	various offects on contractile	underpippings of UCM Entered	
		material and the second	
	runction.	proteomics studies covering a	

_

	Hypertrophic cardiomyopathy (HCM) is the most common inherited disease and a leading cause of sudden death in young adults. It is characterized by abnormal thickening of the myocardium, which imposes a mechanical burden on the heart. HCM is highly heterogeneous and has been linked to mutations in the genes that encode proteins of the sarcomere.	wide range of HCM phenotypes will hold promise to help define disease progression and prognosis based on the proteoform landscape.	
NISTmAB proteoform	The NISTmAb is a recombinant humanized monoclonal antibody Reference Material from the National Institute of Standards and Technology. It is a class representative IgG1 κ intended to serve as a pre- competitive platform for harmonization and technology development in the biopharmaceutical industry.	Complex biotherapeutics, in particular monoclonal antibodies, increasingly dominate the arena of new drugs submitted for regulatory approval. Quality control mechanisms and their accompanying analytics are still evolving to meet increasingly sophisticated needs of the biotherapeutics market. Deviations in quality control may be linked to pathological conditions, such as immunogenicity or toxicity.	[73]
Natriuretic peptides proteoforms	The natriuretic peptide family consists of three biologically active peptides: atrial natriuretic peptide (ANP), brain (or B-type) natriuretic peptide (BNP), and C-type natriuretic peptide (CNP). Among these, ANP and BNP are secreted by the heart and act as cardiac hormones. ProANP and B-ANP are minor forms in the circulation but increases in patients with heart failure.	The human BNP precursor proBNP is proteolytically processed to BNP1-32 and N- terminal proBNP (NT-proBNP) within ventricular myocytes. Uncleaved proBNP as well as mature BNP1-32 and NT-proBNP are secreted from the heart, and its secretion is increased in patients with heart failure. Furthermore, NT-proBNP is O- glycosylated in the plasma of patients with severe heart failure.	[74,75]
Transthyretin proteoforms	Transthyrein (TTR) is a serum and cerebrospinal fluid (CSF) protein with metalloprotease activity. It is one of the most abundant proteins in CSF. Despite being best known for transporting thyroxine (T4) and retinol through the blood–brain barrier (BBB), TTR has been suggested to play a role in a broad range of functions in the central nervous system.	Dysregulations of TTR levels characterize several neurological disorders. In patients with stroke, the detection of TTR has been described as a positive prognostic indicator of clinical outcomes. TTR proteoforms are changed in the CSF of patients affected with spinal muscular atrophy (SMA) type, after treatment with the antisense oligonucleotide nusinersen.	[76]
C-reactive protein proteoforms	C-reactive protein (CRP) is an acute inflammatory protein that	CRP isoforms have distinct biological properties, with nCRP	[77]

	increases up to 1,000-fold at sites	often having more anti-	
	of infection or inflammation.	inflammatory activities	
	CRP is produced as a	compared to mCRP. The nCRP	
	homopentameric protein,	isoform activates the classical	
	termed native CRP (nCRP),	complement pathway, induces	
	which can irreversibly dissociate	phagocytosis, and promotes	
	at sites of inflammation and	apoptosis On the other hand	
	infection into five monomers	mCRP promotes the chemotaxis	
	termed monomorie CPP	and recruitment of circulating	
	(mCDD) CDD is symthesized		
	(mCRP). CRP is synthesized	ieukocytes to areas or	
	primarily in liver nepatocytes	inflammation and can delay	
	but also by smooth muscle cells,	apoptosis. mCRP increases	
	macrophages, endothelial cells,	interleuin-8 and monocyte	
	lymphocytes, and adipocytes.	chemoattractant protein-1	
	Having been traditionally	production.	
	utilized as a marker of infection		
	and cardiovascular events, there		
	is now growing evidence that		
	CRP plays important roles in		
	inflammatory processes and		
	host responses to infection		
	including the complement		
	nathway apontosis		
	pharocytosis piric oxide (NO)		
	release and the production of		
	autokinos, portigularlu		
	cytokines, particularly		
	interleukin-6 and tumor necrosis		
	factor- α .		
	Human SHC-Transforming		
	protein 1 (Shc1) has been found		
	to be important in the regulation		
	of apoptosis and drug resistance		
	in mammalian cells. Shc1 is an		
	intracellular scaffold protein,	Evaluation of p66shc/Shc1 has	
	which is involved in	been reported to be a useful	
	downstream pathways of cell	addition to the regularly used	
	surface signaling receptors, such	biomarkers, such as HbA1c, to	
	as the insulin signaling	increase diagnostic sensitivity	
	pathway. One of the isoforms of	for the identification of	
SHC-Transforming protein 1	Shc1 is p66shc. a mitochondrial	prediabetes (T2DM).	
proteoforms	associated oxidative stress	Knowledge of the presence of	[78,79]
r	biomarker, that may be involved	oxidative stress and	
	in regulating the life span and	inflammatory processes	
	the effects of reactive avy gon	forming a complex pattern of	
	sporios (ROS) Péésha has har	discass progression is therefore	
	species (KOS). Poosne has been	uisease progression, is therefore	
	implicated in several metabolic a	an important step towards early,	
	pathways, being able to act as an	effective treatment.	
	adaptor protein as well.		
	Furthermore, p66shc is		
	significantly altered in patients		
	with elevated blood glucose		
	levels.		
	Mammalian prolyl 4-	P4H catalyzes selective proline-	
	hydroxylase (P4H) is a tetramer	containing peptides to hydroxy-	
Proiyi nydroxylase alpha	composed of two unique	proline-containing peptides. The	[80-88]
subunit proteoforms	subunits, alpha and beta. One	best-known role of	
	gene makes the beta subunit	hydroxyproline is in stabilizing	
	o she had be be a bub diffe		

	that functions independently as a protein disulfide isomerase, and the other genes make three alpha subunit isoforms.	the collagen triple helix. A prolyl hydroxylase domain protein acts on the hypoxia inducible factor alpha subunits, which plays a key role in sensing molecular oxygen. Prolyl hydroxylases are essential for breast cancer metastasis, and a prolyl hydroxylase inhibitor decreases tumorogenesis.	
	Beyond their ability to	decreases taillorogenesis.	
Immunoglobulin G (IgG) proteoforms	neutralize pathogens, antibodies can mediate an array of effector functions through their interactions with Fc-receptors, complement molecules, and mammalian lectin-like molecules. Immunoglobulin G (IgG) fragment antigen binding (Fab) region binds a specific antigen, while its fragment crystallizable (Fc) region binds different receptors on the surface of various immune cells, thereby dictating the type of immune response elicited by the antigen binding.	Analysis of Fc-specific IgG glycosylation is critical for population-level studies of how antibodies may vary in response to vaccination or infection, and across disease states ranging from autoimmunity to cancer in both clinical and animal studies. IgG glycans are an excellent biomarker of biological age. Sialylation of Fc glycan has been reported to have anti- inflammatory activity.	[89–91]
Histone proteoforms	Chromatin is the structural framework that packages DNA into chromosomes within the nucleus of a cell. Histones comprises the principal protein component of chromatin and are involved in the regulation of gene expression. This epigenetic configuration is achieved through complex patterns of post-translational modifications, the incorporation of histones variants, and through controlled histone proteolysis.	Histone post-translational modifications (PTMs) are one of the main mechanisms of epigenetic regulation. Dysregulation of histones PTMs leads to many diseases, such as cancer.	[92–95]
KRAS proteoforms	Mutations of the KRAS gene are found in human cancers with high frequency and result in the constitutive activation of its protein products.	Mutations affecting post- translational modifications leads to aberrant regulation of downstream pathways, promoting cell survival, proliferation, and tumorigenesis that drive cancer progression and negatively affect treatment outcomes.	[96]
Huntingtin proteoforms	Huntington's disease (HD) is a rare neurodegenerative disorder caused by the aberrant expression of mutant Huntingtin (HTT) protein containing an expanded polyglutamine tract. The	HTT is now known to have a wide variety of post- translational modifications, including phosphorylation, sumoylation, acetylation, ubiquitination, and protease cleavage. Correlating these HTT	[97]

	expression of this protein is highly enriched in the brain relative to other tissues with highest expression in neurons.	proteoforms with functions should be fruitful in identifying mechanism of pathology that can be targeted for intervention.	
Alpha-synuclein proteoforms	Cumulative evidence suggests that lysosomal dysfunction contributes to neurodegenerative diseases, especially if amyloid proteins are involved. Among these, alpha-synuclein that progressively accumulates and aggregates in Lewis bodies is undisputedly a main culprit in Parkinson disease.	Alpha-synuclein can possess diverse post-translational modifications, aggregate formations, and truncations, all of which contribute to a growing set of proteoforms. These interfere directly or indirectly with lysosome function, reducing their own degradation, and thereby accelerating the protein aggregation and disease process.	[98]
Carbonic anhydrases proteoforms	Metalloenzymes carbonic anhydrases intervene in the second and rate limiting step of the catalytic mechanism of carbon dioxide reversible hydration. Genetic deficiencies of several carbonic anhydrases have been reported to be associated with diseases such as osteopetrosis, cerebral calcifications, hyperammonemia, retinal problems, and hyperchlorhydrolysis.	The loss of function of carbonic anhydrases would be in principle be treatable with selective activators of these enzymes. Carbonic anhydrase proteoforms represent a crucial family of new targets for improving cognition, but also in therapeutic, such as phobias, obsessive compulsive disorder, generalized anxiety, and post- traumatic disorders, for which few therapies are available. Studies on carbonic anhydrase proteoforms are important to elucidate the role of these modified proteins and their potential activators in brain processes.	[99]

Posttranslational modifications (PTMs) of proteins can happen at any step of the protein lifespan in both physiological and pathophysiological states. However, aberrant states of PTMs are frequently implicated in diseases [102,104,105]. Some of the most common protein modifications are hydroxylation, phosphorylation, acetylation, methylation, glycosylation, deamidation, prenylation, ubiquitylation, sumoylation, and proteolysis, among many others [104]. In particular, proteolysis generates protein isoforms or proteoforms with function and properties distinct from their precursors [105,106]. These proteolytic proteoforms regulate various biological processes and act as hormones, neurotransmitters, and disease pathogens. Unlike signal peptide or prodomain removal, however, protease-generated proteoforms can rarely be predicted from gene sequences [106]. Proteolytic processing is therefore a pervasive and irreversible post-translational modification that expands the protein universe by generating new proteoforms [106]. Deregulated proteolysis is a key driver for human diseases, and proteases are promising targets for tailored therapeutics [105,107].

Alterations in proteolytic systems underlie multiple pathological conditions [108]. For example, when viruses such as the SARS-CoV-2 virus infect a host, several proteolytic enzymes, either of the host or the virus, act in a concerted fashion to regulate and coordinate specific steps of viral replication and assembly [109]. To mitigate the high rate of infectivity of SARS-CoV-2, researchers have suggested that targeting host proteases involved in the entry of the viral genome can act as a potential means to block the ability of

the virus to infect host cells [109,110]. Clinicians have even suggested the use of a clinically approved protease inhibitor as a potential treatment option for those infected with SARS-CoV-2 [111]. An advantage of targeting host proteases to interfere with enveloped-virus entry is that this mechanism is common to several viruses and therefore may be applied to the treatment and prevention of infection with future novel viruses [109].

In another example, matrix metalloproteinases, a class of metal-linked zinc-dependent proteases that cleave internal peptide bonds of proteins to degrade extracellular matrices, have been linked to the development of cardiovascular disease in obese individuals [112]. For example, MMP-2 activity has been noted to be increased in obese rats with spontaneous hypertensive heart failure [113]. By targeting MMP directly through pharmacological inhibition, researchers found improved left ventricular function and reduced cardiac remodeling, thus suggesting that MMP activity contributed to left ventricle dysfunction in these obese rats [113]. This correlation has been seen in obese woman and adolescents as well, thereby demonstrating that direct inhibition of matrix metalloproteinases may potentially be used in the treatment of heart failure with reduced left ventricular function in obese individuals [114,115].

While the identification and inhibition of many proteases and proteoforms have demonstrated to be clinically useful in hampering disease development and progression, detecting these proteins through existing laboratory methods can be extremely challenging.

3. Limitations of Existing Immunoassays used in Clinical Diagnostics

Immunoassays are one of the most widely used detection methods for detecting protein biomarkers in clinical diagnostics, biopharmaceutical analysis, environmental monitoring, and food testing [116]. For most part, enzyme-linked immunosorbent assay (ELISA), immunohistochemistry (IHC) and flow cytometer assays in particular are immunoassays routinely found in diagnostic laboratories. ELISA continues to be the gold standard for biomarker detection [117-123] because of its high precision, throughput, and sensitivity (In clinical diagnostic chemistry, a sensitive test refers to a test that will correctly identify almost all individuals who likely have a disease, and will rarely yield a false negative result. A specific test refers to a test that will almost always correctly rule out those who do not have a disease, and will rarely yield a false positive result [117]. In analytical chemistry, sensitivity is the minimal detectable concentration of an analyte, expressed as the limit of detection (LoD), that can be accurately measured.). While the last two decades has seen tremendous innovation in ELISA technology, including automation with robotic workstations, improved microtiter plate readers, higher throughput well formats, and better antibody immobilization techniques, it has not been able to overcome for several reasons [119–123] limitations of diagnostic sensitivity[117] and specificity (Specificity is the ability to accurately assess a single analyte in the presence of components which may be expected to be present in the sample matrix, such as interferences. Similar to specificity, selectivity is the ability to accurately separate out and assess multiple components present in a sample mixture or matrix [119]).

First, ELISA and other immunoassays are prone to generate false positive results due to the polyreactivity of antibodies and antibody-like molecules. In theory, ELISA uses monoclonal antibodies that should provide higher specificity than polyclonal antibodies because they bind to a single epitope and usually have high affinity. In practice, there are several cases in which monoclonal antibodies have polyreactivity (Figure 3) [28,124–128]. One of the most prominent causes of antibody cross-reactivity or multi-specificity is molecular mimicry. Molecular mimicry is structural, functional or immunological similarities shared between macromolecules found in infectious pathogens and in host tissues [129–132]. Therefore, a false positive can result from binding of a substance different from the one of interest due to structural similarity in its antigens.



Figure 3. Schematic diagram showing the conformation hypothesis explaining polyreactivity. The classic 'lock and key' model of antigen-antibody interaction proposes rigid structures of interacting components and has been used for years to explain the monoreactivity of antibodies. Recent studies suggest that the antigen-binding 'pocket' of many antibody molecules is more flexible than previously thought and thus accommodate different antigenic configurations. Figure 3 shows four different antigens interacting with different amino acid residues within the antigen-binding pocket of a single broadly polyreactive antibody molecule. Each interaction alters the folding or conformation of the antigen-binding pocket in a different way. Figure modified from [28].

In addition to the polyreactivity of antibodies that are coated to the surface of a well that leads to binding of incorrect analytes of interest, interfering substances are another reason for inaccuracy of results in ELISA [133,134]. An interfering substance is one that remains in a test sample, after the washing steps in ELISA, and is able to bind a detection antibody and can elicit an enzyme-substrate reaction, thus producing a false positive result. Interfering substances include both endogenous substances, such as heterophile antibodies and binding proteins, and exogenous substances, such as antibodies administered to a patient for immunotherapy [133–138]. An example of an interference substance causing a false-positive result is a study of 12 women who were diagnosed with postgestational choriocarcinoma on the basis of persistently positive human chorionic gonadotropin (hCG) test results in the absence of pregnancy [139]. In one of these cases, a young woman underwent a hysterectomy, chemotherapy, radiotherapy, and a partial pneumonectomy before the interference was discovered and her diagnosis was corrected [139,140].

Lastly, antibodies used in ELISA can have difficulty binding biomarkers of interest when a signature protein is in the presence of other endogenous proteins where the concentration in some cases is a billion (1×10^{12}) times higher [141]. Such a dilutional factor makes it extremely difficult for a binding interaction to occur, thus resulting in a false-

negative result. While this may seem trivial, crucial proteins with diagnostic value are found in the very low abundance range ($<10^{-6}$ g/L) in biological fluids [141–143].

In order to improve clinical utility in detecting and monitoring proteins, existing testing must therefore progress in ways that allow for enhanced sensitivity, to overcome antibody polyreactivity and interference substances, and robust quantification of biomarkers, to detect those found in low concentration levels in biological samples. One valuable strategy to do this is to employ capillary electrophoresis, a reliable separation technique, with affinity-capture.

4. Capillary Electrophoresis as a Useful Technology for the Separation of Structurally Related Small Molecular-Weight Substances and Biomolecules

Capillary electrophoresis (CE) is a family of electrokinetic separation methods performed in submillimiter capillaries (conventional capillary electrophoresis) and in micro and nanofluidic channels (microchip electrophoresis) filled with a conductive fluid of a certain pH value. The basic principle of capillary electrophoresis is the separation of ions based on their electrophoretic mobility (mass/charge ratio) with the use of an applied voltage. Capillary electrophoresis can separate a wide range of chemical and biochemical entities, including neutral substances, cellular and subcellular entities, and nanoparticles or nanovesicles using one or more of 9 different separation modes [144–151]. Because of this, CE is known for its high-power resolution, high sensitivity and rapidity, high peak capacity, high separation efficiency with reproducible quantification, the use of small sample volumes with low waste generation, on-column and off-column detection, automation, low cost of operation, use of miniaturized components, option of multiple detection systems, and a diverse range of applications [144–151]. Compared to many other laboratory techniques, capillary electrophoresis can discern minor changes in substances that are structurally very similar.

In general, capillary electrophoresis in conventional or microchip format offers practical as well as technical advantages over traditional electrophoresis and other chromatographic techniques. When using narrow capillary tubes, the lateral diffusion effects are reduced, and the temperature differences across the tubes are also reduced. Furthermore, using narrow capillaries helps to reduce band-broadening seen in the peaks generated in other techniques such as high-performance liquid chromatography (HPLC). As a result, separation efficiency in CE is very high [144–151]. As opposed to conventional electrophoresis, which is labor-intensive and can take several hours to perform, CE can be completely automated and only a few microliters of sample are needed for an analysis to be carried out in a few minutes [123,151]. Although some commercial instruments are quite expensive, there are now many possibilities to manufacture miniaturized laboratory-made instruments for less than \$5000 with inexpensive components readily available [123,152]. Reagents and consumables are in general also inexpensive since minute amounts of reagents, buffers and solvents are used than traditional techniques. When performing molecular genotyping, CE detected more alleles and provided higher discriminatory power in comparative studies with agarose gel electrophoresis [153]. Regarding sodium dodecyl sulfate-polyacrylamide gel electrophoresis, a classical technique used for more than four decades but is time-consuming and labor intensive, CE shows many advantages such as on-column detection, automated operation, great resolving power, and capability of accurate protein quantification and molecular weight determination [154]. Regarding the throughput of sample analysis, most commercial analytical separation instruments, including HPLC and CE, are equipped to operate with one column making the process a low-throughput system. A recent report [89] comparing HPLC to CE for the analysis of glycans, were able to demonstrate using a CE ABI 3500 instrument (Applied Biosystems-Thermo Fisher) a high throughput for CE analysis. While the HPLC took as long as an hour to run a single sample, the ABI instrument was able to analyze 24 samples simultaneously in the span of approximately 15 min, with the ability to load two 96 well plates simultaneously, thus enabling the processing of as many as one thousand samples in a day. In addition to its remarkable throughput, this approach demonstrates robust sensitivity, permitting full glycan analysis with small quantities of high-value samples, requiring as little as 1 ul of plasma [89]. If IACE were to be incorporated in sample preparation prior to sample introduction into the CE system, the entire analysis system could be automated and faster.

Typical examples of CE electropherograms for small molecular weight substances and a biomolecule are presented in Figure 4. The electropherogram, or plot of detector signal over time, depicted in Figure 4A shows the simultaneous separation of thalidomide and its hydroxylated metabolites by chiral capillary chromatography (CCC), a mode of capillary electrophoresis separation for the separation of enantiomers [155–158]. The prerequisite of separation of enantiomers in CE is their enantioselective interaction with a chiral selector. Cyclodextrins have the shape of a torus with a hydrophobic interior and a hydrophilic outside. The hydrophilic external shell allows dissolution in the background electrolyte, whereas the lipophilic cavity is responsible for host–guest type interactions leading to the formation of inclusion complexes. In essence, each enantiomer will have a different binding (affinity) constant to the chiral selector, resulting in different effective mobilities and the ability to separate structurally related compounds [159]. In the example presented in Figure 4, in order to achieve optimal separation, the background electrolyte or running buffer contains two cyclodextrins as the chiral selector.

Figure 4B shows an electropherogram of a therapeutic monoclonal antibody. The electropherogram depicted in Figure 4B shows the different charge heterogeneity of the monoclonal antibody separated by imaged capillary isoelectric focusing (iCIEF) [160–163], a system in which the entire capillary is monitored by the whole-column detection system. An advantage of this technique over conventional CIEF is that mobilization of the protein zones is not required; therefore, the various charged variants of an antibody sample can be simultaneously recorded by the whole-column detector without disturbing the separation resolution [164].

As shown in Figures 4-A and 4-B, the electropherograms provides only separation and quantification of each entity represented by a peak intensity or peak area. However, partial structural information of each peak can be achieved by using the proper known standards employing the same experimental conditions. Conversely, a more complete structural analysis is accomplished by coupling CE to a mass spectrometer, a method capable of measuring mass-to-charge ratios of ions using a single instrument (MS) or in tandem (MS/MS), thereby increasing the ability to analyze chemical samples.





pl

7.0

6.5

5.8 pl marker

6.0

5.5

Although the separation of molecules using the various modes of capillary electrophoresis described above can achieve high resolution of a wide range of analytes in a short analysis time, the limits of detection (LOD) are constrained by the small sample volumes

7.5

80

(nanoliters) introduced into the capillaries and the small dimensions (micrometers) of the capillaries. The shorten pathlength of the capillary hinders common optical detection methods, such as UV detection [151,165–168]. Usually, optimal electropherograms can be obtained when using one or more of the 9 separation modes of CE if the target analytes are introduced into the capillary at high concentrations. However, working with diluted samples or complex mixtures where the target analytes are found at subnanomolar concentrations poses a significant limitation for detection of these analytes via CE; overcoming the poor sensitivity of CE has remained to be a challenge for many investigators [123,169–180]. We will therefore discuss immunoaffinity capillary electrophoresis, a method of CE that prevail over many of these limitations. If IACE were to be incorporated in sample preparation prior to sample introduction into the capillary or microchip, the entire system could be automated, reduce analysis time, and overcome limitations in poor detection sensitivity. We will emphasize the advantages for the determination of proteoforms when compared with other methods described to enhance the analytical sensitivity of analytes, such as field-amplified sample stacking, large-volume sample stacking, pH-mediated stacking, isotachophoreris (ITP), stacking techniques in micellar electrokinetic chromatography (MEKC), soli-phase extraction, and tagging target analytes with various chromophores and detecting the derivatized substances with sensitive detectors [123,151,152,173–190]. In some instances, IACE can be used in combination with other methods to enhance analytical sensitivity even further.

5. Advantages of Immunoaffinity Capillary Electrophoresis for the Detection of Proteoforms

Immunoaffinity capillary electrophoresis (IACE) is one of the three methods under the classification of affinity capillary electrophoresis. In general, affinity capillary electrophoresis is a broad term referring to the separation by capillary electrophoresis of substances that participate in specific or non-specific noncovalent affinity interactions during electrophoresis. The interacting molecules can be found in solution or can be immobilized to a solid support [123,146,174–180,191–203]. Immunoaffinity capillary electrophoresis, is the method referred to when using immobilized affinity ligands or probes to capture one or more analytes found in simple or complex matrices. Once captured in a portion of the capillary, or in a channel in microchip electrophoresis, the target analyte is isolated by eliminating the remaining sample, then releasing it from the immobilized ligand using an appropriate eluting buffer or solution. As a result, the target analyte is purified from the matrix and concentrated, significantly enhancing its limits of detection. To avoid confusion when referring to affinity capillary electrophoresis, we have defined the terminology as free-solution affinity capillary electrophoresis (FSACE) to the two methods of affinity capillary electrophoresis whose interactions are carried out in solution, and immobilized affinity capillary electrophoresis or immunoaffinity capillary electrophoresis (IACE) when the ligand or probes are immobilized to a surface within a portion of the capillary or channel. However, IACE not only uses antibodies or antibody fragments as highly selective capture agents, but also other biorecognition affinity reagents can act as ligands or probes, such as lectins, aptamers, metal-organic and other substances that have affinity for other analytes [122,174–180,191–203].

In order to better represent the principle by which the two techniques, ELISA and IACE, provide qualitative and quantitative information, Figure 5 depicts an apple with 6 different slices, each having a different characteristic represented by color and size. ELISA would be able to generate information of the entire apple, and not for each individual slice. Conversely, IACE would be able to provide qualitative and quantitative data for each individual slice, represented in the figure by each modified isoform of an individual proteoform.



Figure 5. Depicts a schematic representing the main differences between the traditional mono-dimensional sandwich ELISA technique with the two-dimensional immuno-capture-separation IACE. The two-dimensional IACE technology can achieve the capture, separation, and partial characterization of each individual isoform.

IACE is therefore considered to be a disruptive technique capable of discerning minor changes in substances such as metabolites of a substance or drug, or modified proteins such as proteoforms. As such, the applications of IACE are numerous, including safety regulation of new pharmaceuticals, identification of specific pathophysiologic changes that occur in inflammatory and other pathologic states, and early detection of disease through biomarker isolation and quantification. Developing biomarkers that can provide an accurate diagnosis and can predict whether patients are likely to benefit from the correct treatment is a pressing objective in many areas of medicine. IACE can be an excellent problem-solving technique, providing a new approach to monitor molecular changes throughout the evolution of a disease and through an assessment of each post-translational modification. Every modified protein component of a proteoform can change as the disease progressed, in particular the glycosylation pattern [62,200,204–206].

6. Applications of Immunoaffinity Capillary Electrophoresis. The Importance of Detecting Molecular Modifications of Proteins during the Progression of a Disease

Recombinant DNA-derived protein therapeutic products are often interchangeably referred to as biopharmaceuticals, biotherapeutics, biologicals, or biologics. Biologics are an important class of medicines, which are administered to treat an array of different diseases such as cancers, autoimmune disorders, infectious diseases, and many others [207–211]. The demand for biologics has risen considerable over the past few decades, owing to increase in prevalence of chronic diseases and development of plasma-derived therapies. To keep up with the demand and meet FDA and other worldwide regulatory agencies and standards for safety, production of biologics in a commercial setting requires

careful control of the manufacturing processing [212,213]. This strict control necessitates a battery of sensitive analytical methodologies to both elucidate the physicochemical properties and to monitor the purity and stability of the products.

Biologics are far from ideal molecules to be used as drugs, given their potential to create toxicity and immunogenicity concerns [212–214]. They are often glycosylated, and their carbohydrate profiles can impact solubility, tertiary structural stability, pharmacokinetics, cellular interactions, and immunogenicity [215]. Therefore, the ability to detect subtle physicochemical differences between purified batches is therefore critical in achieving a controlled process, as well as a consistent, safe, and efficacious product [216–218]. This process of detecting such structural differences can be applied to a variety of pharmaceuticals that have different proteoforms, particular important glycoproteins that can affect their clinical potential and risk profiles.

Figure 6 depicts the analysis by CE of two proteoforms, heparan-N-sulfatase and erythropoietin. Heparan-N-sulfatase (HNS) is a highly glycosylated lysosomal enzyme involved in the degradation of heparan sulfate used to treat patients with mucopolysaccharidosis type IIIA or Sanfilippo syndrome type A, a rare autosomal recessive disease occurring in 1 of 10,000 live births [216]. A deficiency in HNS causes the genetic disorder. Figure 6A shows an electropherogram of heparan-N-sulfatase by capillary zone electrophoresis (CZE). At present, no treatment options is available for this neurodegenerative disorder, although the potential use of enzyme replacement therapy is currently being evaluated. Erythropoietin (EPO), also known as hematopoietin or hemopoietin, is a glycoprotein hormone produced by the kidney in response to cellular hypoxia, that promotes the formation of red blood cells by the bone marrow [219,220]. It is used to treat patients with anemia due to chronic kidney disease. Figure 6B shows an electropherogram of erythropoietin. The electropherogram represent the separation of the different isoforms of erythropoietin by capillary zone electrophoresis (CZE). Until 1990, EPO was considered to have a single biological purpose and action, the stimulation of red blood cell growth and differentiation. EPO additionally is now known to be a pleiotropic growth factor that exhibits an anti-apoptotic action on numerous cells and tissues, including malignant ones [221]. Erythropoietin is highly glycosylated. Endogenous and recombinant human EPO (rHuEPO) only differ in the composition of their N- and O-glycan moieties. The use of lectins has therefore been used to characterize subtle differences in amongst glycoproteoforms of EPO [222].



Figure 6. Depicts the electropherograms of 2 glycoproteins. (**A**) depicts an electropherogram of the various glycoforms of heparan-N-sulfatase separated by capillary zone electrophoresis (CZE). (**B**) depicts an electropherogram showing the various glycoforms of erythropoietin separated by CZE. Figures modified from [216,223].

Aside from biopharmaceuticals, significant glycosylation is seen in most serum proteins and is implicated in many disease states. However, the glycome describes the complete repertoire of glycoconjugates composed of carbohydrate chains, or glycans, that are covalently linked not only to protein molecules, but also lipids [224–226]. More recently, it was reported that small RNAs which are modified with N-glycans are displayed on the surface of living cells [227]. Many malignancies are characterized by abnormal glycosylation, which is commonly associated with the oncogenesis and cancer progression [224– 226,228–238]. Furthermore, persistent non-malignant inflammatory conditions cause continual upregulation of enzymes that promote numerous molecular changes, including glycosylation as well [228–237]. For example, differences in the PSA glycosylation pattern have been found between benign prostate alterations and prostate cancer, and between aggressive prostate cancer and indolent prostate cancer [236]. Additionally, differences in the proportions of PSA fucosylation, sialyation and GalNAc are found in aggressive stages of prostate cancer [236–238]. Although significant progress has been made in identifying changes in the glycosylation patterns of some proteoforms, obtaining inadequate amounts of PSA glycoforms for separation and detection has limited its clinical use in the screening and diagnosis of prostate cancer [236].

Figure 7 depicts comparative results of two examples of proteoforms, transferrin and prostate specific antigen, using ELISA, a mono-dimensional immunoassay, and IACE, a two-dimensional immunoassay. Figure 7A depicts the analysis of transferrin using the ELISA method, and Figure 7B depicts the analysis of transferrin using the IACE method. Transferrin (Tf) is an abundant serum metal-binding protein that delivers iron through blood circulation to cells [239-241]. Studies of congenital atransferrinemia in mice and humans highlight the essential role of transferrin in erythropoiesis and iron metabolism. Patients and mice deficient in transferrin exhibits anemia and a paradoxical iron overload attribute to deficiency in hepticidin, a peptide hormone synthesized largely by the liver that inhibits iron absorption and macrophage iron efflux [242]. Transferrin is also a biomarker of chronic alcohol use [243,244]. Figure 7C depicts the analysis of prostate specific antigen (PSA) using the ELISA method, and Figure 7D depicts the analysis of PSA using the IACE method. Prostate specific antigen is a glycoprotein and is expressed by both normal and neoplastic prostate tissue, making increases in PSA nonspecific [245]. However, isolation of different glycosylated forms of PSA can aid in improving the sensitivity and specificity for prostate cancer diagnosis.



Figure 7. Depicts comparative results of two isoforms using ELISA, a monodimensional immunoassay, and IACE, a twodimensional immunoassay. (**A**) depicts the analysis of transferrin using the ELISA method, and (**B**) depicts the analysis of transferrin using the IACE method. (**C**) depicts the analysis of prostate specific antigen (PSA) using the ELISA method, and (**D**) depicts the analysis of PSA using the IACE method. Figures modified from [246,247].

Improvements in the characterization of proteoforms, using an ultra-low background system that combines an advanced detector technology with signal processing algorithms, have provided various sialoform profiles of transferrin. For example, as depicted in Figure 8, two aliquots of transferrin subjected to sialic acid cleavage were separated by capillary zone electrophoresis [248]. In the first aliquot, transferrin was incubated for 1 h with neuraminidase, and the resulting electropherogram shows primarily 3 isoforms, P0 (asialo-Tf), P1 (monosialo-Tf), and P2 (disialo-Tf). The second aliquot, transferrin was incubated for 5 h with neuraminidase, and the resulting electropherogram shows primarily electropherogram shows 4 isoforms, but changes occur in migration times and quantification. The resulting electropherogram shows P2, P3 (trisialo-Tf), P4 (tetrasialo-Tf) and P5 (pentasialo-Tf). Variation in serum sialoform content can be correlated to different pathological states, such as carbohydrate-deficient transferrin (CDT) [249], congenital disorders of glycosylation (CDG) [250], and cancer [251]. These improvements having better data acquisition, in combination with IACE, might result in superior data reproducibility to monitor minor protein changes during the course of progression of a disease.



Figure 8. Depicts some changes occurring in an artificially created microenvironment where an intact transferrin proteoform can be altered by adding neurominidase and removing the terminal sialic acids attached to transferrin. The separation and quantification profile of the intact transferrin isoforms changes due to the removal of sialic acids which alter the mass/charge ratio of the molecules. This electropherogram was performed with an ultra-low background system that combines an advanced detector technology with signal processing algorithms. Figure modified from [248].

As depicted in the above figures, IACE has the capability of capturing, separating and detecting glycoproteins, including those present in minute quantities. The target proteoform is captured and bound to antibodies immobilized to a solid support within a portion of the capillary or microchip channel, or directly into the wall of the capillary or channel. If enough protein substance is bound to the antibody, it is then eluted and detected on-line usually by ultraviolet detection. Alternatively, if not enough protein substance is bound, the protein can be labeled in situ by a fluorescent chromophore, separated by one or more of the CE separation modes, and detected on-line using a laser-induced fluorescence. Ideally, labeling and preconcentrating of a protein sample prior to introducing the sample into the capillary, will enhance even further the detection of a protein found at low abundance in complex mixtures. Furthermore, the CE instrument can be coupled to a mass spectrometer providing additional protein information. Peptide mapping of the proteoform can also be carried out, as well as identification of the sugar moieties of the proteoform [123,151,174,197–203,246,252,253]. Greater implementation of IACE therefore has the potential to further understand protein biomarkers, their application to disease states and pharmaceuticals, and have the potential to improve health outcomes through prevention and early diagnosis of disease.

7. Future of Immunoaffinity Capillary Electrophoresis for the Analysis of Proteoforms

With the growing geriatric population and the increasing prevalence of chronic diseases across the globe, growing demand for improving diagnostic laboratory techniques will continue. Proteomics should ideally be the driver for improving diagnostics and will serve as a primary method for detection of molecular changes indicative of disease states. Numerous laboratories have reported a plethora of assays using capillary electrophoresis in conventional format and in microfluidic format for detection and quantification of several proteoforms [254–267]. However, in order for proteoforms to reach their full clinical potential, it is necessary to be able to accurately detect and quantify near all proteoforms in biological fluids using advanced and sophisticated analytical instrumentation, such as IACE.

Use of immunoaffinity capillary electrophoresis in conventional CE instruments or in a point-of-care instrument has a significant potential to profile proteoform patterns in biological fluids, and in proteins obtained from circulating tumor cells, circulating immunological cells, exosomes, and other cellular or vesicles present in biofluids [123,174,175,195–203,246,247]. The ideal proteoform biomarker would be cost-effective, objective, fast to process and easy to interpret, with high clinical diagnostic sensitivity and specificity. IACE can capture and separate one or more proteoforms simultaneously, with the capability to discern minor molecular changes at high analytical sensitivity and specificity, and the data can be obtained in a short period of time with higher accuracy than conventional immunoassays. Furthermore, IACE is a more sustainable and affordable technology, given that it minimizes its use of samples and reagents. Ideally, it can also be used in tandem with telemedicine, to further improve access to and cost of healthcare.

The future of the 21st century medicine will include the use of POCT instruments, allowing home-based monitoring of wellness and disease using small samples of biological fluids for liquid biopsy. This will incorporate conveniently obtained specimens such as oral fluid, sputum, urine, sweat and tears. In summary, IACE technology will shorten the clinical decision path, offering a quick and accurate means of detecting disease and selecting the best treatment for a particular illness without compromising safety or effectiveness.

Author Contributions: Both authors contributed to data analysis and manuscript writing. Both authors have read and agreed to the published version of the manuscript.

Funding: There was no external financial support to conduct the research project.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data is available, upon request due to privacy/ethical restrictions, directly from the corresponding author of the publication where the information was obtained. Alternatively, the data is openly available in a public repository that does not issue DOIs.

Acknowledgments: In Memoriam: We would like to dedicate this paper to the memory of a friend, colleague, pioneer and educator, Alejandro Venegas, whose life was unfortunately cut short, for his great contributions to molecular biology, biochemistry and vaccines in Chile.

Conflicts of interest: Norberto Guzman is the inventor of several patents issued and pending related to the IACE technology.

References

- Stone, J.; Carson, A.; Sharpe, M. Functional symptoms and signs in neurology: Assessment and diagnosis. J. Neurol. Neurosurg. Psychiatry 2005, 76 (Suppl. S1), i2–i12, https://doi.org/10.1136/jnnp.2004.061655.
- 2. Davis, M.; Votey, S. *Sign and Symptoms in Emergency Medicine 2006*, 2nd ed.; Mosby-Elsevier: Amsterdam, The Netherlands, 2006; Paperback 9780323036450.
- 3. King, L.S. Signs and symptoms. J. Am. Med. Assoc. 1968, 206, 1063–1065, https://doi.org/10.1001/jama.1968.03150050051011.
- Llewelyn, H.; Ang, H.A.; Lewis, K.; Al-Abdullah, A. Oxford Handbook of Clinical Diagnosis, 3rd ed.; Oxford University Press: Oxford, UK, 2014; Print ISBN-13: 9780199679867, https://doi.org/10.1093/med/9780199679867.001.0001.
- National Center for Chronic Disease Prevention and Health Promotion. *Health and Economic Costs of Chronic Diseases*; Centers for Disease Control and Prevention: Atlanta, GA, USA; US Department of Health and Human Services: Washington, DC, USA, 2018; Available online: https://www.cdc.gov/chronicdisease/about/costs/index.htm (accessed on 1 January 2021).
- 6. Partnership to Fight Chronic Disease. Available online: https://www.fightchronicdisease.org/sites/de-fault/files/docs/PFCD_ChronDisease_FactSheet3Final.pdf (accessed on 1 January 2020).
- Golics, C.J.; Basra, M.K.; Salek, M.S.; Finlay, A.Y. The impact of patients' chronic disease on family quality of life: An experience from 26 specialties. *Int. J. Gen. Med.* 2013, *6*, 787–798.
- 8. Raghupathi, W.; Raghupathi, V. An empirical study of chronic diseases in the United States: A visual analytics approach to public health. *Int. J. Environ. Res. Pub. Health* **2018**, *15*, 431, https://doi.org/10.3390/ijerph15030431.
- 9. Prince, A.E.R.; Berkman, B.E. When does an illness begin: Genetic discrimination and disease manifestation. *J. Law Med. Ethics* **2012**, 40, 655–664, https://doi.org/10.1111/j.1748-720X.2012.0069.x.
- 10. Hunter, P. The inflammation theory of disease. The growing realization that chronic inflammation is crucial in many diseases opens new avenues for treatment. *EMBO Rep.* **2012**, *13*, 968–970, https://doi.org/10.1038/embor.2012.142.
- Menozzi, E.; Mcnaughtan, J.; Schapira, A.H.V. The gut-brain axis and Parkinson disease: Clinical pathogenetic relevance. *Ann. Med.* 2021, 53, 611–624, https://doi.org/10.1080/07853890.1890330.
- 12. Cinque, A.; Vago, R.; Trevisani, F. Circulating RNA in kidney cancer: What we know and what we still suppose. *Genes* **2021**, *12*, 835, https://doi.org/10.3390/genes12060835.
- 13. Özgur, Y. Relationship between vitamin D deficiency, albuminuria, peripheral artery disease and 5-year mortality in chronic kidney disease. J. Coll. Physicians Surg. Pak. 2021, 30, 644–650, https://doi.org/10.29271/jcpsp.2021.06.64412.
- 14. Carmichael, J.; Fadavi, H.; Ishibashi, F.; Shore, A.C.; Tavakoli, M. Advances in screening, early diagnosis and accurate staging of diabetic neuropathy. *Front. Endocrinol.* **2021**, *12*, 671257, https://doi.org/10.3390/fendo.2021.671257. eCollection 2021.
- 15. Shaw, G. The silent disease. *Nature* **2016**, *537*, S98–S99, https://doi.org/10.1038/537S98a.
- Hanmer, J.; Yu, L.; Li, J.; Kavalieratos, D.; Peterson, L.; Hess, R. The diagnosis of asymptomatic disease is associated with fewer healthy days: A cross sectional analysis from the national health and nutrition examination survey. *Br. J. Health Psychol.* 2019, 24, 88–101, https://doi.org/10.1111/bjhp.12341.
- 17. Saquib, N.; Saquib, J.; Ioannidis, J.P.A. Does screening for disease save lives in asymptomatic adults? Systemic review of metaanalyses and randomized trials. *Int. J. Epidemiol.* 2015, 44, 264–277, https://doi.org/10.1093/ije/dyu140.
- 18. Cebey-López, M.; Salas, A. Recognising the asymptomatic enemy. *Lancet Infect. Dis.* 2021, 21, 305–306, https://doi.org/10.1016/S1473-3099(20)30587-9.
- 19. King, P.; Peacock, I.; Donnelly, R. The UK Prospective Diabetes Study (UKPDS): Clinical and therapeutic implications for type 2 diabetes. *Br. J. Clin. Pharmacol.* **1999**, *48*, 643–648, https://doi.org/10.1046/j.1365-2125.1999.00092.x.
- Russo, G.T.; Giorda, C.B.; Cercone, S.; Nicolucci, A.; Cucinotta, D.; on behalf of BetaDecline Group. Factors associated with betacell dysfunction in type 2 diabetes: The BetaDecline study. *PLoS ONE* 2014, 9, e109702, https://doi.org/10.1371/journal.pone.0109702.
- 21. Kahn, S.E. Clinical review 135: The importance of beta-cell failure in the development and progression of type 2 diabetes. *J. Clin. Endocr. Metabol.* **2001**, *86*, 4047–4058, https://doi.org/10.1210/jcem.86.9.7713.
- 22. White, M.G.; Shaw, J.A.M.; Taylor, R. Type 2 diabetes: The pathologic basis of reversible β-cell dysfunction. *Diabetes Care* **2016**, 39, 2080–2088, https://doi.org/10.2337/dc16-0619.
- 23. Holman, R.R. Assesing the potential for alpha-glucosidase inhibitors in prediabetic states. *Diab. Res. Clin. Pract.* **1998**, 40 (Suppl. S1), S21–S25, https://doi.org/10.1016/s0168-8227(98)00038-2.
- 24. Gerstung, M.; Jolly, C.; Leshchiner, I.; Dentro, S.C.; Gonzalez, S.; Rosebrock, D.; Mitchell, T.J.; Rubanova, Y.; Anur, P.; Yu, K.; et al. The evolutionary history of 2,658 cancers. *Nature* 2020, *578*, 122–128, https://doi.org/10.1038/s41586-019-1907-7.
- 25. Tang, Y.; Qiao, G.; Xu, E.; Xuan, Y.; Liao, M.; Ying, G. Biomarkers for early diagnosis, prognosis, prediction, and recurrence monitoring of non-small cell cancer. *OncoTargets Ther.* **2017**, *10*, 4527–4534, https://doi.org/10.2147/OTT.S142149.
- 26. Hassaneim, M.; Callison, J.C.; Callaway-Lane, C.; Aldrich, M.C.; Grogan, E.L.; Massion, P.P. The state of molecular biomarkers for the early detection of lung cancer. *Cancer Prev. Res.* **2012**, *5*, 992–1006, https://doi.org/10.1158/1940-6207.CAPR-11-0441.
- 27. Vashit, S.K.; Luong, J.H.T. (Eds.) Handbook of Immunoassay Technologies: Approaches, Performances, and Applications; Academic Press-Elsevier: San Diego, CA, USA, 2018; ISBN-13: 9780128117620.

- 28. Notkins, A.L. Polyreactivity of antibody molecules. Trends Immunol. 2004, 25, 174–179, https://doi.org/10.1016/j.it.2004.02.004.
- U.S. Preventive Services Task Force; Grossman, D.C.; Curry, S.J.; Owens, D.K.; Bibbins-Domingo, K.; Caughey, A.B.; Davidson, K.W.; Doubeni, C.A.; Ebell, M.; Epling, J.W.J.; et al. Screening for prostate cancer: US Preventive Task Force recommendation statement. J. Am. Med. Assoc. 2018, 319, 1901–1913, https://doi.org/10.1001/jama.2018.3710.
- Woloshin, S.; Patel, N.; Kesselheim, A.S. False negative test for SARS-CoV-2 infection–Challenges and implications. N. Engl. J. Med. 2020, 383, e38, https://doi.org/10.1056/NEJMp2015897.
- Lin, J.; Dai, W.; Li, W.; Xiao, L.; Luo, T.; Guo, Y.; Yang, Y.; Han, Y.; Zhu, P.; Wu, Q.; et al. Potential false-positive and false-negative results for COVID-19 IgG/IgM antibody testing after heat-inactivation. *Front. Med.* 2021, 7, 589080, https://doi.org/10.3389/fmed.2020.589080.
- 32. Kassirer, J.P. Our stubborn quest for diagnostic certainty. A cause of excessive testing. N. Engl. J. Med. 1989, 320, 1489–1491, https://doi.org/10.1056/NEJM198906013202211.
- Macklin, A.; Khan, S.; Kislinger, T. Recent advances in mass spectrometry based clinical proteomics: Applications to cancer research. *Clin. Proteom.* 2020, 17, 1–25, https://doi.org/10.1186/s12014-020-09283-w.
- Katathiya, U.; Padariya, M.; Faktor, J.; Coyaud, E.; Alfaro, J.A.; Fahraeus, R.; Hupp, T.R.; Goodlett, D.R. Interfaces with structure dynamics of the workhorses from cells revealed through cross-linking mass spectrometry (CLMS). *Biomolecules* 2021, *11*, 382, https://doi.org/10.3390/biom11030382.
- 35. Roehrl, M.H.; Roehrl, V.B.; Wang, J.Y. Proteome-based pathology: The next frontier in precision medicine. *Expert Rev. Precis. Med. Drug Dev.* **2021**, *6*, 1–4, https://doi.org/10.1080/23808993.2021.1854611.
- Smith, L.M.; Kelleher, N.L.; Consortium for Top Down Proteomics. Proteoform: A single term describing protein complexity. Nat. Methods 2013, 10, 186–187, https://doi.org/10.1038/nmeth.2369.
- Smith, L.M.; Kelleher, N.L. Proteoforms as the next proteomics currency. *Science* 2018, 359, 1106–1107, https://doi.org/10.1126/science.aat1884.
- 38. Smith, L.; Agar, J.; Chamot-Rooke, J.; Danis, P.; Ge, Y.; Loo, J.; Pasa-Tolic, L.; Tsybin, Y.; Kelleher, N. The human proteoform project: A plan to define the Human Proteome. *Preprints* **2020**, 2020100368, https://doi.org/10.20944/preprints202010.0368.v1.
- 39. Aebersold, R.; Agar, J.N.; Amster, I.J.; Bager, M.S.; Bertozzi, C.R.; Boja, E.S.; Costello, C.E.; Cravatt, B.J.; Fenselau, B.F.; Garcia, B.A.; et al. How many human proteoforms are there? *Nat. Chem. Bio.* **2018**, *14*, 206–214, https://doi.org/10.1038/nchembio.2576.
- 40. Hoff, F.W.; van Dijk, A.D.; Kornblau, S.M. Proteoforms in acute leukemia: Evaluation of age- and disease-specific proteoform patterns. *Intechopen* **2019**, *5*, 62–79; https://doi.org/10.5772/intechopen.90329.
- 41. Bogaert, A.; Fernandez, E.; Gevaert, K. N-terminal proteoforms in human disease. Trends Biochem. Sci. 2020, 45, 308–320.
- Ponomarenko, E.A.; Poverennaya, E.V.; Ilgisonis, E.V.; Pyatnitskiy, M.A.; Kopylov, A.T.; Zgoda, V.G.; Lisitsa, A.V.; Archakov, A.I. The size of the Human Proteome: The width and the depth. Intern. J. Anal. Chem. 2016, 2016, 7436849, https://doi.org/10.1155/2016/7436849.
- 43. Da Silva-Araújo, J.; Tavares-Machado, O.L. Proteoforms: General concepts and methodological process for identification. *Intechopen* **2019**, *2*, 523–555; https://doi.org/10.5772/intechopen.89914.
- 44. Gregorich, Z.R.; Ge, Y. Top-down proteomics in health and disease: Challenges and opportunities. *Proteomics* **2014**, *14*, 1195–1210; https://doi.org/10.1002/pmic.201300432.
- 45. Wu, A.H. Cardiac markers: From enzymes to proteins, diagnosis, laboratory to bedside. Ann. Clin. Lab. Sci. 1999, 29, 18–23.
- 46. Mayeux, R. Biomarkers: Potential uses and limitations. *NeuroRx* 2004, *1*, 182–188, https://doi.org/10.1602/neurorx.1.2.182.
- 47. Liang, S.L.; Chan, D.W. Enzymes and related proteins as cancer biomarkers: A proteomic approach. *Clin. Chim. Acta* 2007, 381, 93–97, https://doi.org/10.1016/j.cca.2007.02.017.
- Yassine, H.N.; Trenchevska, O.; Dong, Z.; Bashawri, Y.; Koska, J.; Reaven, P.D.; Nelson, R.W.; Nedelkov, D. The association of plasma cystatin C proteoforms with diabetic chronic kidney disease. *Proteome Sci.* 2016, 14, 7, https://doi.org/10.1186/s12953-016-0096-7.
- 49. Trenchevska, O.; Sherma, N.D.; Oran, P.E.; Reaven, P.D.; Nelson, R.W.; Nedelkov, D. Quantitative mass spectrometric immunoassay for the chemokine RANTES and its variants. *J. Proteom.* **2015**, *116*, 15–23, https://doi.org/10.106/j.jprot.2014.12.011.
- 50. Noor, A.; Zafar, S.; Zerr, I. Neurodegenerative proteinopathies in the proteoform spectrum–Tools and challenges. *Int. J. Mol. Sci.* **2021**, *22*, 1085; doi.10.3390/ijms22031085.
- 51. Muchtar, E.; Blauwet, L.A.; Gert, M.A. Restrictive cardiomyopathy. Genetics, pathogenesis, clinical manifestations, diagnosis, and therapy. *Circul. Res.* 2017, 121, 819–837, https://doi.org/10.1161/CIRCRESAHA.117.310982.
- 52. Wildburger, N.C.; Esparza, T.J.; LeDuc, R.D.; Fellers, R.T.; Thomas, P.M.; Cairns, N.J.; Kelleher, N.L.; Bateman, R.J.; Brody, D.L. Diversity of amyloid-beta proteoforms in the Alzheimer's disease brain. *Sci. Rep.* **2017**, *7*, 9520, https://doi.org/10.1038/s41598-017-10422-x.
- Georgiev, G.A.; Gh, M.S.; Romano, J.; Teixeira, K.L.D.; Struble, C.; Ryan, D.S.; Sia, R.K.; Kitt, J.P.; Harris, J.M.; Hsu, K.-L.; et al. Lacritin proteoforms prevent tear film collapse and maintain epithelial homeostasis. *J. Biol. Chem.* 2021, 296, 100070, https://doi.org/10.1074/jbc.RA120.015833.
- Justis, B.M.; Coburn, C.E.; Tyler, E.M.; Showalter, R.S.; Dissler, B.J.; Li, M.; McNamara, N.A.; Laurie, G.M.; McKown, R.L. Development of a quantitative immunoassay for tear lacritin proteoforms. *Trans. Vis. Sci. Tech.* 2020, *9*, 13, doi.10.1167/tvst.9.9.13.
 Malaney, P.; Uversky, V.; Davé, V. PTEN proteoforms in biology and disease. *Cell Mol. Life Sci.* 2017, *74*, 2783–2794.
- 56. Tiambeng, T.N.; Tucholski, T.; Wu, Z.; Zhu, Y.; Mitchell, S.D.; Roberts, D.S.; Jin, Y.; Ge, Y. Analysis of cardiac troponin proteoforms by top-down mass spectrometry. *Methods Enzymol.* **2019**, *626*, 347–374, https://doi.org/10.1016/bs.mie.2019.07.029.

- 57. Soetkamp, D.; Raedschelders, K.; Mastali, M.; Sobhani, K.; Bairey Merz, C.N.; Van Eyk, J. The continuing evolution of cardiac troponin I biomarker analysis: From protein to proteoform. *Expert. Rev. Proteom.* **2017**, *14*, 973–986, https://doi.org/10.1080/14789450.2017.1387054.
- Nuti, E.; Rossello, A.; Cuffaro, D.; Camodeca, C.; Van Bael, J.; van der Matt, D.; Martens, E.; Fiten, P.; Pereira, R.V.S.; Ugarte-Berzal, E.; et al. Bivalent inhibitor with selectivity for trimeric MMP-9 amplifies neutrophil chemotaxis and enables functional studies on MMP-9 proteoforms. *Cells* 2020, *9*, 1634, https://doi.org/10.3390/cells9071634.
- 59. Ruhaak, L.R.; Cobbaert, C.M. Quantifying apolipoprotein(a) in the era of proteoforms and precision medicine. *Clin. Chim. Acta* **2020**, *511*, 260–268, https://doi.org/10.1016/j.cca.2020.10.010.
- 60. Fania, C.; Arosio, B.; Capitanio, D.; Torretta, E.; Gussago, C.; Ferri, E.; Mari, D.; Gelfi, C. Protein signature in cerebrospinal fluid and serum of Alzheimer's disease patients: The case of apolipoprotein A-1 proteoforms. *PLoS ONE* **2017**, *12*, e0179280, https://doi.org/10.1371/journal.pone.0179280.
- Scumaci, D.; Olivo, E.; Fiumara, V.; La Chimia, M.; De Angelis, M.T.; Mauro, S.; Costa, G.; Ambrosio, F.A.; Alcaro, S.; Agosti, V.; et al. DJ-1 proteoforms in breast cancer cells: The escape of metabolic epigenetic misregulation. *Cells* 2020, *9*, 1968, https://doi.org/10.3390/cells9091968.
- 62. Moran, A.B.; Domínguez-Vega, E.; Nouta, J.; Pongracz, T.; de Reijke, T.M.; Wuhrer, M.; Lageveen-Kammeijer, G.S.M. Profiling the proteoforms of urinary prostate-specific antigen by capillary electrophoresis-mass spectrometry. *J. Proteom.* **2021**, *238*, 104148, https://doi.org/10.1016/j.jprot.2021.104148.
- 63. Tamara, S.; Franc, V.; Heck, A.J.R. A wealth of genotype-specific proteoforms fine-tunes hemoglobin scavenging by haptoglobin. *Proc. Natl. Acad. Sci. USA* **2020**, *117*, 15554–15564.
- Falck, D.; Haberger, M.; Plomp, R.; Hook, M.; Bulau, P.; Wuhrer, M.; Reusch, D. Affinity purification of erythropoietin from cell culture supernatant combined with MALDI-TOF-MS analysis of erythropoietin N-glycosylation. *Sci. Rep.* 2017, *7*, 5324, https://doi.org/10.1038/s41598-017-05641-1.
- Ng, T.; Marx, G.; Littlewood, T.; Macdougall, I. Recombinant erythropoietin in clinical practice. *Postgrad. Med. J.* 2003, 79, 367–376; https://doi.org/10.1136/pmj.79.933.367.
- Čaval, T.; Tian, W.; Yang, Z.; Clausen, H.; Heck, A.J.R. Direct quality control of glycoengineered erythropoietin variants. *Nat. Commun.* 2018, *9*, 3342, https://doi.org/10.1038/s41467-018-05536-3.
- Savaryn, J.P.; Toby, T.K.; Catherman, A.D.; Fellers, R.T.; LeDuc, R.D.; Thomas, P.M.; Friedewald, J.J.; Salomon, D.R.; Abecassis, M.M.; Kelleher, N.L. Comparative top down proteomics of peripheral blood mononuclear cells from kidney transplant recipients with normal kidney biopsies or acute rejection. *Proteomics* 2016, *16*, 2048–2058, https://doi.org/10.1002/pmic.201600008.
- Ahn, C.-S.; Han, X.; Bae, Y.-A.; Ma, X.; Kim, J.-T.; Cai, H.; Yang, H.-J.; Kang, I.; Wang, H.; Kong, Y. Alteration of immunoproteome profile of Echinococcus granulosus hydatid fluid with progression of cystic echinococcosis. *Parasites Vectors* 2015, *8*, 10; 10.1186/s13071-014-0610-7.
- 69. Li, B.; Wang, X.; Yang, C.; Wen, S.; Li, J.; Li, N.; Long, Y.; Mu, Y.; Liu, Q.; Li, X.; et al. Human growth hormone proteoform pattern changes in pituitary adenomas: Potential biomarkers for 3P medical approaches. *EPMA J.* **2021**, *12*, 67–89, https://doi.org/10.1007/s13167-021-00232-7.
- 70. Garranzo-Asensio, M.; Guzmán-Aránguez, A.; Povés, C.; Fernández-Aceñero, M.J.; Montero-Calle, A.; Ceron, M.A.; Fernandez-Diez, S.; Rodríguez, N.; de Cedrón, M.; Ramírez de Molina, A.; et al. The specific seroreactivity to ΔNp73 isoforms shows higher diagnostic ability in colorectal cancer patients than the canonical p73 protein. *Sci. Rep.* 2019, *9*, 13547, https://doi.org/10.1038/s41598-019-49960-z.
- 71. Banfi, C.; Brioschi, M.; Mapelli, M.; Gianazza, E.; Mallia, A.; Zoanni, B.; Salvioni, E.; Gugliando, P.; Capra, N.; Veglia, F.; et al. Immature circulating SP-B, bound to HDL, represents an early sign of smoke-induced pathophysiological alterations. *Biomolecules* 2021, *11*, 551, https://doi.org/10.3390/biom11040551.
- Tucholski, T.; Cai, W.; Gregorich, Z.R.; Bayne, E.F.; Mitchell, S.D.; Mcllwain, S.J.; de Lange, W.J.; Wrobbel, M.; Karp, H.; Hite, Z.; et al. Distinct hypertrophic cardiomyopathy genotypes result in convergent sarcomeric proteoform profiles revealed by topdown proteomics. *Proc. Natl. Acad. Sci. USA* 2020, *117*, 24691–24700. https://doi.org/10.1073/pnas.2006764117.
- 73. Turner, A.; Schiel, J.E. Qualification of NISTmAb charge heterogeneity control assays. *Anal. Bioanal. Chem.* 2018, 410, 2079–2093, https://doi.org/10.1007/s00216-017-0816-6.
- 74. Nakagawa, Y.; Nishikimi, T.; Kuwahara, K. Atrial and brain natriuretic peptides: Hormones secreted from the heart. *Peptides* **2019**, *111*, 18–25, https://doi.org/10.1016/j.peptides.2018.05.012.
- 75. Hammerer-Lercher, A.; Halfinger, B.; Sarg, B.; Mair, J.; Puschendorf, B.; Griesmacher, A.; Guzman, N.A.; Lindner, H.H. Analysis of circulating forms of proBNP and NT-proBNP in patients with severe heart failure. *Clin. Chem.* **2008**, *54*, 858–865, https://doi.org/10.1373/clinchem.2007.090266.
- Bianchi, L.; Sframeli, M.; Vantaggiato, L.; Vita, G.L.; Ciranni, A.; Polito, F.; Oteri, R.; Gitto, E.; Di Giuseppe, F.; Angelucci, S.; et al. Nursinersen modulates proteomics profiles of cerebrospinal fluid in spinal muscular atrophy type 1 patients. *Int. J. Mol. Sci.* 2021, 22, 4329, https://doi.org/10.3390/ijms22094329.
- 77. Sproston, N.R.; Ashworth, J.J. Role of C-reactive protein at sites of inflammation and infection. *Front. Immunol.* **2018**, *9*, 754, https://doi.org/10.3389/fimmu.2018.00754.
- 78. Mir, H.A.; Ali, R.; Mushtaq, U.; Khanday, F.A. Structure-functional implications of longevity protein p66Shc in health and disease. *Ageing Res. Rev.* **2020**, *63*, 101139, https://doi.org/10.1016/j.arr.2020.101139.

- 79. Jelinek, H.F.; Helf, C.; Khalaf, K. Human SHC-transforming protein 1 and its isoform p66shc: A novel marker for prediabetes. *J. Diabetes Investig.* **2021**, https://doi.org/10.1111/JDI.13551.
- Cutroneo, K.R.; Guzman, N.A.; Liebelt, A.G. Elevation of peptidylproline hydroxylase activity and collagen synthesis in spontaneous primary mammary cancers of inbred mice. *Cancer Res.* 1972, 32, 2828–2833.
- Gorres, K.L.; Raines, R.T. Prolyl 4-hydroxylase. Crit. Rev. Biochem. Mol. Biol. 2010, 45, 106–124, https://doi.org/10.3109/10409231003627991.
- Guzman, N.A.; Oronsky, A.L.; Desnick, R.J.; Cutroneo, K.R.; Prockop, D.J. A new isoenzyme of prolyl 4-hydroxylase from human placenta. Purification and partial characterization. In *Federation Proceedings*; Federation of American Societies for Experimental Biology: St. Louis, MO, USA, 1981; Volume 40, p. 1706.
- 83. Guzman, N.A.; Ascari, W.Q.; Cutroneo, K.R.; Desnick, R.J. Comparison between avian and human prolyl 4-hydroxylases: Studies on the holomeric enzymes and their constituent subunits. *J. Cell. Biochem.* **1992**, *48*, 172–189, https://doi.org/10.1002/jcb.240480209.
- 84. Guzman, N.A. Prolyl hydroxylase: An overview. In *Prolyl Hydroxylase, Protein Disulfide Isomerase, and Other Structurally Related Proteins*; Guzman, N.A., Ed.; Marcel Dekker Inc: New York, NY, USA, 1998; Chapter 1, pages 1–64.
- Gilkes, D.M.; Chaturvedi, P.; Bajpai, S.; Wong, C.C.-L.; Wei, H.; Pitcairn, S.; Hubbi, M.E.; Wirtz, D.; Semenza, G.L. Collagen prolyl hydroxylases are essential for breast cancer metastasis. *Cancer Res.* 2013, 73, 3285–3296, https://doi.org/10.1158/0008-5472.CAN-12-3963.
- 86. Hirota, K. HIF-a prolyl hydroxylase inhibitors and their implications for biomedicine: A comprehensive review. *Biomedicines* **2021**, *9*, 468, https://doi.org/10.3390/biomedicines9050468.
- Li, M.; Wang, Q.; Zheng, Q.; Wu, L.; Zhao, B.; Wu, Y. Prognostic and diagnostic roles of prolyl 4-hydroxylase subunit alpha members in breast cancer. *Biomark. Med.* 2021, 15, 1085–1095, https://doi.org/10.2217/bmm-2020-0323.
- Shi, R.; Gao, S.; Zhang, J.; Xu, J.; Graham, L.M.; Yang, X.; Li, C. Collagen prolyl-4-hydroxylases modify tumor progression. *Acta Biochim. Biophys. Sin.* 2021, 53, 805–814; doi:10.1093/abbs/gmab065.
- Mahan, A.E.; Tedesco, J.; Dionne, K.; Baruah, K.; Cheng, H.D.; De Jager, P.L.; Barouch, D.H.; Suscovich, T.; Ackerman, M.; Crispin, M.; et al. A method for high-throughput, sensitive analysis of IgG Fc and Fab glycosylation by capillary electrophoresis. J. Immunol. Meth. 2015, 417, 34–44, https://doi.org/10.1016/j.jim.2014.12.004.
- 90. Gudelj, I.; Lauc, G.; Pezer, M. Immunoglobulin G glycosylation in aging and diseases. *Cell. Immunol.* 2018, 333, 65–79; doi:10.1016/j.cellimm.2018.07.009.
- 91. Markina, Y.V.; Gerasimova, E.V.; Markin, A.M.; Glanz, V.Y.; Wu, W.-K.; Sobenin, I.A.; Orekhov, A.N. Sialylated immunoglobulins for the treatment of immune-inflammatory diseases. *Int. J. Mol. Sci.* 2020, *21*, 5472; doi.10.3390/ijms21155472.
- Anderson, L.C.; Karch, K.R.; Ugrin, S.A.; Coradin, M.; Engish, A.M.; Sidoli, S.; Shabanowitz, J.; Garcia, B.A.; Hunt, D.F. Analyses of histone proteoforms using front-end electron transfer dissociation-enabled orbitrap instruments. *Mol. Cell. Proteom.* 2016, 15, 975–988, https://doi.org/10.1074/mcp.O115.053843.
- 93. Lu, C.; Coradin, M.; Porter, E.G.; Garcia, B.A. Accelerating the field of epigenetic histone modification through mass spectrometry-based approaches. *Mole. Cell. Proteom.* **2021**, *20*, 10006, https://doi.org/10/1074/mcp.R120.002257.
- 94. Martire, S.; Banaszynski, L.A. The roles of histone variants in fine-tuning chromatin organization and function. *Nat. Rev. Mol. Cell. Biol.* **2020**, *21*, 522–541, https://doi.org/10.1038/s41580-020-0262-8.
- Lindner, H.H. Analysis of histones, histones variants, and their post-translational modified forms. *Electrophoresis* 2008, 29, 2516–2532, https://doi.org/10.1002/elps.200800094.
- 96. Ntai, I.; Fornelli, L.; DeHart, C.J.; Hutton, J.E.; Doubleday, P.F.; LeDuc, R.D.; van Nispen, A.J.; Fellers Whiteley, G.; Boja, E.S.; Rodriguez, H.; et al. Precise characterization of KRAS4b proteoforms in human colorectal cells and tumor reveals mutation/modification cross-talk. *Proc. Natl. Acad. Sci. USA* 2018, 115, 4140–4145, https://doi.org/10.1073/pnas.1716122115.
- 97. Seeley, C.; Kegel-Gleason, K.B. Taming the Huntington's disease proteome: What have we learned? *J. Huntingt. Dis.* **2021**, *10*, 239–257, https://doi.org/10.3233/JHD-200465.
- Wildburger, N.C.; Hartle, A.-S.; Schidlitzki, A.; Richter, F. Current evidence for a bidirectional loop between the lysosome and alpha-synuclein proteoforms. *Front. Cell. Dev. Biol.* 2020, *8*, 598446, https://doi.org/10.3389/fcell.2020.598446.
- Nocentini, A.; Cuffaro, D.; Ciccone, L.; Orlandini, E.; Nencetti, S.; Nuti, E.; Rosello, A.; Supuran, C.T. Activation of carbonic anhydrases from human brain by alcohol oxime ethers: Towards human carbonic anhydrase VII selective activators. *J. Enzyme Inhib. Med. Chem.* 2021, 36, 48–57, https://doi.org/10.1080/14756366.2020.1838501.
- Nedelkov, D. Mass spectrometric studies of apolipoprotein proteoforms and their role in lipid mebabolism and type 2 diabetes. *Proteomes* 2017, 5, 27, https://doi.org/10.3390/proteomes5040027.
- Lorentzian, A.; Uzozie, A.; Lange, P.F. Origins and clinical relevance of proteoforms in pediatric malignancies. *Expert Rev. Pro*teom. 2019, 16, 185–200, https://doi.org/10.1080/14789450.2019.1575206.
- Mnatsakanyan, R.; Shema, G.; Basik, M.; Batist, G.; Borchers, C.H.; Sickmann, A.; Zahedi, R.P. Detecting post-translational modification signatures as potential biomarkers in clinical mass spectrometry. *Expert Rev. Proteom.* 2018, 15, 515–535, https://doi.org/10.1080/14789450.2018.1483340.
- Toby, T.K.; Fornelly, L.; Kelleher, N.L. Progress in top-down proteomics and the analysis of proteoforms. *Annu. Rev. Anal. Chem.* 2016, 9, 499–519, https://doi.org/10.1146/annurev-anchem-071015-041550.
- Xu, H.; Wang, Y.; Lin, S.; Deng, W.; Peng, D.; Cui, Q.; Xue, Y. PTMD: A database of human disease-associated post-translational modifications. *Genom. Proteom. Bioinf.* 2018, 16, 244–251, https://doi.org/10.1016/j.gpb.2018.06.004.

- 105. Weng, S.S.H.; Demir, F.; Ergin, E.K.; Dirnberger, S.; Uzozie, A.; Tuscher, D.; Nierves, L.; Tsui, J.; Huesgen, P.F.; Lange, P.F. Sensitive determination of proteolytic proteoforms in limited microscale proteome samples. *Mol. Cell. Proteom.* 2019, *18*, 2335–2347, https://doi.org/10.1074/mcp.TIR119.001560.
- 106. Eckhard, U.; Marino, G.; Butler, G.S.; Overall, C.M. Positional proteomics in the era of the human proteome project on the doorstep of precision medicine. *Biochimie* **2016**, *122*, 110–118, https://doi.org/10.1016/j.biochi.2015.10.018.
- Vandenbroucke, R.E.; Libert, C. Is there new hope for therapeutic matrix metalloproteinase inhibition? *Nat. Rev. Drug Discov.* 2014, 13, 904–927, https://doi.org/10.1038/nrd4390.
- López-Otín, C.; Bond, J.S. Proteases: Multifunctional enzymes in life and disease. J. Biol. Chem. 2008, 283, 30433–30437, https://doi.org/10.1074/jbc.R800035200.
- 109. Gioia, M.; Ciaccio, C.; Calligari, P.; De Simone, G.; Sbardella, D.; Tundo, G.; Fasciglione, G.F.; Di Masi, A.; Di Pierro, D.; Bocedi, A.; et al. Role of proteolytic enzymes in the COVID-19 infection and promising therapeutic approaches. *Biochem. Pharmacol.* 2020, *182*, 114225, https://doi.org/10.1016/j.bcp.2020.114225.
- 110. Seth, S.; Batra, J.; Srinivasan, S. COVID-19: Targeting proteases in viral invasion and host immune response. *Front. Mol. Biosci.* **2020**, *7*, 215, https://doi.org/10.3389/fmolb.2020.0025.
- 111. Hoffmann, M.; Kleine-Weber, H.; Schroeder, S.; Krüger, N.; Herrler, T.; Erichsen, S.; Schiergens, T.S.; Herrler, G.; Wu, N.-H.; Nitsche, A.; et al. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell* 2020, 181, 271–280.e8, https://doi.org/10.1016/j.cell.2020.02.052.
- 112. Hua, Y.; Nair, S. Proteases in cardiometabolic diseases: Pathophysiology, molecular mechanisms and clinical applications. *Biochem. Biophys. Acta* 2015, 185, 195–208, https://doi.org/10.1016/j.bbadis.2014.04.032.
- 113. Peterson, J.T.; Hallak, H.; Johnson, L.; Li, H.; O'Brien, P.M.; Sliskovic, D.R.; Bocan, T.M.; Coker, M.L.; Etoh, T.; Spinale, F.G. Matrix metalloproteinase inhibition attenuates left ventricular remodeling and dysfunction in a rat model of progressive heart failure. *Circulation* 2001, 103, 2303–2309; doi:10.1161/01.cir.103.18.2303.
- 114. Miksztowicz, V.; Siseles, N.; Fernandez-Machulsky, N.; Schreier, L.; Berg, G. Increase in MMP-2 activity in overweight and obese women is associated with menopausal status. *Climateric* 2012, *15*, 602–606, https://doi.org/10.3109/13697137.2012.667174.
- 115. Glowinska-Olszewska, B.; Urban, M. Elevated matrix metalloproteinase 9 and tissue inhibitor of metalloproteinase 1 in obese children and adolescents. *Metabolism* **2007**, *56*, 799–805, https://doi.org/10.1016/j.metabol.2007.01.011.
- 116. Wild, D.G. (Ed.). The Immunoassay Handbook: In *Theory and Applications of Ligand Binding*, ELISA and Related Techniques, 4th ed.; Elsevier Science: Amsterdam, The Netherlands, 2013; ISBN: 9780080970370.
- 117. Saah, A.J.; Hover, D.R. "Sensitivity" and "specificity" reconsidered: The meaning of these terms in analytical and diagnostic settings. *Ann. Intern. Med.* **1997**, *126*, 91–94.
- 118. Li, Y.; Cassone, V.M. A simple, specific high-throughput enzyme-linked immunosorbent assay (ELISA) for quantitative determination of melatonin in cell culture medium. *Int. Immunopharmacol.* 2015, 28, 230–234, https://doi.org/10.1016/j.intimp.2015.06.012.
- Rao, T.N. Validation of Analytical Methods. In Calibration and Validation of Analytical Methods—A Sampling of Current Approaches, Chapter 7; IntechOpen Limited: London, UK, 2018, https://doi.org/10.5772/intechopen.72087.
- 120. Kwok, J.; Yang, M.; Wu, J.K.; Carter, C.J.; Jackson, S. High false positive rate of an ELISA screen for the detection of an antifactor VIII antibodies in congenital hemophilia A. *Blood* 2013, 122, 3586, https://doi.org/10.1182/blood.V122.21.3586.3586.
- 121. Wai, C.T.; Tambyah, P.A. False-positive HIV-1 ELISA in patients with hepatitis B. Am. J. Med. 2002, 112, 737, https://doi.org/10.1016/S0002-9343(02)01113-0.
- 122. Kharlamova, N.; Dunn, N.; Bedri, S.K.; Jerling, S.; Almgren, M.; Faustini, F.; Gunnarsson, I.; Rönnelid, J.; Pullerits, R.; Gjertsson, I.; et al. False positive results in SARS-CoV-2 serological tests for samples from patients with chronic inflammatory diseases. *Front. Immunol.* 2021, 12, 666114, https://doi.org/10.3389/fimmu.2021.666114.
- 123. Guzman, N.A.; Guzman, D.E. An emerging micro-scale immune-analytical diagnostic tool to see the unseen. Holding promise for precision medicine and P4 medicine. *J. Chromatogr. B* 2016, 1021, 14–29, https://doi.org/10.1016/j.jchromb.2015.11.026.
- 124. Jarossay, A.; Hadzhieva, M.; Kaveri, S.V.; Lacroix-Desmazes, S.; Dimitrov, J.D. Natural and induced antibody polyreactivity. *Anti-Cancer Agents Med. Chem.* 2015, 15, 1230–1241, https://doi.org/10.2174/1871520615666150716104935.
- 125. Hadzhieva, M.; Vassilev, T.; Bayry, J.; Kaveri, S.; Lacroix-Desmazes, S.; Dimitrov, J.D. Relationship between natural and hememediated antibody polyreactivity. *Biochem. Biophys. Res. Commun.* 2016, 472, 281–286, https://doi.org/10.1016/j.bbrc.2016.02.112.
- 126. Reyes-Ruiz, A.; Dimitrov, J.D. How can polyreactive antibodies conquer rapidly evolving viruses? *Trends Immunol.* **2021**, *42*, 654–657, https://doi.org/10.1016/j.it.2021.06.008.
- 127. Augustyniak, D.; Majkowska-Skrobek, G.; Roszkowiak, J.; Dorotkiewics-Jach, A. Defensive and offensive cross-reactive antibodies elicited by pathogens: The good, the bad and the ugly. *Curr. Med. Chem.* 2017, 24, 4002–4437, https://doi.org/10.2174/0929867324666170508110222.
- 128. Jain, D.; Salunke, D.M. Antibody specificity and promiscuity. Biochem. J. 2019, 476, 433–447, https://doi.org/10.1042/BCJ20180670.
- 129. Thaper, D.; Prabha, V. Molecular mimicry: An explanation for autoimmune diseases and infertility. *Scand. J. Immunol.* 2018, *88*, e12697, https://doi.org/10.1111/sji.12697.
- 130. Madhavaram, H.; Patel, T.; Kyle, C. Kavain interference with amphetamine immunoassay. J. Anal. Toxicol. 2020, bkaa178, https://doi.org/10.1093/jat/bkaa178 (accessed on 1 December 2020).
- Tapryal, S.; Gaur, V.; Kaur, K.J.; Salunke, D.M. Structural evaluation of a mimicry-recognizing paratope: Plasticity in antigenantibody interactions manifests in molecular mimicry. *J. Immunol.* 2013, 191, 456–463, https://doi.org/10.4049/jimmunol.1203260.

- 132. Obando-Pereda. Can molecular mimicry expain the cytokine storm of SARS-CoV-2?: An in silico approach. J. Med. Virol. 2021, 93, 5350–5357, https://doi.org/10.1002/jmv.27040.
- 133. Bergman, D.; Larsson, A.; Hansson-Hamlin, H.; Ahlén, E.; Holst, B.S. Characterization of canine anti-mouse antibodies highlights that multiple strategies are need to combat immunoassay interference. *Sci. Rep.* **2019**, *9*, 14521, https://doi.org/10.1038/s41598-019-51598-019-51228-3.
- Jara-Aguirre, J.C.; Baumann, N.A.; Block, D.R.; Algeciras-Schimnich, A. Human chorionic gonadotrophin suspected heterophile interference investigations in immunoassays: A recommended approach. *Clin. Chem. Lab. Med.* 2019, 57, 1192–1196, https://doi.org/10.1515/cclm-2018-1142.
- 135. Emerson, J.F.; Ngo, G.; Emerson, S.S. Screening for interference in immunoassays. *Clin. Chem.* 2003, 49, 1163–1169, https://doi.org/10.1373/49.7.1163.
- 136. Selby, C. Interference in immunoassay. Ann. Clin. Biochem. 1999, 36, 704–721, https://doi.org/10.1177/000456329903600603.
- 137. Tate, J.; Ward, G. Interferences in immunoassays. Clin. Biochem. Rev. 2004, 25, 105–120.
- 138. Sturgeon, C.M.; Viljoen, A. Analytical error and interference in immunoassay: Minimizing risk. *Ann. Clin. Biochem.* 2011, 48, 418–432, https://doi.org/10.1258/acb.2011.011073.
- 139. Rotmensch, S.; Cole, L.A. False diagnosis and needless therapy of presumed malignant disease in women with false-positive human chorionic gonadotropin concentrations. *Lancet* 2000, 355, 712–715, https://doi.org/10.1016/S0140-6736(00)01324-6.
- 140. Emerson, J.F.; Lai, K.K.Y. Endogenous antibody interferences in immunoassays. *Lab. Med.* 2013, 44, 69–73, https://doi.org/10.1309/LMMURCFQHKSB5YEC.
- Halvorsen, T.G.; Reubsaet, L. Determination of very low abundance diagnostic proteins in serum using immunocapture LC-MS/MS. Spectroscopy 2017, 15, 16–22.
- 142. Vehus, T. Performing quantitative determination of low-abundant proteins by targeted mass spectrometry liquid chromatography. *Intechopen* **2017**, *3*, 79–94, https://doi.org/10.5772/intechopen.68713.
- 143. Harney, D.J.; Hutchison, A.T.; Su, Z.; Hatchwell, L.; Heilbronn, L.K.; Hocking, S.; James, D.E.; Larance, M. Small-protein enrichment assay enables the rapid, unbiased analysis of over 100 low abundance factors from human plasma. *Mol. Cell. Proteom.* 2019, 18, 1899–1915, https://doi.org/10.1074/mcp.TIR119.001562.
- Voeten, R.L.C.; Ventouri, I.K.; Haselberg, R.; Somsen, G.W. Capillary electrophoresis: Trends and recent advances. *Anal. Chem.* 2018, 90, 1464–1481, https://doi.org/10.1021/acs.analchem.8b00015.
- 145. Sastre-Toraño, J.; Ramautar, R.; de Jong, G. Advances in capillary electrophoresis for life sciences. J. Chromatogr. B 2019, 1118, 116–136, https://doi.org/10.1016/j.jchromb.2019.04.020.
- 146. Wang, Y.; Adeoye, A.; Ogunkunle, E.O.; Wei, I.-A.; Filla, R.T.; Roper, M.G. Affinity capillary electrophoresis: A critical review of the literature from 2018 to 2020. *Anal. Chem.* **2021**, *93*, 295–310, https://doi.org/10.1021.acs.analchem.0c04526.
- 147. Dortez, S.; Sierra, T.; Crevillén, A.G.; Escarpa, A. CE/microchip electrophoresis of carbohydrates and glycoconjugates with electrochemical detection, 2nd ed.; In *Carbohydrate Analysis by Modern Liquid Phase Separation Techniques*; Rassi, Z.E., Ed.; Elsevier: Amsterdam, The Netherlands, 2021; Chapter 12, pp. 563–594; ISBN: 9780128214473.
- 148. Vitorino, R.; Guedes, S.; Pinto da Costa, J.; Kasicka, V. Microfluidics for peptidomics, proteomics, and cell analysis. *Nanomaterials* **2021**, *11*, 1118, https://doi.org/10.3390/nano11051118.
- El Deeb, S.; Fonseca-Silva, C.; Nascimento-Junior, C.; Hanafi, R.S.; Borges, K.B. Chiral capillary electrokinetic chromatography: Principle and applications, detection and identification, design of experiment, and exploration of chiral recognition using molecular modeling. *Molecules* 2021, 26, 2841, https://doi.org/10.3390/molecules26102841.
- 150. Peluso, P.; Chankvetadze, B. Native and substituted cyclodextrins as chiral selectors for capillary electrophoresis enantioseparations: Structures, features, application, and molecular modeling. *Electrophoresis* **2021**, https://doi.org/10.1002/elps.202100053.
- 151. Guzman, N.A. (Ed.) Capillary Electrophoresis Technology; Marcel Dekker Inc.: New York, NY, USA, 1993.
- 152. Pan, J.-Z.; Fang, P.; Fang, X.-X.; Hu, T.-T.; Fang, J.; Fang, Q. A low-cost palmtop high-speed capillary electrophoresis bioanalyzer with laser induced fluorescence detection. *Sci. Rep.* **2018**, *8*, 1791, https://doi.org/10.1038/s41598-018-20058-0.
- Gupta, V.; Dorsey, G.; Hubbard, A.E.; Rosenthal, P.J.; Greenhouse, B. Gel versus capillary electrophoresis genotyping for categorizing treatment outcomes in two anti-malarial trials in Uganda. *Malar. J.* 2010, *9*, 19:. https://doi.org/10.1186/1475-2875-9-19.
- 154. Zhu, Z.; Lu, J.J.; Liu, S. Protein separation by capillary electrophoresis: A review. Anal. Chim. Acta 2012, 709, 21–31, https://doi.org/10.1016/j.aca.2011.10.022.
- 155. Yu, R.B.; Quirino, J.P. Chiral selectors in capillary electrophoresis: Trends during 2017. *Molecules* 2019, 24, 1135, https://doi.org/10.3390/molecules24061135.
- 156. Kalíková, K.; Riesová, M.; Tesařová, E. Recent chiral selectors for separation in HPLC and CE. Open Chem. 2012, 10, 450–471, https://doi.org/10.2478/s11532-011-0142-3.
- Francotte, E.R. Polysaccharide derivatives as unique chiral selectors for enantioselective chromatography. *Chimia* 2017, 71, 430–450, https://doi.org/10.2533/chimia.2017.430.
- 158. Fang, L.; Du, Y.; Hu, X.; Luo, L.; Guo, X.; Guo, X.; Yu, J. Carboxymethyl β-cyclodextrin as chiral selector in capillary electrophoresis: Enantioseparation of 16 basic chiral drugs and its chiral recognition mechanism associated with drugs' structural features. *Biomed. Chromatogr.* **2017**, *31*, e3991, https://doi.org/10.1002/bmc.3991.

- 159. Blaschke, G.; Meryring, M.; Muhlenbrock, C.; Chankvetadze, B. Recent results of biotransformation of drugs: Investigation of the in vitro biotransformation of thalidomide using a dual cyclodextrin system in capillary electrophoresis. *Il Farmaco* 2020, *57*, 551–554, https://doi.org/10.1016/s0014-827x(02)01258-2.
- 160. Suba, D.; Urbányl, Z.; Salgó, A. Capillary isoelectric focusing method development and validation for investigation of recombinant therapeutic monoclonal antibody. J. Pharm. Biomed. Anal. 2015, 114, 53–61, https://doi.org/10.1016/j.jpba.2015.04.037.
- 161. Xu, T.; Sun, L. A mini review on capillary isoelectric focusing-mass spectrometry for top-down proteomics. *Front. Chem.* **2021**, *9*, 651757, https://doi.org/10.3389/fchem.201.651757.
- Costello, M.A.; Woititz, C.; DeFeo, J.; Stremlo, D.; Wen, L.-F.L.; Palling, D.J.; Iqbal, K.; Guzman, N.A. Characterization of humanized anti-Tac monoclonal antibody by traditional separation techniques and capillary electrophoresis. *J. Liq. Chromatogr. Rel. Technol.* 1992, 15, 1081–1097, https://doi.org/10.1080/10826079208018851.
- 163. Dadouch, M.; Ladner, Y.; Perrin, C. Analysis of monoclonal antibodies by capillary electrophoresis: Sample preparation, separation, and detection. *Separations* **2021**, *8*, 4, https://doi.org/10.3390/separations8010004.
- 164. Salas-Solano, O.; Kennel, B.; Park, S.S.; Roby, K.; Sosic, Z.; Boumajny, B.; Free, S.; Reed-Bogan, A.; Michels, D.; McElroy, W.; et al. Robustness of iCIEF methodology for the analysis of monoclonal antibodies: An interlaboratory study. *J. Sep. Sci.* 2012, 35, 3124–3129, https://doi.org/10.1002/jssc.201200633.
- 165. Osbourn, D.M.; Weiss, D.J.; Lunte, C.E. On-line preconcentration methods for capillary electrophoresis. *Electrophoresis* **2000**, *21*, 2768–2779, https://doi.org/10.1002/1522-2683(2000801)21:14<2768:AID-ELPS2768>3.0.CO;2-P.
- 166. Glatz, Z. On-capillary derivatization as an approach to enhancing sensitivity in capillary electrophoresis. *Electrophoresis* **2015**, 36, 744–763, https://doi.org/10.1002/elps.201400449.
- 167. Zhang, Z.; Zhang, F.; Liu, Y. Recent advances in enhancing the sensitivity and resolution of capillary electrophoresis. *J. Chromatogr. Sci.* 2013, *51*, 666–683.
- 168. Kawai, T. Recent advances in trace bioanalysis by capillary electrophoresis. *Anal. Sci.* 2021, 37, 27–36, https://doi.org/10.2116/analsci.20SAR12.
- Thomas, D.H.; Rakestraw, D.J.; Schoeniger, J.S.; Lopez-Avila, V.; Van Emon, J. Selective trace enrichment by immunoaffinity capillary electrochromatography on-line with capillary electrophoresis–laser induced fluorescence. *Electrophoresis* 1999, 20, 57– 66, https://doi.org/10.1002/(SICI)1522-2683(19990101)10:1<57::AID-ELPS57>3.0.CO;2-J.
- 170. Figeys, D.; Zhang, Y.; Aebersold, R. Optimization of solid phase microextraction-capillary zone electrophoresis-mass spectrometry for high sensitivity protein identification. *Electrophoresis* **1998**, *19*, 2338–2347, https://doi.org/10.1002/elps.1150191314.
- Zhang, H.; Zhou, L.; Chen, X. Improving sensitivity by large-volume sample stacking combined with sweeping without polarity switching by capillary electrophoresis coupled to photodiode array ultraviolet detection. *Electrophoresis* 2008, 29, 1556–1564; 10.1002/elps.200700759.
- 172. Wang, Y.-C.; Su, P.; Zhang, X.-X.; Chang, W.B. Enhancement of the sensitivity of a capillary electrophoresis immunoassay for estradiol with laser-induced fluorescence based on a fluorescein-labeled secondary antibody. *Anal. Chem.* **2001**, *73*, 5616–5619, https://doi.org/10.1021/ac010537a.
- 173. Marie, A.L.; Ray, S.; Lu, S.; Jones, J.; Ghiran, I.; Ivanov, A.R. High-sensitivity glycan profiling of blood-derived immunoglobulin G, plasma, and extracellular vesicles isolates with capillary zone electrophoresis-mass spectrometry. *Anal. Chem.* 2021, 93, 1991– 2002, https://doi.org/10.1021/acs.analchem.0c03102.
- 174. Guzman, N.A.; Guzman, D.E. A two-dimensional affinity capture and separation mini-platform for the isolation, enrichment, and quantification of biomarkers and its potential use for liquid biopsy. *Biomedicines* 2020, *8*, 255, https://doi.org/10.3390/biomedicines8080255.
- 175. Guzman, N.A.; Guzman, D.E. From a central laboratory to the beside: A point-of-care instrument to monitoring wellness and disease using two-dimensional immunoaffinity capillary electrophoresis technology. *Archiv. Biomed. Res.* **2018**, *1*.
- 176. Gao, Z.; Zhong, W. Recent (2018–2020) development in capillary electrophoresis. Anal. Bioanal. Chem. 2021, 1–16, https://doi.org/10.1007/s00216-021-03290-y.
- 177. Ta, H.Y.; Collin, F.; Perquis, L.; Poinsot, V.; Ong-Meang, V.; Couderc, F. Twenty years of amino acid determination using capillary electrophoresis: A review. *Anal. Chim. Acta* **2021**, *1174*, 338233, https://doi.org/10.1016/j.aca.2021.338233.
- 178. John, A.S.; Sidek, M.M.; Thang, L.Y.; Sami, S.; Tey, H.Y.; See, H.H. Online sample preconcentration techniques in nonaqueous capillary and microchip electrophoresis. J. Chromatogr. A 2021, 1638, 461868, https://doi.org/10.1016/j.chroma.2020.461868.
- 179. Kristoff, C.J.; Bwanali, L.; Veltri, L.M.; Gautam, G.P.; Rutto, P.K.; Newton, E.O.; Holland, L.A. Challenging bioanalyses with capillary electrophoresis. *Anal. Chem.* 2020, *92*, 49–66, https://doi.org/10.1021/acs.analchem.9b04718.
- Ouahabi, O.E.; Salim, H.; Pero-Gascon, R.; Benavente, F. A simple method for the analysis of extracellular vesicles enriched for exosomes from human serum by capillary electrophoresis with ultraviolet diode array detection. J. Chromatogr. A 2021, 1635, 461752, https://doi.org/10.1016/j.chroma.2020.461752.
- Oedit, A.; Hankemeier, T.; Lindenburg, P.W. On-line coupling of two-phase microelectroextraction to capillary electrophoresismass spectrometry for metabolomics analyses. *Microchem. J.* 2021, *162*, 105741, https://doi.org/10.1016/j.microc.2020.105741.
- Lim, R.R.X.; Fung, F.M.; Feng, H.-T.; Li, S.F.Y. Analysis of lipopolysaccharides by coupling microscale solid-phase extraction with capillary electrophoresis-laser induced fluorescence. *Microchem. J.* 2021, 161, 105771, https://doi.org/10.1016/j.microc.2020.105771.

- 183. Zhang, Z.; Park, J.; Barrett, H.; Dooley, S.; Davies, C.; Verhagen, M.C. Capillary electrophoresis-sodium dodecyl sulfate with laser-induced fluorescence detection as a highly sensitive and quality control-friendly method for monitoring adeno-associated virus capsid protein purity. *Hum. Gene Ther.* 2021, *32*, 628–637, https://doi.org/10.1089/hum.2020.233.
- Claude, B.; Nehmé, R.; Morin, P. Analysis of urinary neurotransmitters by capillary electrophoresis: Sensitivity enhancement using field-amplified sample injection and molecular imprinted polymer solid phase extraction. *Anal. Chim. Acta* 2011, 699, 242– 248, https://doi.org/10.1016/j.aca.2011.05.014.
- 185. Simonet, B.M.; Ríos Valcárcel, M. Enhancing sensitivity in capillary electrophoresis. *Trends Anal. Chem.* 2003, 22, 605–614, https://doi.org/10.1016/S0165-9936(03)01014-8.
- 186. Almeda, S.; Arce, L.; Valcárcel, M. The more and less common approaches to enhancing sensitivity in capillary electrophoresis. *Curr. Anal. Chem.* **2010**, *6*, 126–143, https://doi.org/10.2174/157341110790945508.
- 187. Drevinskas, T.; Telksnys, L.; Maruska, A.; Gorbatsova, J.; Kaljurand, M. Capillary electrophoresis sensitivity enhancement based on adaptive moving average method. *Anal. Chem.* **2018**, *90*, 6773–6780, https://doi.org/10.1021/acs.analchem.8b00664.
- 188. Galievsky, V.A.; Stasheushi, A.S.; Krylov, S.K. "Getting the best sensitivity from on-capillary fluorescence detection in capillary electrophoresis" A tutorial. Anal. Chim. Acta 2016, 935, 58–81, https://doi.org/10.1016/j.aca.2016.06.015.
- Wang, Y.; Fonslow, B.R.; Wong, C.C.; Nakorchevsky, A.; Yates, J.R. III. Improving the comprehensiveness and sensitivity of sheathless CE-MS/MS for proteomic analysis. *Anal. Chem.* 2012, *84*, 8505–8513, https://doi.org/10.1021/ac301091m.
- 190. Chen, D.; McCool, E.N.; Yang, Z.; Shen, X.; Lubeckyj, R.A.; Xu, T.; Wang, Q.; Sun, L. Recent advances (2019–2021) of capillary electrophoresis-mass spectrometry for multilevel proteomics. *Mass Spectrom. Rev.* **2021**, https://doi.org/10.1002/mas.21714.
- 191. Heegaard, N.H.H.; Nilsson, S.; Guzman, N.A. Affinity capillary electrophoresis: Important applications areas and some recent developments. *J. Chromatogr. B* 1998, 715, 29–54; 10.1016/s0378-4347(98)00258-8.
- 192. Olabi, M.; Stein, M.; Wätzig, H. Affinity capillary electrophoresis for studying interactions in life sciences. *Methods* **2018**, *146*, 76–92, https://doi.org/10.1016/j.ymeth.2018.05.006.
- 193. Nevídalová, H.; Michalcová, L.; Glatz, Z. Capillary electrophoresis-based approaches for the study of affinity interactions combined with various sensitive and nontraditional detection techniques. *Electrophoresis* **2019**, *40*, 625–642, https://doi.org/10.1002/elps.201800367.
- 194. Nevídalová, H.; Michalcová, L.; Glatz, Z. Capillary electrophoresis-based immunoassay and aptamer assay: A review. *Electrophoresis* **2020**, *41*, 414–433, https://doi.org/10.1002/elps.201900426.
- 195. Guzman, N.A. Disease Detection System and Method. U.S. Patent 10,408,789, 10 September 2019.
- 196. Guzman, N.A.; Guzman, D.E. A home-based portable instrument to monitor wellness and disease. Atlas of Science–Another View on Science. Avaliable online: https://atlasofscience.org/a-home-based-portable-instrument-to-monitor-wellness-and-disease/ (accessed on 21 March 2016).
- 197. Pero-Gascon, R.; Pont, L.; Sanz-Nebot, V.; Benavente, F. On-line immunoaffinity solid-phase extraction capillary electrophoresis-mass spectrometry for the analysis of transthyretin. *Methods Mol. Biol.* **2019**, 1972, 57–76, https://doi.org/10.1007/978-1-4939-9213-3_5.
- Puerta, A.; Martin-Alvarez, P.J.; Ongay, S.; Diez-Masa, J.C.; de Frutos, M. Immunoaffinity, capillary electrophoresis, and statistics for studying intact alpha 1-acid glycoprotein isoforms as an atherothrombosis biomarkers. *Methods Mol. Biol.* 2013, 919, 215–230, https://doi.org/10.10007/978-1-62703-029-8_20.
- Qi, X.-H.; Zhang, L.W.; Zhang, X.-X. Simultaneous determination of nandrolone, testosterone, and methyltestosterone by multiimmunoaffinity column and capillary electrophoresis. *Electrophoresis* 2008, 29, 3398–3405, https://doi.org/10.1002/elps.200700673.
- Garrido-Medina, R.; Farina-Gomez, N.; Diez-Masa, J.C.; de Frutos, M. Immunoaffinity chromatographic isolation of prostatespecific antigen from seminal plasma for capillary electrophoresis analysis of its isoforms. *Anal. Chim. Acta* 2014, 820, 47–55, https://doi.org/10.1016/j.aca.2014.02.031.
- 201. Phillips, T.M.; Wellner, E.F. Analysis of inflammatory mediators in newborn dried blood spot samples by chip-based immunoaffinity capillary electrophoresis. *Methods Mol. Biol.* 2019, 1972, 185–198, https://doi.org/10.1007/978-1-4939-9213-3_13.
- Phillips, T.M.; Wellner, E.F. Detection of cerebral spinal fluid associated chemokines in birth traumatized premature babies by chip-based immunoaffinity CE. *Electrophoresis* 2013, 34, 1530–1538, https://doi.org/10.1002/elps.201200634.
- 203. Guzman, N.A. Method and System for Simultaneous Determination of Multiple Measurable Biomarkers during the Development of Communicable Disease. U.S. Patent Application No. 2020/16/894,316. 5 June 2020.
- 204. Vermassen, T.; Van Praet, C.; Vanderschaeghe, D.; Maenhout, T.; Lumen, N.; Callewaert, N.; Hoebeke, P.; Van Belle, S.; Rottey, S.; Delanghe, J. Capillary electrophoresis of urinary prostate glycoproteins assists in the diagnosis of prostate cancer. *Electrophoresis* 2014, 35, 1017–1024, https://doi.org/10.1002/elps.201300332.
- 205. Kammeijer, G.S.M.; Nouta, J.; de la Rosette, J.J.M.C.H.; de Reijke, T.M.; Wuher, M. An in-depth glycosylation assay for urinary prostate-specific antigen. *Anal. Chem.* **2018**, *90*, 4414–4421, https://doi.org/10.1021/acs.analchem.7b04281.
- 206. Reider, B.; Gacsi, E.; Jankovics, H.; Vonderviszt, F.; Szarvas, T.; Guttman, A.; Jarvas, G. Integrated workflow for urinary prostate specific antigen N-glycosylation anlysis using sdAb partitioning and downstream capillary electrophoresis separation. *Anal. Chim. Acta* 2021, 338892, in press, https://doi.org/10.1016/j.aca.2021.338892.
- 207. Dimitrov, D.S. Therapeutic proteins. Methods Mol. Biol. 2012, 899, 1–26, https://doi.org/10.1007/978-1-61779-921-1_1.
- Leader, B.; Baca, Q.J.; Golan, D.E. Protein therapeutics: A summary and pharmacological classification. *Nat. Rev. Drug Discov.* 2008, 7, 21–39, https://doi.org/10.1038/nrd2399.

- Chennamsetty, N.; Voynov, V.; Kayser, V.; Helk, B.; Trout, B.L. Design of therapeutic proteins with enhanced stability. *Proc. Natl. Acad. Sci. USA* 2009, 106, 11937–11942, https://doi.org/10.1073/pnas.0904191106.
- Kariolis, M.S.; Wells, R.C.; Getz, J.A.; Kwan, W.; Mahon, C.S.; Tong, R. Brain delivery of therapeutic proteins using Fc fragment blood-brain barrier transport vehicle in mice and monkeys. *Sci. Transl. Med.* 2020, 12, eaay1359, https://doi.org/10.1126/scitranslmed.aay1359.
- 211. Schadt, S.; Hauri, S.; Lopes, F.; Edelmann, M.R.; Staack, R.F.; Villaseñor, R.; Kettenberger, H.; Roth, A.B.; Schuler, F.; Richter, W.F.; et al. Are biotransformation studies of therapeutic proteins needed? Scientific considerations and technical challenges. *Drug Metabol. Dispos.* 2019, 47, 1443–1456, https://doi.org/10.1124/dmd.119.088997.
- Scientific Advice from European Medicines Agency (EMA). Guideline on Immunogenicity Assessment of Therapeutic Proteins. 2017. Available online: https://www.ema.europa.eu/en/documents/scientific-guideline/guideline-immunogenicity-assessmenttherapeutic-proteins-revision-1_en.pdf (accessed on 18 May 2017).
- Rosenberg, A.S. Immunogenicity Risk Assessment for Therapeutic Proteins: A Guide to Risk Assessment for Therapeutic Peptides and Generic Peptides. 2014. Available online: https://www.fda.gov/media/146112/download (accessed on 31 August 2014).
- 214. Ratanji, K.D.; Derrick, J.P.; Dearman, R.J.; Kimber, I. Immunogenicity of therapeutic proteins: Influence of aggregation. J. Immunotoxicol. 2014, 11, 99–109, https://doi.org/10.3109/1547691X.2013.821564.
- Solá, R.J.; Griebenow, K. Glycosylation of therapeutic proteins: An effective strategy to optimize efficacy. *BioDrugs* 2010, 24, 9– 21, https://doi.org/10.2165/11530550-000000000-00000.
- Roseman, D.S.; Weinberger, R. Quantitative capillary zone electrophoresis method for the precise determination of charge differences arising from the manufacture of heparan-N-sulfatase. J. Pharm. Biomed. Anal. 2013, 85, 67–73, https://doi.org/10.1016/j.jpba.2013.05.018.
- 217. Papathanasiou, M.M.; Kontoravdi, C. Engineering challenges in therapeutic protein product and process design. *Curr. Opin. Chem. Engin.* **2020**, *27*, 81–88, https://doi.org/10.1016/j.coche.2019.11.010.
- 218. Morrow, T.; Felcone, L.H. Defining the difference: What makes biologics unique. Biotechnol. Health 2004, 1, 24–29.
- Suresh, S.; Rajvanshi, P.K.; Noguchi, C.T. The many facets of erythropoietin physiologic and metabolic response. *Front. Physiol.* 2020, 10, 1534, https://doi.org/10.3389/fphys.2019.01534.
- Peng, B.; Kong, G.; Yang, C.; Ming, Y. Erythropoietin and its derivatives: From tissue protection to immune regulation. *Cell Death Dis.* 2020, 11, 79, https://doi.org/10.1038/s41419-020-2276-8.
- 221. Debeljak, N.; Solár, P.; Sytkowski, A.J. Erythropoietin and cancer: The unintended consequences of anemia correction. *Front. Immunol.* **2014**, *5*, 563, https://doi.org/10.3389/fimmu.2014.00563.
- 222. Nagano, M.; Stübiger, G.; Marchetti, M.; Gmeiner, G.; Allmaier, G.; Reichel, C. Detection of isoforms of recombinant human erythropoietin by various plant lectins after isoelectric focusing. *Electrophoresis* **2005**, *26*, 1633–1645, https://doi.org/10.1002/elps.200410214.
- Bristow, A.; Charton, E. Assessment of the suitability of a capillary electrophoresis method for determining isoform distribution of erythropoietin. *Pharmeuropa* 1999, 11, 290–300.
- 224. Reily, C.; Stewart, T.J.; Renfrow, M.B.; Novak, J. Glycosylation in health and disease. *Nat. Rev. Nephrol.* 2019, 15, 346–366, https://doi.org/10.1038/s41581-019-0129-4.
- Alocci, D.; Ghraichy, M.; Barletta, E.; Gastaldello, A.; Mariethoz, J.; Lisacek, F. Understanding the glycome: An interactive view of glycosylation from glycocompositions to glycoepitopes. *Glycobiology* 2018, 28, 349–362, https://doi.org/10.1093/glycob/cwy019.
- 226. Lauc, G.; Pezer, M.; Rudan, I.; Campbell, H. Mechanisms of disease: The human N-glycome. *Biochim. Biophys. Acta* 2016, 1860, 1574–1582, https://doi.org/10.1016/j.bbagen.2015.10.016.
- 227. Flynn, R.A.; Pedram, K.; Malaker, S.A.; Batista, P.J.; Smith, B.A.H.; Johnson, A.G.; George, B.M.; Majzoub, K.; Villalta, P.W.; Carette, J.E.; Bertozzi, C.R. Small RNAs are modified with N-glycans and displayed on the surface of living cells. *Cell* 2021, 184, 3109–3124, https://doi.org/10.1016/j.cell.2021.04.023.
- Franc, V.; Zhu, J.; Heck, A.J.R. Comprehensive proteoform characterization of plasma complement component C8abj by hybrid mass spectrometry approaches. J. Am. Soc. Mass Spectrom. 2018, 29, 1099–1110, https://doi.org/10.1007/s13361-018-1901-6.
- 229. Pang, X.; Li, H.; Guan, F.; Li, X. Multiple roles of glycans in hematological malignancies. *Front. Oncol.* 2018, *8*, 364, https://doi.org/10.3389/fonc.2018.00364.
- 230. Meany, D.L.; Chan, D.W. Aberrant glycosylation associated with enzymes as cancer biomarkers. Clin. Proteom. 2011, 8, 2–14.
- Poiroux, G.; Barre, A.; Rougé, P.; Benoist, H. Targeting glycosylation aberrations to improve the efficiency of cancer phototherapy. *Curr. Cancer Drug Targ.* 2019, 19, 349–359, https://doi.org/10.2174/1568009618666180628101059.
- 232. Tuccillo, F.M.; de Laurentiis, A.; Palmieri, C.; Fiume, G.; Bonelli, P.; Borrelli, A.; Tassone, P.; Scala, I.; Buonaguro, F.M.; Quinto, I.; et al. Aberrant glycosylation as biomarker for cancer: Focus on CD43. *BioMed Res. Int.* 2014, 2014, 742831, https://doi.org/10.1155/2014/742831.
- Ip, C.K.M.; Yin, J.; Ng, P.K.S.; Lin, S.-Y.; Mills, G. Genomic-glycosylation aberrations in tumor initiation, progression and management. AIMS Med. Sci. 2016, 3, 386–416, https://doi.org/10.3934/medsci.2016.4.386.
- 234. Venkitachalam, S.; Revoredo, L.; Varadan, V.; Fecteau, R.E.; Ravi, L.; Lutterbaugh, J.; Markowitz, S.D.; Willis, J.E.; Gerken, T.A.; Guda, K. Biochemical and functional characterization of glycosylation-associated mutational landscapes in colon cancer. *Sci. Rep.* 2016, *6*, 23642, https://doi.org/10.1038/srep23642.

- 235. Hu, C.W.; Qiu, Y.; Ligeralde, A.; Raybon, A.Y.; Yoo, S.Y.; Coombes, K.R.; Qutub, A.A.; Kornblau, S.M. A quantitative analysis of heterogeneities and hallmarks in acute myelogenous leukaemia. *Nat. Biomed. Enginer.* **2019**, *3*, 889–901, https://doi.org/10.1038/s41551-019-0387-2.
- 236. Gratacós-Mulleras, A.; Duran, A.; Shehni, A.A.; Ferrer-Batallé, M.; Ramírez, M.; Comet, J.; de Llorens, R.; Saldova, R.; Llop, E.; Peracaula, R. Characterisation of the main PSA glycoforms in aggressive prostate cancer. *Sci. Rep.* 2020, 10, 18974:. https://doi.org/10.1038/s41598-020-75526-3.
- 237. Saldova, R.; Fan, Y.; Fitzpatrick, J.M.; Watson, R.W.G.; Rudd, P.M. Core fucosylation and α2-3 sialylation in serum N-glycome is significantly increased in prostate cancer comparing to benign prostate hyperplasia. *Glycobiology* 2011, 21, 195–205, https://doi.org/10.1093/glycob/cwq147.
- 238. Zhou, J.; Yang, W.; Hu, Y.; Höti, N.; Liu, Y.; Shah, P.; Sun, S.; Clark, D.; Thomas, S.; Zhang, H. Site-specific fucosylation analysis identifying glycoproteins associated with aggressive prostate cancer cell lines using tandem affinity enrichments of intact glycopeptides followed by mass spectrometry. *Anal. Chem.* 2017, *89*, 7623–7630, https://doi.org/10.1021/acs.analchem.7b01493.
- 239. Besarab, A.; Drueke, T.B. The problem with transferrin saturation as a key indicator of iron 'sufficient' in chronic kidney disease. *Nephrol. Dial. Transplant.* **2021**, *36*, 1377–1383, https://doi.org/10,1093/ndt/gfaa048.
- 240. Formanowicz, D.; Formanowicz, P. Transferrin changes in haemodialised patients. Int. Urol. Nephrol. 2012, 44, 907–919, https://doi.org/10.1007/s11255-011-9947-4.
- 241. Vogt, A.-C.; Arsiwala, T.; Mohsen, M.; Manolova, V.; Bachmann, M.F. On iron metabolism and its regulation. *Int. J. Mol. Sci.* **2021**, 22, 4591, https://doi.org/10.3390/ijms22094591.
- 242. Bartnikas, T.B. Known and potential roles of transferrin in iron biology. *Biometals* 2012, 25, 677–686, https://doi.org/10.1007/s10534-012-9520-3.
- 243. Arndt, T. Carbohydrate-deficient transferrin as a marker of chronic alcohol abuse: A critical review of preanalysis, analysis, and interpretation. *Clin. Chem.* **2001**, 47, 13–27, https://doi.org/10.1093/clinchem/47.1.13.
- 244. Zühlsdorf, A.; Said, M.; Seger, C.; Park, J.H.; Reunert, J.; Rust, S.; Wada, Y.; Grüneber, M.; DuChesne, I.; Marquardt, T. It is not always alcohol abuse-A transferrin variant impairing the CTC test. *Alcohol Alcohol.* 2016, 51, 148–153, https://doi.org/10.1093/alcalc/agv099.
- 245. Lilja, H.; Ulmert, D.; Vickers, A.J. Prostate-specific antigen and prostate cancer: Prediction, detection and monitoring. *Nat. Rev. Cancer* 2008, *8*, 268–278, https://doi.org/10,1038/nrc2351.
- Medina-Casanellas, S.; Benavente, F.; Giménez, E.; Barbosa, J.; Sanz-Nebot, V. On-line immunoaffinity solid phase extraction capillary spectrometry for the analysis of large biomolecules: A preliminary report. *Electrophoresis* 2014, 35, 2130–2136, https://doi.org/10.1002/elps.201400119.
- 247. Barrabés, S.; Farina-Gomez, N.; Llop, E.; Puerta, A.; Diez-Masa, J.C.; Perry, A.; de Llorens, R.; de Frutos, M.; Peracaula, R. Comparative analysis of prostate-specific antigen by two-dimensional gel electrophoresis and capillary electrophoresis. *Electrophoresis* **2017**, *38*, 408–416, https://doi.org/10.1002/elps.201600432.
- 248. Horiba Scientific-Capillary Electrophoresis Application Note CEAN02-Protein Analysis-Analysis of Transferrin Glycoforms Using EVA and GST Algorithms. Available online: https://www.horiba.com/fileadmin/uploads/Scientific/Documents/Capillary_Electrophoresis/ce02-app-note.pdf (accessed on 1 January 2021).
- Nuevo-Ordonez, Y.; Anton, R.F.; Davis, W.C. Quantification of total serum transferrin and transferrin sialoforms in human serum; an alternative method for the determination of carbohydrate-deficient transferrin in clinical samples. *Anal. Methods* 2014, 6, 3967, https://doi.org/10.1039/c4ay00159a.
- 250. Zühlsdorf, A.; Park, J.H.; Wada, Y.; Rust, S.; Reunert, J.; DuChesne, I.; Grüneberg, M.; Marquardt, T. Transferrin variants: Pitfalls in the diagnosis of congenital disorders of glycosylation. *Clin. Biochem.* 2015, 48, 11–13, https://doi.org/10.1016/j.clinbiochem.2014.09.022.
- 251. Gruszewska, E.; Cylwik, B.; Gudowska, M.; Kedra, B.; Szmitkowski, M.; Chrostek, L. Changes in transferrin isoforms in pancreatic cancer. *Ann. Clin. Lab. Sci.* **2016**, *46*, 286–290.
- 252. Guzman, N.A. Consecutive protein digestion and peptide derivatization employing an on-line analyte concentrator to map proteins using capillary electrophoresis. In *CRC Series in Analytical Biotechnology;* Righetti, P.G., Ed.; CRC Press: Boca Raton, FL, USA, 1996; Chapter 4, pp. 101–121.
- 253. Guzman, N.A.; Blanc, T.; Phillips, T.M. Immunoaffinity capillary electrophoresis as a powerful strategy for the quantification of low-abundance biomarkers, drugs, and metabolites in biological matrices. *Electrophoresis* 2008, 29, 3259–3278, https://doi.org/10.1002/elps.200800058.
- 254. Chang, W.W.P.; Hobson, C.; Bomberger, D.C.; Schneider, L.V. Rapid separation of protein isoforms by capillary zone electrophoresis with dynamic coatings. *Electrophoresis* **2005**, *26*, 2179–2186, https://doi.org/10.1002/elps.200410283.
- Carillo, S.; Jakes, C.; Bones, J. In-depth analysis of monoclonal antibodies using microfluidic capillary electrophoresis and native mass spectrometry. J. Pharm. Biomed. Anal. 2020, 185, 113218, https://doi.org/10.1016/j.jpba.2020.113218.
- 256. Stolz, A.; Hedeland, Y.; Salzer, L.; Römer, J.; Heiene, R.; Leclercq, L.; Cottet, H.; Bergquist, J.; Neusüß, C. Capillary zone electrophoresis-top-down tandem mass spectrometry for in-depth characterization of hemoglobin proteoforms in clinical veterinary samples. *Anal. Chem.* **2020**, *92*, 10531–10539, https://doi.org/10.1021/acs.analchem.0c01350.
- Shen, X.; Yang, Z.; McCool, E.N.; Lubeckyj, R.A.; Chen, D.; Sun, L. Capillary electrophoresis-mass spectrometry for top-down proteomics. *Trends Analyt. Chem.* 2019, 120, 115644, https://doi.org/10.1016/j.trac.2019.115644.

- 258. Chen, C.-H.; Feng, H.; Guo, R.; Li, P.; Laserna, A.K.C.; Ji, B.; Ng, B.H.; Li, S.F.Y.; Khan, S.H.; Paulus, A.; et al. Intact NIST monoclonal antibody characterization-Proteoforms, glycoforms-using CE-MS and CE-LIF. *Cogent Chem.* 2018, 4, 1480455, https://doi.org/10.1080/23312009.2018.1480455.
- Thomas, S.L.; Thacker, J.B.; Schug, K.A.; Maráková, K. Sample preparation and fractionation techniques for intact proteins for mass spectrometric analysis. J. Sep. Sci. 2021, 44, 211–246, https://doi.org/10.1002/jssc.2020000936.
- 260. Kaur, H.; Beckman, J.; Zhang, Y.; Li, Z.J.; Szigeti, M.; Guttman, A. Capillary electrophoresis and the biopharmaceutical industry: Therapeutic protein analysis and characterization. *Trends Anal. Chem.* 2021, 144, 116407; https://doi.org/10.1016/j.trac.2021.116407.
- Giorgetti, J.; Beck, A.; Leize-Wagner, E.; François, Y.-N. Combination of intact, middle-up and bottom-up levels to characterize 7 therapeutic monoclonal antibodies by capillary electrophoresis – Mass spectrometry. J. Pharm. Biomed. Anal. 2020, 182, 113107, doi:10.1016/j.jpba.2020.113107.
- McCool, E.N.; Lubeckyj, R.A.; Shen, X.; Chen, D.; Kou, Q.; Liu, X.; Sun, L. Deep top-down proteomics using capillary zone electrophoresis-tandem mass spectrometry: Identification of 5700 proteoforms from the *Escherichia coli* proteome. *Anal. Chem.* 2018, 90, 5529–5533, https://doi.org/10.1021/acs.analchem.8b00693.
- 263. Wu, H.; Tang, K. Highly sensitive and robust capillary electrophoresis-electrospray ionization-mass spectrometry: Interfaces, preconcentration techniques and applications. *Rev. Anal. Chem.* **2020**, *39*, 45–55, https://doi.org/10.1515/revac-2020-0112.
- 264. Sarg, B.; Faserl, K.; Kremser, L.; Halfinger, B.; Sebastiano, R.; Lindner, H.H. Comparing and combining capillary electrophoresis electrospray ionization mass spectrometry and nano-liquid chromatography electrospray ionization mass spectrometry for the characterization of post-translationally modified histones. *Mol. Cell. Proteom.* 2013, 12, 2640–2656, https://doi.org/10.1074/mcp.M112.024109.
- Ren, T.-J.; Zhang, X.-X.; Li, X.; Chen, H.-X. Isoforms of recombinant human erythropoietin by polarity-reversed capillary isoelectric focusing. *Electrophoresis* 2020, 41, 2055-2061, https://doi.org/10.1002/elps.202000165.
- 266. Andrasi, M.; Pajaziti, B.; Sipos, B.; Nagy, C.; Hamidli, N.; Gaspar, A. Determination of deamidated isoforms of human insulin using capillary electrophoresis. *J. Chromatogr. A* 2020, *1626*, 461344, https://doi.org/10.1016/j.chroma.2020.461344.
- 267. Crihfield, C.L.; Kristoff, C.J.; Veltri, L.M.; Penny, W.M.; Holland, L.A. Semi-permanent cationic coating for protein separations. *J. Chromatogr. A* **2019**, *1607*, 460397, https://doi.org/10.1016/j.chroma.2019.460397.