

Supplementary Material


Red blood cells thermodynamic behavior in neurodegenerative pathologies and aging

Svetla Todorova¹, Sashka Krumova¹, Desislava Bogdanova², Avgustina Danailova¹, Elena Zlatareva², Nikolay Kalaydzhev², Ariana Langari¹, Ivan Milanov², Stefka G. Taneva^{1,*}


¹Institute of Biophysics and Biomedical Engineering, Bulgarian Academy of Sciences, "Acad. G. Bonchev" Str. 21, 1113 Sofia, Bulgaria

²University multiprofile hospital for active treatment in neurology and psychiatry "Sv. Naum", Louben Roussev Str. 1, Sofia 1113, Bulgaria


Supplementary Figures




Supplementary Figure S1. Series of calorimetric curves of normal healthy RBCs measured along the cells aging (denoted HC1 – HC9), the days of the DSC measurements are indicated for each healthy donor in the corresponding panel.



Supplementary Figure S2. Series of calorimetric curves of RBCs derived from the nine PD patients along the cells aging (denoted PD1 – PD9), for each studied PD case the days of the DSC measurements are indicated in the corresponding panel.



Supplementary Figure S3. Series of calorimetric curves of RBCs derived from the eight ALS patients during cells aging (denoted ALS1 – ALS8), the days of the DSC measurements are indicated for each studied ALS case in the corresponding panel.



Supplementary Figure S4. Series of calorimetric curves of RBCs derived from three AD patients along the cells aging (denoted AD1 – AD3), the days of the DSC measurements are indicated for each studied AD case in the corresponding panel.