

Article

Relationship between Brain-Derived Neurotrophic Factor and Cognitive Decline in Patients with Mild Cognitive Impairment and Dementia

Matea Nikolac Perkovic ¹, Fran Borovecki ^{2,3}, Igor Filipcic ⁴, Barbara Vuic ¹, Tina Milos ¹, Gordana Nedic Erjavec ¹, Marcela Konjevod ¹, Lucija Tudor ¹, Ninoslav Mimica ^{3,5}, Suzana Uzun ^{3,5}, Oliver Kozumplik ⁵, Dubravka Svob Strac ¹ and Nela Pivac ^{1,6,*}

¹ Laboratory for Molecular Neuropsychiatry, Division of Molecular Medicine, Ruder Boskovic Institute, 10000 Zagreb, Croatia; mnikolac@irb.hr (M.N.P.); barbara.vuic@irb.hr (B.V.); tina.milos@irb.hr (T.M.); gnedic@irb.hr (G.N.E.); mkonjevod@irb.hr (M.K.); ltudor@irb.hr (L.T.); dsvob@irb.hr (D.S.S.)

² Department of Neurology, University Hospital Centre Zagreb, 10000 Zagreb, Croatia; fbor@mef.hr

³ School of Medicine, University of Zagreb, 10000 Zagreb, Croatia;

ninoslav.mimica@bolnica-vrapce.hr (N.M.); suzana.uzun@bolnica-vrapce.hr (S.U.)

⁴ Psychiatric Hospital “Sveti Ivan”, 10090 Zagreb, Croatia; igor.filipcic@pbsvi.hr

⁵ Department for Biological Psychiatry and Psychogeriatrics, University Psychiatric Hospital Vrapče, 10090 Zagreb, Croatia; oliver.kozumplik@bolnica-vrapce.hr

⁶ University of Applied Sciences Hrvatsko Zagorje Krapina, 49000 Krapina, Croatia

* Correspondence: npivac@irb.hr




Figure S1. Comparison of the results (Kruskal-Wallis ANOVA by ranks), concerning BDNF plasma concentration in MCI subjects and patients diagnosed with AD, between current study (cohort 2) and previous study by Domitrovic Sudic and colleagues [59] (cohort 1).

AD, Alzheimer's Disease; BDNF, Brain-Derived Neurotrophic Factor; HC, Healthy Control; MCI, Mild Cognitive Impairment

* $p \leq 0.001$ vs. HC (Dunn's post-hoc test)

** $p \leq 0.001$ vs. MCI, Cohort 2 (Dunn's post-hoc test)

*** $p < 0.050$ vs. MCI, Cohort 1 (Dunn's post-hoc test)