

Article

# Open Geospatial System for LUCAS In Situ Data Harmonization and Distribution

Martin Landa <sup>\*,†</sup> , Lukáš Brodský <sup>†</sup> , Lena Halounová , Tomáš Bouček  and Ondřej Pešek 

Department of Geomatics, Faculty of Civil Engineering, Czech Technical University in Prague, 166 29 Prague, Czech Republic; lukas.brodsky@fsv.cvut.cz (L.B.); lena.halounova@fsv.cvut.cz (L.H.); tomas.boucek@fsv.cvut.cz (T.B.); ondrej.pesek@fsv.cvut.cz (O.P.)

\* Correspondence: martin.landa@fsv.cvut.cz

† These authors contributed equally to this work.

**Abstract:** The use of in situ references in Earth observation monitoring is a fundamental need. LUCAS (Land Use and Coverage Area frame Survey) is an activity that has performed repeated in situ surveys over Europe every three years since 2006. The dataset is unique in many aspects; however it is currently not available through a standardized interface, machine-to-machine. Moreover, the evolution of the surveys limits the performance of change analysis using the dataset. Our objective was to develop an open-source system to fill these gaps. This paper presents a developed system solution for the LUCAS in situ data harmonization and distribution. We have designed a multi-layer client-server system that may be integrated into end-to-end workflows. It provides data through an OGC (Open Geospatial Consortium) compliant interface. Moreover, a geospatial user may integrate the data through a Python API (Application Programming Interface) to ease the use in workflows with spatial, temporal, attribute, and thematic filters. Furthermore, we have implemented a QGIS plugin to retrieve the spatial and temporal subsets of the data interactively. In addition, the Python API includes methods for managing thematic information. The system provides enhanced functionality which is demonstrated in two use cases.

**Keywords:** LUCAS; in situ; data harmonization; data distribution; web services; QGIS plugin



**Citation:** Landa, M.; Brodský, L.; Halounová, L.; Bouček, T.; Pešek, O. Open Geospatial System for LUCAS In Situ Data Harmonization and Distribution. *ISPRS Int. J. Geo-Inf.* **2022**, *11*, 361. <https://doi.org/10.3390/ijgi11070361>

Academic Editors: Dev Raj Paudyal and Wolfgang Kainz

Received: 6 May 2022

Accepted: 20 June 2022

Published: 23 June 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

The use of independent in situ references in Earth observation monitoring is a fundamental need [1]. Currently, the volume of generated geospatial datasets is increasing significantly into big data [2]. Such data are characterized by their large volume, high value, high variety, and potentially high velocity and high veracity [3]. In order to provide the geoscience community the full opportunity to uncover previously unknown insights, such datasets should be assessed through a standardized, scalable, and extensible technology allowing full integration into end-to-end scientific workflows.

The European continental land cover mapping activities, which use Earth observation images, require reliable and representative in situ references for the validation and calibration of automated mapping across the whole of Europe [4]. Such references are typically developed in land cover production campaigns ad hoc from aerial orthophotos or very high resolution images, but in situ surveys are rarely performed. The difficulties with representative references are magnified when land cover change detection mapping is performed.

LUCAS, the Land Use and Coverage Area frame Survey, is an activity managed by Eurostat, which has performed in situ surveys over Europe every three years since 2006 [5]. Today, there is a series of 2006, 2009, 2012, 2015, and 2018 observations. The surveyors mainly examine the in situ land cover (76 classes) and land use (41 classes) and take photos (one facing photo and four landscape photos in the cardinal compass directions), but also evaluate agro-environmental information and take a 500-gram topsoil sample at one out

of ten points. Recently (2018 survey), the Copernicus, INSPIRE (Infrastructure for Spatial Information in Europe), and EUNIS (European Nature Information System) attributes were added. The primary objective of the sampling was to provide area estimates for spatial and territorial analyses such as agricultural statistics [6]. However, the use of the LUCAS dataset is manifold. There are 168,402 (2006), 234,623 (2009), 270,272 (2012), 339,696 (2015), and 337,854 (2018) sample points surveyed in the LUCAS dataset. This gives a set of 651,377 individual surveyed points on the territory of the European Union (EU) member states, while 319,150 samples have been visited at least twice and 35,204 samples visited repeatedly in all the surveyed years, allowing land changes to be evaluated. The datasets total 1,350,847 individual in situ analyzed points.

There are other sampling activities, such as Geo-Wiki.org [7] and ESA GlobCover [8], providing validation datasets; these are working, however, with a considerably smaller amount of collected data and without consistently repeated observations over time. In this respect, the LUCAS dataset is unique in several aspects, including its spatial sampling density ( $2 \times 2$  km grid) covering the EU member states, the total number of in situ visited sites, its temporal resolution (three-year frequency), and its thematic coverage.

The LUCAS dataset has been used in numerous research activities. Close et al. [9] and Weigand et al. [10] used the LUCAS dataset as training and validation reference data together with Sentinel-2 for a per-pixel supervised classification at the national level of land use and land cover. Pflugmacher et al. [11] mapped the pan-European land cover using the Landsat spectral-temporal metrics based on the European LUCAS 2018 survey. Gao et al. [12] used LUCAS data to evaluate the current global land cover maps over the EU. d'Andrimont et al. [13] used LUCAS data, together with Sentinel-1 images, to map crop types at a 10 m spatial resolution at the European level for 2018. Borrelli et al. [14] integrated a soil erosion module with the 2018 LUCAS topsoil survey to monitor the soil health status across the EU and to support actions to prevent soil degradation. All the above-mentioned authors used the LUCAS dataset as a status dataset in their analysis; however, they only used one year of the survey, while the potential of repeated in situ measurements remained unused. We believe that the data maintain a far higher potential.

Nevertheless, as the LUCAS activity has evolved since 2006, there are differences in the separate surveys that need to be harmonized. d'Andrimont et al. [13] proposed and applied a number of harmonization steps related to the renaming of the database columns, re-coding of the variables, and correcting the theoretical location coordinates. Weigand et al. [10] proposed several pre-processing schemes for the LUCAS data (two positioning approaches and three semantic selection approaches). Use of the LUCAS data showed a positive effect on the accuracy of land cover classification, and especially the positional correction of points.

We further argue the importance of the open reference data sets being available on-line using state-of-the-art geospatial technology to allow full integration of the data (machine-to-machine) into automated processing pipelines, for instance, in cloud environments. It is natural to use an OGC compliant interface, allowing wide integration using standards. Interoperability and standardization are crucial for achieving information system modularity in order to seamlessly connect various software modules or components as presented in the publication of Jeppesen et al. [15]. The OGC standards have been designed to assure the interoperability of Geographical Information Systems (GIS) and the Spatial Data Infrastructure (SDI) in general [16]. Many OGC compliant implementations of scientific workflows are available [17–20]. An example of the automated coordination of OGC web services to produce thematic maps is described in the publication of Rautenbach et al. [21]. OGC specifies several open data formats and web services suitable for geospatial data distribution and data exchange. Of particular interest are the Web Feature Service (WFS) [22] and the OGC API Features [23] suited for requesting raw vector data. In OGC-based SDI, a user is able to seamlessly access data stored in different file-based or database-oriented geospatial formats [24]. Vector features are commonly stored and maintained by object-relational database management systems with a geospatial exten-

sion installed. In open-source settings, the PostgreSQL database system with a geospatial PostGIS extension is widely adopted as a OGC compliant database [25,26].

Currently, individual LUCAS datasets are available for downloading in Comma Separated Values (CSV) files from the Eurostat portal [27], and the photos are available through the online LUCAS Viewer [28]. This type of distribution technology lacks integration convenience, which prevents the data from being employed in wider end-to-end workflows. We reviewed the individual surveys while preparing spatio-temporal Land Cover mapping over the period between 2000 and 2019 on the European scale [29] and identified the following three areas of possible enhancement: (1) harmonization of the data values in the database (including variable names and data types), (2) spatio-temporal aggregation of the individual datasets (including harmonization of the spatial coordinates), and (3) establishment of an OGC compliant distribution system to provide interactive machine-to-machine accessibility.

This paper proposes and develops a technological solution to retrieve the spatial and temporal subsets of harmonized LUCAS data as part of the Geo-harmonizer project development and shares the system as an open-source code with the geoscience community.

Here, we present a SpaceTime\_LUCAS (ST\_LUCAS) system that we developed to process the three areas mentioned above. We divided the overall aim into several objectives:

- O1 data storage in a persistence layer;
- O2 full and configurable automation of the harmonization process for past and future LUCAS survey updates and space–time aggregation for change analysis;
- O3 development of software to access the data via a standardized (OGC) web service;
- O4 development of a client Python API and QGIS plugin to retrieve the subsets of LUCAS data based on spatial, temporal, and thematic filters;
- O5 development of translation methods to provide LUCAS land cover data in other nomenclatures and allow user-defined analytics such as legend aggregation.

Finally, we demonstrate the use of the developed system in two use cases in Section 5.

## 2. Materials and Methods

The architectural design of the ST\_LUCAS system was derived from the objectives listed in the introduction. The methodological process follows several steps. Given the objectives (system requirements), we designed a high-level static architecture with generic functionality. For each of the system components, we specified the purpose, function, inputs, and outputs. This enabled us to evaluate the overall functionality of the system. Next, we identified the interfaces between the components and designed the data flow. After the static architecture, we designed a system dynamic architecture to evaluate the system interaction in a sequential manner. We also defined a set of tests to validate all the components, interfaces, and data.

We designed the system using general workflow diagrams in the Unified Modeling Language (UML) 2.5, especially the UML component and sequential diagrams. The system was coded using Bash, Python 3, and Structured Query Language (SQL). The overall system was deployed using the Docker virtualization technology.


### *LUCAS Data Harmonization*

The Eurostat LUCAS primary data collected during the five surveys of 2006, 2009, 2012, 2015, and 2018 [30–34] are the main inputs to the system. The individual surveys distributed as CSV files were downloaded to the file system along with the respective technical documentation. The theoretical LUCAS 2018  $2 \times 2$  km<sup>2</sup> grid from Eurostat [35] was also added.

Having defined the second objective (O2) as being to harmonize the LUCAS dataset across the survey years, we first reviewed the instructions for surveyors and classification technical references [36–39] and the record descriptors of the primary data [40–44] to identify the changes between individual LUCAS surveys, which evolved between 2006 and 2018. These findings formed the basis for the harmonization process. Generally,


the process involved the harmonization of spatial coordinates, renaming of attributes, harmonization of data values, and unification of data types. We consider the 2018 survey to be the most developed, and it also consists new thematic information; hence we used this year as a reference. All the database attributes and their values are harmonized to this reference. These harmonization steps were cross-checked with assumptions made by d'Andrimont et al. [13] during the development of the system in the Geo-harmonizer project. The harmonized attributes of the ST\_LUCAS dataset are listed in Appendix A Table A2. Users may also explore the coding and mapping tables online on the ST\_LUCAS website ([https://geoforall.fsv.cvut.cz/st\\_lucas](https://geoforall.fsv.cvut.cz/st_lucas), release 1.0 published by the authors on 9 June 2022).

Individual harmonized datasets from 2006, 2009, 2012, 2015, and 2018 were consequently merged into a common database through the space–time aggregation. The aim was to create a common LUCAS database where the vector of thematic information which allows direct evaluation of, for instance, land cover changes over individual years may be retrieved by the user for each point visited repeatedly in situ. Given the fact that the respective visits were measured by GPS, the spatial coordinates of the collected information differ in time for each point with a unique ID. The differences vary from meters to hundreds of meters (Figure 1 below).



**Figure 1.** Distance between GPS measured and theoretical points (OBS\_DIST attribute). A total of 35,509 points with a distance of more than 1000 m are not shown in this figure.

Therefore, we calculated the representative geometry as a geometric median of the repeated GPS measurements (Figure 2). The median was computed by the PostGIS function [45] using the Weiszfeld algorithm [46] to avoid the influence of possible outliers that are present. In addition, we introduced a new attribute (SURVEY\_DIST, Appendix A Table A2), which measures the distance between the survey GPS coordinates and the geometric median coordinates. All the spatial coordinates (measured, theoretical, and median) were transformed to the common ETRS89-extended/LAEA Europe coordinate reference system (EPSG 3035).



**Figure 2.** Space–time aggregation of observed LUCAS GPS locations (circle symbol) using the geometrical median (diamond symbol). The theoretical location snapped to a LUCAS  $2 \times 2$  km<sup>2</sup> grid is represented by a rectangle symbol. The distances between the GPS and median locations are shown by arrowed dashed lines.


### 3. System Design

This section presents the system design. It is split into several linked sub-sections defining the system at different abstraction levels. The design starts with a high-level static architecture and the main system components, their logical structure, and decomposition. Next, the interfaces are identified and elaborated, and, finally, the system behavior is designed in the dynamic architecture.

#### 3.1. System High-Level Architecture

The system architecture follows a typical multi-layer client-server model customized to meet the objectives defined in Section 1. The system partitions the tasks between the server side with the LUCAS dataset and clients, and the users consuming the spatio-temporal subsets of the LUCAS data. Moreover, the design also follows the idea of Component-Based Software Engineering (CBSE) as described, for instance, by Vale et al. [47].

The system, as depicted in Figure 3, is decomposed into three main layers consisting of several components (Table 1) in each of the layers. The persistence layer provides a means to store the input, intermediate (separate years), and final harmonized space–time aggregated LUCAS data (O1). The main role of this layer is to preserve the data in non-volatile storage for further distribution. The main component of this layer is a database system (spatially enabled SQL database). The application layer provides the system deployment and harmonization (O2) procedures and the data distribution web service (O3). The distribution functionality is achieved by the OGC Web Feature Service. The client layer communicates through an internal interface with the data distribution service that provides harmonized LUCAS data to the end user. The client layer’s main functionality is provided by the Python software package. The main requirement is to create requests based on user-defined spatial, attribute, thematic, and temporal filters. The client layer provides two ways to retrieve and filter the subsets of the data (O4), through a command line interface (CLI) or through a graphical user interface (GUI). The client layer has an additional defined functionality to translate the LUCAS land cover nomenclature to other legends (O5) or aggregate the legend based on customer requirements.



**Figure 3.** Static high-level architecture of the ST\_LUCAS system. The Jupyter Notebook (dotted line) is not intended to be a software component. The Jupyter Notebook is used in this work to present the functionality of the Python package (C1) and to provide manual verification.

**Table 1.** Overview of the ST\_LUCAS system components.


ID	Name	Layer	Role	Objective
P1	File system	Persistence	Store primary data	O1
P2	Database	Persistence	Store and provide harmonized data	O1
A1	Deployment package	Application	Deploy the system including data harmonization	O2
A2	Web service	Application	Provide access to harmonized data through a web service	O3
C1	Python package	Client	API interface to a web service and a set of analytical functions	O4, O5
C2	Local file system	Client	Store locally harmonized LUCAS data	O4, O5
C3	QGIS plugin	Client	Provide GUI interface via GIS to a web service and a set of selected analytical functions	O4, O5

The system is designed for three potential high-level users. The system administrator is able to deploy the system as it is designed or configure it for customized application use. The geospatial developer is able to access the harmonized LUCAS dataset through the Python API (e.g., using the Jupyter Notebook or directly by the Python code) in a fully automated processing pipeline. The GIS user is able to interactively explore and download the subsets of the LUCAS dataset via the desktop QGIS application (QGIS plugin).

The ST\_LUCAS system is devised to allow full scalability. The vertical scalability may be achieved by deploying the system in a cloud environment. The horizontal scalability of the presented system may be achieved by adding multiple map server instances in the application layer.

### 3.2. System Interfaces

The overall system is designed as encapsulated individual software components, which provide a set of related functions. The software components communicate with each other via interfaces [48]. There are two types of interfaces: internal interfaces and external interfaces. The internal and external interfaces are illustrated using the UML component diagram in Figure 4. The internal interface provides communication between the persistence layer and the application layer (IF1). It is an interface between the database system and the web service using the OGC WFS specification where harmonized LUCAS data flow. The other internal interface provides communication between the application layer (web service) and the client Python package (IF2), which is defined by the OGC WFS specification. The external interfaces are between the client application (CLI or GUI) and the client layer Python package (IF3). Here, the web service sends requests to the spatio-temporal database and the service provides harmonized LUCAS data to the end users' client in the form of the Geography Markup Language (GML) features. A QGIS plugin/Jupyter Notebook (client layer) communicates with the ST\_LUCAS system via the Python package API. This interface provides the access and functionalities to retrieve and filter harmonized spatio-temporal LUCAS data subsets from the application layer.



**Figure 4.** ST\_LUCAS system software components with defined interfaces.

### 3.3. System Dynamic Architecture

The dynamic architecture was designed by means of UML sequential diagrams to plan the interactions of the above-designed system components over time. The dynamic architecture was divided into two phases: the system deployment phase and the user interaction process.

Figure 5 presents a UML sequential diagram in the deployment phase, which is the interaction between the main components of the application layer (deployment package and web service) and the persistence layer (file system and database system). The process starts with the configuration of the initial system, according to the administrator's requirements. Next, the deployment package (A1) performs three major operations. Firstly, the LUCAS primary data distributed in the plain CSV format are automatically downloaded from the data provider specified in the system configuration. Secondly, after successfully

downloading the primary CSV files to the server file system (P1), the database system (P2) is deployed. Deployment of the database system is split into ten sub-operations starting with a database initialization step and followed by importing the primary LUCAS data into the initialized database. The harmonization process includes the unification of spatial coordinates, attribute names, attribute data types, and data values. These operations are performed separately for each survey year. Subsequently, space–time aggregation of the harmonized LUCAS observations is applied. The deployment process is completed by creating a database dump file for the creation of the spatio-temporal database. Thirdly, the system web service (A2) component is deployed. Next, the harmonized LUCAS data in the database system may be published by the web service. Each of the sequential steps is followed by tests to evaluate the system performance.




Figure 5. ST\_LUCAS dynamic architecture (system deployment).

Figure 6 presents a UML sequential diagram of the user interaction process. This process is the interaction between the main components of the client side (CLI/GUI clients—C3, Python package—C1) and the server side represented by the application layer web service (A2) providing the data from the persistence layer (database system—P2). In addition, the users' local file system (C2) is introduced in Figure 3 to illustrate the dynamic interaction with the ST\_LUCAS system.




Figure 6. ST\_LUCAS dynamic architecture of the system user interaction.

The end user is able to interact with the system using a GUI designed for the open-source QGIS platform or through the Python API. The QGIS plugin (C3) is built on the top of the Python package (C1) allowing easy access to harmonized space–time LUCAS data. Alternatively, the Jupyter Notebook may be used for code-based data interaction in the Python programming language.

Initially, the server access has to be configured to access a web service (A2) through the Python package (C1) on the client side. As soon as the server (web service) provides information on the successful connection, the user is able to define the data requests. The user builds a request based on spatial, attribute, thematic, and temporal filters in order to obtain a subset of the harmonized LUCAS dataset. The request is sent to the server-side web service component (A2) by the Python package (C1). The response from the server comprises a subset of the harmonized LUCAS dataset that corresponds to the filters defined in the user's request. The data retrieved from a web service (A2) are stored by the Python package (C1) on the local file system (C2).

### 3.4. System Validation

System validation is a constituent part of the system design. The validation is performed at two levels. At the design (documentation) level, which allows traceability between the model components in the architecture phase, and the software level to test each component and the interface of the designed system at the system deployment, execution, and external users' access.

We defined unit tests for each software component according to the detailed architectural design of the system (Figures 3–5). The ST\_LUCAS system deployment process is controlled by the A1 software component. The process is formed by three major steps as shown in Figure 5: (1) download primary data (P1), (2) deploy database (P2), and (3) deploy web service (A2). Unit tests (1) validate the LUCAS primary data download process as specified in the configuration. The following set of unit tests (2) validates whether DB (P2) was initialized (2a), the primary data imported (2b), the LUCAS data harmonization process applied (2c–2h), the harmonized LUCAS data prepared for publication (2i), and the DB recovery file created (2j). The last set of unit tests (3) validates whether the web service

(A2) is operational (3a) and the ST\_LUCAS data (3b) and metadata (3c) are published. For the client layer, the unit tests cover the Python package (C1) and the local file system (C2) components. For the QGIS plugin (C3), manual verification is performed. An overview of the unit tests is depicted in Table 2. Each test is evaluated as successful if the test result is in compliance with the system configuration; otherwise, it is evaluated as failed.

**Table 2.** Overview of unit tests. Test IDs reflect the system dynamic architecture as presented in Figure 5.

Component ID	Test IDs	Description
A1, P1	1_001	Primary data are downloaded according to the system configuration.
	2a_001	DB is initialized according to the system configuration.
	2b_001-003	Primary data are imported according to the system configuration.
	2c_001-002	Coordinates are harmonized according to the system configuration.
A1, P2	2d_001	Attributes are harmonized according to the system configuration.
	2e_001-002	Data values are harmonized according to the system configuration.
	2f_001	Data types are harmonized according to the system configuration.
	2g_001-004	Harmonized data are merged according to the system configuration.
	2h_001-003	Data are space–time aggregated according to the system configuration.
	2i_001-004	Publication views are created according to the system configuration.
A1, A2	2j_001	DB recovery file is created according to the system configuration.
	3a_001-003	Test case consists of checking OGC WFS operations: GetCapabilities, DescribeFeatureType and GetFeature.
	3b_001-003	ST_LUCAS dataset available via WFS.
A1, A2	3c_001-003	The test cases consist of checking that ST_LUCAS metadata are published according to the deployed database.
	C1, C2	001-007

The interaction between the software components is validated by integration tests. For each interface (Figure 4), a set of integration tests was designed (Table 3). Various combinations of spatial, temporal, attribute, and thematic filters are tested. A request based on a specified set of filters is built by the Python package (C1) and sent to the server. The web service (A2) returns a relevant subset of harmonized LUCAS data (IF2). The subset is compared with a recordset provided by a direct query to the database (P2) using SQL statements (IF1). The integration tests are evaluated with a pass result only if both interfaces (IF1 and IF2) return the same subset of the LUCAS data. The system administrator is notified about the integration test results regularly by email notifications. It ensures that system administrators will be informed about a potential system failure in real time, which is an important aspect of the operational deployment of the ST\_LUCAS system. For the IF3 interface, manual verification was performed via the Jupyter Notebook.

**Table 3.** Overview of integration tests. Interface IDs reflect the system architecture as presented in Figure 4.

Interface ID	Test IDs	Description
IF1, IF2	001–004	Test cases consist of checking WFS responses retrieved by the Python package (IF2) covering various combinations of spatial, attribute, thematic, and temporal filters. The responses are compared with the subsets retrieved from spatio-temporal DB via SQL statements (IF1). Test cases pass only if there is no difference between the WFS responses and the subsets retrieved from DB.

The unit (Table 2) and integration (Table 3) tests are automated with the Python pytest package [49]. Verification is, therefore, performed during the software execution, which dynamically checks the software behavior. The unit test results are verified by the system administrator at the deployment phase. In the event that one of the unit tests fails, the deployment process is terminated, and the system is not deployed. The integration test results are verified repeatedly at the system execution phase. The result of each test operation is sorted in a log file, which is publicly available (Demonstration of the ST\_LUCAS system, [https://geoforall.fsv.cvut.cz/st\\_lucas](https://geoforall.fsv.cvut.cz/st_lucas), release 1.0 published by the authors on 9 June 2022) and may, therefore, be verified by the user.

#### 4. System Implementation

The ST\_LUCAS system implementation and deployment follow the design as described in Section 3. The system is completely based on open-source software components to ensure its reproducibility, transparency, and extensibility (Appendix A Table A1). Overall, the architecture splits the system into the client (frontend) and the server side (backend); hence, the implementation aspects are described separately.

##### 4.1. Backend

The backend, the server side consisting of the persistence and application layers, is composed of software components that are responsible for the data storage, harmonization of LUCAS data, and data distribution. The components (represented in Figure 3) are implemented as a collection of services (see Section 4.3). Each service is managed in an isolated Docker container running on any operating system supported by the Docker virtualization technology [50]. There are two core backend software components: the database management system, operating a spatio-temporal DB (P2), and the map server, providing a web service (A2). The database management system is represented by an object-relational open-source PostgreSQL server [51]. This database system has a strong reputation for reliability, data integrity, and correctness. Moreover, it provides extension to geospatial data, the so-called PostGIS, which “spatially enables” the PostgreSQL database [52]. PostGIS also follows the OpenGIS “Simple Features Specification for SQL” by OGC. Here, we use the PostGIS version 3.1. The P2 and A2 components are deployed by the system deployment package (A1).

The database population process is controlled by a collection of scripts implemented in the Python 3 and Bash programming languages covering all steps as described in the dynamic system architecture (Figure 5). The harmonization process of the LUCAS data is procedural. It is decomposed as a series of computational steps carried out at the database level by scripts written in SQL. The configuration of the harmonization process is managed by a collection of CSV and JavaScript Object Notation (JSON) files. The robust but still easy-to-use configuration allows system administrators to affect the result of the harmonization process. In the same way, the system may be easily extended by new LUCAS observations planned to be published in 2022. The GDAL library [53] is used to import the primary LUCAS data from the file system (P1) to the PostGIS database tables (P2).

The web service (A2) is implemented as a map server, provided by the open-source GeoServer software version 2.19 [54]. The GeoServer is an OGC compliant implementation of a number of open standards such as the Web Map Service (WMS) and the Web Coverage Service (WCS). Additional formats and publication options are available as extensions, including the Web Processing Service (WPS) and the Web Map Tile Service (WMTS). It also conforms to the OGC WFS standard, which allows the sharing of the vector features to be used on the client side. Users are able to incorporate the LUCAS dataset into their processing pipelines and applications, freeing the data and permitting greater transparency. When the data becomes available from the deployment process, the GeoServer provides harmonized LUCAS data via a standardized WFS interface through Hypertext Transfer Protocol (HTTP).

The deployment package source code is available from the GitLab repository ([https://gitlab.com/geoharmonizer\\_inea/st\\_lucas/st\\_lucas-system-deployment](https://gitlab.com/geoharmonizer_inea/st_lucas/st_lucas-system-deployment), release 1.0 published by the authors on 9 June 2022).

#### 4.2. Frontend

The frontend, the client side, is composed of the Python package (C1) and the CLI client or GUI clients, as depicted in Figure 3. The Python package maintains the main functionality of the client side to be used by the end-user, either through the QGIS plugin (C3) without the necessity of any additionally programming/scripting, while the geospatial developer may use the Python API package directly to integrate harmonized LUCAS subsets into their own coded Earth observation pipelines.

##### 4.2.1. Python Package

The frontend is developed based on the core component, the Python package (C1), which provides communication between the web server (A2) and the users' clients through API. The package implemented in the Python 3 programming language consists of three modules: (1) `request` to create a request by specifying spatial, temporal, attribute, or thematic filters; (2) `io` to retrieve harmonized LUCAS data provided by the web service (A2) based on the submitted request, and to store the retrieved data in the user local file system (C2) in a specified data format; and (3) `analyze` to process the received harmonized LUCAS data—LUCAS land cover classes aggregation and nomenclature translation.

To use the Python API, the user creates a request by the `LucasRequest` Python class. In a single request, a combination of spatial, temporal, attribute, and thematic filters may be used. The spatial filter may be defined either by a bounding box, the NUTS0 country code, or by a user-defined polygon vector layer. Spatial coordinates must be specified in the ETRS89-extended/LAEA Europe coordinate reference system (EPSG 3035). The spatial filter is required by request; the other filters are optional. The temporal filter is specified by a list of survey years to be queried. The attribute filter is provided by an operator, an attribute name, and a list of values. The thematic filter defines the subset of harmonized LUCAS attributes to be retrieved. The example below (Listing 1) demonstrates the combination of all possible filters; the spatial filter is defined by a bounding box (`request.bbox`) covering the whole EU territory; the temporal filter (`request.years`) restricts the result to the 2015 and 2018 survey years; and the attribute filter (`request.propertyname`, `request.operator`, `request.literal`, and `request.logical`) selects only the LUCAS observations with land cover classes (LUCAS attribute LC1), C21 (Spruce dominated coniferous woodland), or C22 (Pine dominated coniferous woodland). The thematic filter (`request.group`) defines a subset of harmonized LUCAS attributes only to those which are relevant to the “Land Cover, Land Use” thematic group (LC\_LU code in Appendix A Table A2).

#### Listing 1. Build a request.

---

```

from st_lucas import LucasRequest
from owslib.fes import PropertyIsEqualTo, Or

request = LucasRequest()
request.bbox = (1510105, -2292253, 8582000, 5306000)
request.years = [2015, 2018]
request.propertyname = 'LC1'
request.operator=PropertyIsEqualTo
request.literal = ['C21', 'C22']
request.logical = Or
request.group = 'LC_LU'

```

---

The LUCAS subset retrieval according to the defined filters and data storage is managed by a `LucasIO` Python class. The LUCAS data are retrieved from the web service (A2) by the `LucasIO.download()` method based on the specified `LucasRequest` class instance. The number of retrieved LUCAS samples is returned by the `LucasIO.count()` method (Listing 2).

**Listing 2.** Download LUCAS subset based on the request.

---

```
from st_lucas import LucasIO

lucasio = LucasIO()
lucasio.download(request)
print('Number of LUCAS points:', lucasio.count())
```

---

The downloaded LUCAS data may be stored on a local file system (C2) as an OGC GeoPackage file by the `LucasIO.to_gpkg()` method for further processing in GIS applications or converted by the `LucasIO.to_geopandas()` method into a `GeoDataFrame` object, which may be processed and analyzed in the Python code by the `GeoPandas` library [55].

The `analyze` Python module provides two analytical functionalities. The Python class `LucasClassAggregate` implements the LUCAS land cover classes aggregation. In addition to the aggregation of land cover classes, the module also offers the possibility of translating the LUCAS nomenclature into other nomenclatures using the `LucasClassTranslate` Python class. Complex Python API usage is demonstrated in several use cases and is discussed in Section 5.


The client Python API may be used directly by the Python code (CLI), running the Jupyter Notebook, or by the QGIS developed plugin (GUI). The Jupyter Notebook as a web-based interactive computing platform allows interactive data exploration [56] for geospatial developers. The Python package source code and the Jupyter Notebooks demonstrating the capabilities of the system are available in the GitLab repository ([https://gitlab.com/geoharmonizer\\_inea/st\\_lucas/st\\_lucas-python-package](https://gitlab.com/geoharmonizer_inea/st_lucas/st_lucas-python-package), release 1.0 published by the authors on 9 June 2022).

#### 4.2.2. QGIS Plugin

Another option for harmonized space–time LUCAS dataset exploration is to use the `ST_LUCAS` QGIS plugin. The plugin is a GUI integrated into the open-source QGIS platform. The user interface (Figure 7) is split into three tabs: (1) Download; (2) Analyze; and (3) Photos. The added value of using GIS is to interactively select the spatial, attribute, temporal, and thematic filters from the GUI (Download tab). In particular, it enables selection of an area of interest (spatial filter) by the extent of the map canvas, specification of a country from a list of EU countries or use of a user-defined vector polygon data layer. The plugin interface allows a list of selected years (temporal filter) and a group of attributes (thematic filter) to also be specified. There are five thematic groups in total to choose from, where each group contains specific thematic attributes in addition to the basic attributes. The following groups are available: Land Cover, Land Use; Land Cover, Land Use, Soil; Forestry; Copernicus; and Inspire PLCC (Appendix A Table A2). The user is also able to download LUCAS points with all available attributes.

The Analyze tab integrates functionality for performing a user-defined class aggregation using a JSON file and a nomenclature translation using a CSV file. An example of usage is included in the plugin documentation available online on the `ST_LUCAS` website ([https://geoforall.fsv.cvut.cz/st\\_lucas/qgis\\_plugin/](https://geoforall.fsv.cvut.cz/st_lucas/qgis_plugin/), release 1.0 published by the authors on 9 June 2022).


Users are also able to browse photos (a facing photo and four landscape photos in the cardinal compass directions) of a selected LUCAS point in the Photo tab (Figure 8) as provided by the GISCO service [57].



**Figure 7.** ST\_LUCAS QGIS plugin (highlighted by a red box) retrieving harmonized LUCAS data for the Czech Republic territory (background basemap: OpenStreetMap—public WMS view service).

The subset of harmonized space–time LUCAS data retrieved from the server is stored in the OGC GeoPackage format with a predefined style for further usage. The style of LUCAS points is defined to distinguish the land cover classes at the first level of the LUCAS nomenclature as shown in Figure 7. In addition, points with a circular symbol indicate that photos are available for display. Points with a square symbol indicate the opposite.

The QGIS plugin source code is available in the GitLab repository ([https://gitlab.com/geoharmonizer\\_inea/st\\_lucas/st\\_lucas-qgis-plugin](https://gitlab.com/geoharmonizer_inea/st_lucas/st_lucas-qgis-plugin), release 1.0 published by the authors on 9 June 2022).



**Figure 8.** Showing LUCAS photos from the GISCO service by the ST\_LUCAS QGIS plugin.

### 4.3. ST\_LUCAS System Deployment

In order to enhance the portability of the developed ST\_LUCAS system, the Docker virtualization technology [50] was employed. All the backend components are deployed through virtualization. There are four deployed services managed by the Docker-compose tool, each running in an isolated Docker container: (1) db: responsible for the spatio-temporal database (P2) deployment including harmonized space–time LUCAS data; (2) gs: responsible for running a map server providing the OGC WFS service (A2); (3) gsp: responsible for publishing harmonized space–time LUCAS data as a WFS service. The publication process is performed when the database deployment is successfully completed. At this point, the system metadata and documentation are also populated. The service is terminated when the publishing process is successfully completed; and (4) gst: performs repetitive integration tests (Table 3) controlling whether the system is operational. Logs are publicly available; and the system administrator is notified by email.

The ST\_LUCAS system deployment controls components in the persistence and application layers as shown in Figure 3. The spatio-temporal DB (P2) is managed by a Docker container (db), which mounts the server file system (P1) as a volume. As the system deployment package (A1) manages all the deployment steps, it is available to all Docker containers (db, gs, gsp, and gst) through the mounted volume.

The complete deployment of the ST\_LUCAS system is performed by a single command: `docker-compose up`.

The demonstration ST\_LUCAS system installation, available at [https://geoforall.fsv.cvut.cz/st\\_lucas](https://geoforall.fsv.cvut.cz/st_lucas) (release 1.0 published by the authors on 9 June 2022), is deployed on GNU/Debian operating system version 11.

## 5. Discussion

The developed ST\_LUCAS system has multiple uses. System managers may take the system and install it (Docker containers) in their institutes as it is (objective O1). Geospatial developers may utilize (e.g., configure) the current system and let it run for their own purposes (O2). Geospatial users may access the harmonized spatio-temporal LUCAS dataset, and its subsets, using either the machine-to-machine Python API or the interactive QGIS plugin (O4). The data are provided by a standardized interface defined by OGC WFS (O3). The system offers developers and geospatial users full and configurable automation of the harmonization process for past and future LUCAS survey updates, spatio-temporal and thematic subsetting of the data based on user-defined filters, land cover legend translation to a defined nomenclature, and legend aggregation (O5).

Next, we demonstrate and discuss the use of the ST\_LUCAS system via two use cases (Sections 5.1 and 5.2), in addition to the use of the system described by Witjes et al. [29] for the European scale land cover mapping between 2000 and 2019.

### 5.1. LUCAS Data for Land Cover Change Analysis

In the first use case, we demonstrate how to retrieve a vector of changing land cover from the ST\_LUCAS system. Assuming the geospatial user wants to calibrate the classification change model or validate the existing land cover change product, we need to retrieve a subset of data with repeated visits on the same geographical points. The task may be simply set up by selecting LUCAS points where repeated visits are higher than one. As mentioned in Section 1, a total of 319,150 sample points are visited at least twice, and 35,204 samples are visited five times.

Below, a Python snippet code (Listing 3) demonstrates how to retrieve LUCAS points with repeated visits for land cover change analysis using the SURVEY\_COUNT attribute for AOI (areas of interest) in the Czech Republic. There were 11,084 points retrieved with the initial spatial and temporal filter, while by adding the condition SURVEY\_COUNT > 1 we received a subset of 6175 points.

**Listing 3.** Build a request for land cover change analysis.

---

```

from st_lucas import LucasRequest
from owslib.fes import PropertyIsGreaterThan

request = LucasRequest()


request.countries = ['CZ']
request.st_aggregated = True
request.group = 'LC_LU'

request.propertyname = 'SURVEY_COUNT'
request.operator = PropertyIsGreaterThan
request.literal = 1

```

---

In Figure 9, we illustrate an example of the land cover change vector (LC1 codes: B16, B15, B55, E10, and E20) for the case of five in situ repeated visits (2006, 2009, 2012, 2015, and 2018), together with orthophotos in the background. It clearly shows the changing landscape documented by the orthophotos and the changing LC codes in the LUCAS database.



**Figure 9.** Example of changing land cover over time for POINT\_ID = 46642928 as recorded in the LUCAS dataset (background orthophotos: Czech State Administration of Land Surveying and Cadastre—public WMS view service).

### 5.2. LUCAS Data for Land Product Validation

In the next use case, we demonstrate the use of LUCAS data for land product validation. In this case, validation of the national-level Land Parcel Identification System (LPIS) of the Czech Republic for 2018, which is an open dataset [58]. We simplified the validation into two agriculture classes, cropland (class 1) and grassland (class 2), which represent

the majority of the agricultural land in LPIS. Initially, we retrieved the subset of LUCAS data with the respective spatial and temporal filters in the validation process, similarly to the code snippet in Listing 1. Next, we used the `LucasClassAggregate` Python class from the developed Python package to simplify the nomenclature to the above-defined legend (Listing 4). The method's argument is a Python dictionary defining the class mappings for the aggregation process.

**Listing 4.** Perform land cover class aggregation.

---

```

from st_lucas import LucasClassAggregate

lc1_to_agri = {
"1": ["B11", "B12", "B13", "B14", "B15", "B16", "B17", "B18",
      "B19", "B21", "B22", "B23", "B31", "B32", "B33", "B34",
      "B35", "B36", "B37", "B41", "B42", "B43", "B44", "B45",
      "B51", "B52", "B53", "B54", "B55", "B71", "B72", "B73",
      "B74", "B75", "B76", "B77", "B81", "B82", "B83", "B84"],
"2": ["E10", "E20", "E30"]
}

lucasaggr = LucasClassAggregate(lucasio.data, mappings=lc1_to_agri)
lucasaggr.apply()

```

---

The retrieved vector layer with the aggregated nomenclature was overlaid with the LPIS product and validation indicators were calculated (Table 4). It is possible to conclude that the LPIS product has a high thematic accuracy, with an overall F1-score of 96%, compared to the LUCAS in situ survey of 2018.

**Table 4.** LPIS validation indicators.

Class	Code	Support	F1-Score	Precision	Recall
Cropland	1	1941	98.1	97.1	99.1
Grassland	2	690	94.1	96.1	92.1
Overall		2631	96.1	97.1	95.1

## 6. Conclusions

This article presents a developed ST\_LUCAS geospatial data system. A versatile open-source framework for the LUCAS dataset harmonization, distribution by the OGC compliant interface, Python client API and QGIS plugin to retrieve subsets of data, and methods to manage nomenclature translation, class aggregation, and thematic information. The source code, documentation, and installation instructions are publicly available on GitLab ([https://gitlab.com/geoharmonizer\\_inea/st\\_lucas](https://gitlab.com/geoharmonizer_inea/st_lucas), release 1.0 published by the authors on 9 June 2022). The ST\_LUCAS deployment package and the Python client package are published under the MIT license, the QGIS plugin under GNU GPL v3.

The system is designed in a multi-layer client-server model allowing integration to end-to-end workflows. Integration of the system into full end-to-end workflows is facilitated by the Python API. The transferability of the system to different server end-points is facilitated by OS-level virtualization using Docker containers. This allows a diverse audience of geospatial developers and scientists to use the capabilities of the ST\_LUCAS system in their own environments. Moreover, the system may be configured based on specific user requirements. The system is prepared for the new LUCAS survey of 2022 as soon as it is publicly available.

We discuss the use of the ST\_LUCAS system via two use cases (Section 5). The workflow management capabilities have been already tested to prepare a harmonized spatio-temporal LUCAS dataset for the European scale land cover mapping between 2000 and

2019 [29]. The experience gained from this large-scale use case proved the necessity of auxiliary methods for the land cover nomenclature translation and aggregation, which is not a common feature in purely distribution systems.

Overall, we believe the ST\_LUCAS system provides better accessibility and usability of the European LUCAS dataset through a standardized OGC interface. The specific enhancement is in the space–time aggregation of the data that emphasizes the use of a highly valuable in situ survey database for change analysis as a priority of all land monitoring projects. In addition, the ST\_LUCAS system is fully integrated into the QGIS desktop platform, allowing interactive exploration of the LUCAS data and direct GIS analysis. As a continuation of the ST\_LUCAS development, we plan to run extended demonstration use cases. In particular, we intend to explore the value of repeated in situ visits for land cover change detection. To fulfill this task, we shall develop translation tables from the LUCAS land cover legend to various nomenclatures. Furthermore, we shall explore the spatial representativeness of the sampled points in order to use the data for validation and land cover calibration activities in varying spatial resolutions.

The ST\_LUCAS system was implemented based on current knowledge of the primary data acquired in 2018 and the previous surveys performed since 2006. The system may require future updates according to changes in the LUCAS dataset from subsequent surveys.

**Author Contributions:** Conceptualization, Martin Landa and Lukáš Brodský; methodology, Martin Landa, Lukáš Brodský and Lena Halounová; software, Martin Landa, Tomáš Bouček and Ondřej Pešek; validation, Martin Landa and Tomáš Bouček; formal analysis, Lukáš Brodský; investigation, Lukáš Brodský; resources, Lukáš Brodský; data curation, Tomáš Bouček; writing—original draft preparation, Lukáš Brodský, Martin Landa and Tomáš Bouček; writing—review and editing, Lena Halounová and Ondřej Pešek; visualization, Martin Landa, Lukáš Brodský and Tomáš Bouček; supervision, Lena Halounová; project administration, Martin Landa; and funding acquisition, Martin Landa. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by the Grant Agreement Connecting Europe Facility (CEF) Telecom project grant number 2018-EU-IA-0095 Geo-harmonizer project (2019–2022).

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** The LUCAS primary data are available from EUROSTAT free of charge to all users (<https://ec.europa.eu/eurostat/en/web/lucas/data/primary-data>, accessed on 26 April 2022).

**Acknowledgments:** Administrative and technical support by the Faculty of Civil Engineering, Czech Technical University in Prague. The authors would like to thank reviewers for taking the time and effort necessary to review the manuscript. The authors sincerely appreciate all valuable comments and suggestions, which helped to improve the quality of the manuscript.

**Conflicts of Interest:** The authors declare no conflict of interest.

## Abbreviations

The following abbreviations are used in this manuscript:

API	Application Programming Interface
CBSE	Component-Based Software Engineering
CLI	Command Line Interface
CSV	Comma Separated Values (file format)
EPSG	EPSG Geodetic Parameter Dataset
EU	European Union
EUNIS	European Nature Information System
GIS	Geographic Information System
GUI	Graphical User Interface
HTTP	Hypertext Transfer Protocol
INSPIRE	Infrastructure for Spatial Information in Europe

JSON	JavaScript Object Notation
LC	Land Cover
LUCAS	Land Use and Coverage Area frame Survey
NUTS	Nomenclature of Territorial Units for Statistics
OGC	Open Geospatial Consortium
PLCC	Pure Land Cover Components
SDI	Spatial data Infrastructure
SQL	Structured Query Language
UML	Unified Modeling Language
WCS	Web Coverage Service
WFS	Web Feature Service
WMS	Web Map Service
WMTS	Web Map Tile Service
WPS	Web Processing Service

## Appendix A

**Table A1.** Open-source software used by the ST\_LUCAS system. Component IDs reflect the system architecture as presented in Figure 3.

Component ID	Software	License
P2	PostgreSQL	PostgreSQL licence
P2	PostGIS	GNU GPL
A1	Docker CE	N/A (free of charge)
A1	Docker Compose	Apache License 2.0
A1	psycopg2 *	GNU LGPL v3
A1	gdal *	MIT
A1	pytest *	MIT
A1	owslib *	BSD 3
A1	geoserver-rest *	MIT
A1	requests *	Apache 2.0
A2	GeoServer	GNU GPL
C1	json/os/csv/logging/tempfile/pathlib/shutil *	PSF 2.2/BSD 0
C1	gdal *	MIT
C1	owslib *	BSD 3
C1	requests *	Apache 2.0
C3	QGIS	GNU GPL

\* Python package.

**Table A2.** List of ST\_LUCAS attributes.

Attribute	Group	Description	Units	Origin
POINT_ID	DEFAULT	Unique point identifier		Primary
NUTS0	DEFAULT	NUTS Lvl 0		Primary
NUTS1	DEFAULT	NUTS Lvl 1		Primary
NUTS2	DEFAULT	NUTS Lvl 2		Primary
NUTS3	DEFAULT	NUTS Lvl 3		Primary
SURVEY_DATE	DEFAULT	Date of observation	yyyy-mm-dd	Harmonized
CAR_LATITUDE	DEFAULT	GPS Car parking latitude	°	Primary
CAR_LONGITUDE	DEFAULT	GPS Car parking longitude	°	Primary
CAR_EW	DEFAULT	GPS Car parking East/West	1: East, 2: West, -1: Not Relevant	Primary
GPS_PROJ	DEFAULT	GPS Projection	1: WGS84, 2: GPS Problem, -1: Not Relevant	Harmonized
GPS_PREC	DEFAULT	GPS Precision	m	Primary
GPS_LAT	DEFAULT	GPS Observation latitude	°	Harmonized

Table A2. Cont.

Attribute	Group	Description	Units	Origin
GPS_EW		GPS Observation East/West	1: East, 2: West, -1: Not Relevant	Harmonized
GPS_LONG	DEFAULT	GPS Observation longitude	°	Harmonized
GPS_ALTITUDE	DEFAULT	GPS altitude	m	Primary
GEOG_GPS	DEFAULT	PostGIS geography (EPSG 4326) generated from GPS_LAT, GPS_LONG		New
GEOM_GPS	DEFAULT	PostGIS geometry (EPSG 3035) generated from GPS_LAT, GPS_LONG		New
GEOM_REPR_AREA	DEFAULT	PostGIS geometry (EPSG 3035) of representative area		New
TH_LAT	DEFAULT	Theoretical Latitude	°	Primary
TH_EW		Theoretical East/West	1: East, 2: West, -1: Not Relevant	Harmonized
TH_LONG	DEFAULT	Theoretical Longitude	°	Primary
GEOG_TH	DEFAULT	PostGIS geography (EPSG 4326) generated from TH_LAT, TH_LONG		New
GEOM_THR	DEFAULT	PostGIS geography (EPSG 3035) generated from TH_LAT, TH_LONG snapped to LUCAS grid		New
GEOM	DEFAULT	PostGIS geometry (EPSG 3035) generated from measured GPS location (GEOM_GPS) if no GPS problem detected otherwise theoretical location (GEOM_THR)		New
DIST_THR_GRID	DEFAULT	Distance computed from GEOG_THR and LUCAS grid	m	New
OBS_DIST	DEFAULT	GPS Distance to theoretical point	m	Harmonized
OBS_DIRECT	DEFAULT	Direction of observation in case of linear feature	1: on the point, 2: Look to the North, 3: Look to the East, -1: Not Relevant	Primary
OBS_TYPE	DEFAULT	Observation type	1: In Situ < 100 m, 2: In Situ > 100 m, 3: In Situ PI, 4: In Situ PI not possible, 5: Out of national territory, 6: Out of EU28, 7: In Office PI, -1: Not Relevant	Harmonized
OBS_RADIUS	DEFAULT	Radius of observation circle	1: 1.5 m, 2: 20 m, -1: Not Relevant	Primary
LC1	LAND COVER (LC_LU, LC_LU_SO)	Land Cover 1		Primary
LC1_H	LAND COVER (LC_LU, LC_LU_SO)	Harmonized Land Cover 1 to 2018 nomenclature	-1: Not Relevant	New
LC1_H_L3_MISSING	LAND COVER (LC_LU, LC_LU_SO)	Harmonized Land Cover 1 on lvl 1 or lvl 2 if lvl 3 is missing		New
LC1_H_L3_MISSING _LEVEL	LAND COVER (LC_LU, LC_LU_SO)	Level of available land cover 1 value if lvl 3 is missing	1: Level 1, 2: Level 2	New
LC1_SPEC	LAND COVER (LC_LU, LC_LU_SO)	Land Cover 1 Species	-1: Not Relevant	Harmonized
LC1_PERC	LAND COVER (LC_LU, LC_LU_SO)	Percentage of coverage of Land Cover 1	%, -1: Not Relevant	Harmonized
LC1_PERC_CLS	LAND COVER (LC_LU, LC_LU_SO)	Percentage of coverage of Land Cover 1 by codes	1: 10%, 2: 25%, 3: 50%, 4: 75%, 5: 100%, -1: Not Relevant	New
LC2	LAND COVER (LC_LU, LC_LU_SO)	Land Cover 2		Primary
LC2_H	LAND COVER (LC_LU, LC_LU_SO)	Harmonized Land Cover 2 to 2018 nomenclature	-1: Not Relevant	New

Table A2. Cont.

Attribute	Group	Description	Units	Origin
LC2_H_L3_MISSING	LAND COVER (LC_LU, LC_LU_SO)	Harmonized Land Cover 2 on lvl 1 or lvl 2 if lvl 3 is missing		New
LC2_H_L3_MISSING_LEVEL	LAND COVER (LC_LU, LC_LU_SO)	Level of available land cover 2 value if lvl 3 is missing	1: Level 1, 2: Level 2	New
LC2_SPEC	LAND COVER (LC_LU, LC_LU_SO)	Land Cover 2 Species	-1: Not Relevant	Harmonized
LC2_PERC	LAND COVER (LC_LU, LC_LU_SO)	Percentage of coverage of Land Cover 2	%, -1: Not Relevant	Harmonized
LC2_PERC_CLS	LAND COVER (LC_LU, LC_LU_SO)	Percentage of coverage of Land Cover 2 by codes	1: 10%, 2: 25%, 3: 50%, 4: 75%, 5: 100%, -1: Not Relevant	New
LU1	LAND USE (LC_LU, LC_LU_SO)	Land Use 1		Primary
LU1_H	LAND USE (LC_LU, LC_LU_SO)	Harmonized Land Use 1 to 2018 nomenclature	-1: Not Relevant	New
LU1_TYPE	LAND USE (LC_LU, LC_LU_SO)	Land Use 1 species	-1: Not Relevant	Primary
LU1_PERC	LAND USE (LC_LU, LC_LU_SO)	Percentage of coverage of Land Use 1	%, -1: Not Relevant	Harmonized
LU1_PERC_CLS	LAND USE (LC_LU, LC_LU_SO)	Percentage of coverage of Land Use 1 by codes	1: 5%, 2: 10%, 3: 25%, 4: 50%, 5: 75%, 6: 90%, 7: 100%, -1: Not Relevant	New
LU2	LAND USE (LC_LU, LC_LU_SO)	Land Use 2		Primary
LU2_H	LAND USE (LC_LU, LC_LU_SO)	Harmonized Land Use 2 to 2018 nomenclature	-1: Not Relevant	New
LU2_TYPE	LAND USE (LC_LU, LC_LU_SO)	Land Use 2 species	-1: Not Relevant	Primary
LU2_PERC	LAND USE (LC_LU, LC_LU_SO)	Percentage of coverage of Land Use 2	%, -1: Not Relevant	Harmonized
LU2_PERC_CLS	LAND USE (LC_LU, LC_LU_SO)	Percentage of coverage of Land Use 2 by codes	1: 5%, 2: 10%, 3: 25%, 4: 50%, 5: 75%, 6: 90%, 7: 100%, -1: Not Relevant	New
PARCEL_AREA_HA	LAND USE (LC_LU, LC_LU_SO)	Parcel Area - area of the parcel which the point belongs to	1: <0.1 ha, 2: 0.1–0.5 ha, 3: 0.5–1 ha, 4: 1–10 ha, 5: >10 ha, -1: Not Relevant	Harmonized
TREE_HEIGHT_SURVEY	TREE PROPERTIES (FO)	Height of trees at survey time	1: <5 m, 2: >5 m, -1: Not Relevant	Primary
TREE_HEIGHT_MATURITY	TREE PROPERTIES (FO)	Height of trees at maturity	1: <5 m, 2: >5 m, -1: Not Relevant	Primary
FEATURE_WIDTH	LAND COVER (LC_LU, LC_LU_SO)	Feature width	1: <20 m, 2: >20 m, -1: Not Relevant	Primary
LM_PLOUGH_SLOPE	LAND MANAGEMENT (LC_LU, LC_LU_SO)	Slope of ploughed field	1: Flat, 2: Gently sloping, 3: Steeply sloping, 4: Undulating, -1: Not Relevant	Primary
LM_PLOUGH_DIRECT	LAND MANAGEMENT (LC_LU, LC_LU_SO)	Plough direction	1: Across the slope, 2: Down the slope, 3: Not Applicable, -1: Not Relevant	Primary
LM_STONE_WALLS	LAND MANAGEMENT (LC_LU, LC_LU_SO)	Presence of stone walls	1: No, 2: Stone wall not maintained, 3: Stone wall well maintained, -1: Not Relevant	Primary
LM_GRASS_MARGINS	LAND MANAGEMENT (LC_LU, LC_LU_SO)	Presence of grass margins	1: No, 2: Grass margin < 1 m, 3: Grass margin > 1 m, -1: Not Relevant	Primary

Table A2. Cont.

Attribute	Group	Description	Units	Origin
CPRN_CANDO	COPERNICUS LAND COVER (CO)	Copernicus taken	1: Yes, 2: No, -1: Not Relevant	Primary
CPRN_LC	COPERNICUS LAND COVER (CO)	Copernicus Land Cover		Primary
CPRN_LC1N	COPERNICUS LAND COVER (CO)	Extension of LC North		Primary
CPRNC_LC1E	COPERNICUS LAND COVER (CO)	Extension of LC East		Primary
CPRNC_LC1S	COPERNICUS LAND COVER (CO)	Extension of LC South		Primary
CPRNC_LC1W	COPERNICUS LAND COVER (CO)	Extension of LC West		Primary
CPRN_LC1N_BRDTH	COPERNICUS LAND COVER (CO)	Percentage of breadth North	%, -1: Not Relevant	Primary
CPRN_LC1E_BRDTH	COPERNICUS LAND COVER (CO)	Percentage of breadth East	%, -1: Not Relevant	Primary
CPRN_LC1S_BRDTH	COPERNICUS LAND COVER (CO)	Percentage of breadth South	%, -1: Not Relevant	Primary
CPRN_LC1W_BRDTH	COPERNICUS LAND COVER (CO)	Percentage of breadth West	%, -1: Not Relevant	Primary
CPRN_LC1N_NEXT	COPERNICUS LAND COVER (CO)	Next copernicus Land Cover North		Primary
CPRN_LC1E_NEXT	COPERNICUS LAND COVER (CO)	Next copernicus Land Cover East		Primary
CPRN_LC1S_NEXT	COPERNICUS LAND COVER (CO)	Next copernicus Land Cover South		Primary
CPRN_LC1W_NEXT	COPERNICUS LAND COVER (CO)	Next copernicus Land Cover West		Primary
CPRN_URBAN	URBAN (CO)	Point in Urban area	1: Yes, 2: No, -1: Not Relevant	Primary
CPRN_IMPERVIOUS_PERC	IMPERVIOUS (CO)	Percentage of imperviousness	%, -1: Not Relevant	Primary
INSPIRE_PLCC1	INSPIRE PLCC (IN)	Percentage of Coniferous forest trees	%, -1: Not Relevant	Primary
INSPIRE_PLCC2	INSPIRE PLCC (IN)	Percentage of Broadleaved forest trees	%, -1: Not Relevant	Primary
INSPIRE_PLCC3	INSPIRE PLCC (IN)	Percentage of Shrubs	%, -1: Not Relevant	Primary
INSPIRE_PLCC4	INSPIRE PLCC (IN)	Percentage of herbaceous plants	%, -1: Not Relevant	Primary
INSPIRE_PLCC5	INSPIRE PLCC (IN)	Percentage of Lichens and mosses	%, -1: Not Relevant	Primary
INSPIRE_PLCC6	INSPIRE PLCC (IN)	Percentage of consolidated bare land	%, -1: Not Relevant	Primary
INSPIRE_PLCC7	INSPIRE PLCC (IN)	Percentage of unconsolidated bare land	%, -1: Not Relevant	Primary
INSPIRE_PLCC8	INSPIRE PLCC (IN)	Percentage of other land	%, -1: Not Relevant	Primary
EUNIS_COMPLEX	EUNIS (LC_LU)	EUNIS Complex	6: X06, 9: X09, 10: Other, 11: Unknown, -1: Not Relevant	Primary
GRASSLAND_SAMPLE	GRASS (LC_LU)	Sample Grassland module	0: FALSE, 1: TRUE	Primary
GRASS_CANDO	GRASS (LC_LU)	Grassland taken	1: Yes, 2: No, -1: Not Relevant	Primary
GRAZING	LAND USE (LC_LU, LC_LU_SO)	Signs of grazing	1: Visible sighns of grazing, 2: No sighn of grazing, -1: Not Relevant	Harmonized
WM	LAND USE (LC_LU, LC_LU_SO)	Presence of Water Management	1: Irrigation, 2: Potential irrigation, 3: Drainage, 4: Irrigation and drainage, 5: No visible Water management, -1: Not Relevant	Primary

Table A2. Cont.

Attribute	Group	Description	Units	Origin
WM_SOURCE	LAND USE (LC_LU, LC_LU_SO)	Source of irrigation	1: Well, 2: Pond/Lake/Reservoir, 3: Stream/Canal/Ditch, 4: Lagoon/Wastewater, 5: Other/Not identifiable, -1: Not Relevant	Harmonized
WM_TYPE	LAND USE (LC_LU, LC_LU_SO)	Type of irrigation	1: Gravity, 2: Pressure sprinkler irrigation, 3: Pressure micro-irrigation, 4: Gravity/Pressure, 5: Other/Not identifiable, -1: Not Relevant	Harmonized
WM_DELIVERY	LAND USE (LC_LU, LC_LU_SO)	Delivery System	1: Canal, 2: Ditch, 3: Pipeline, 4: Other/Not identifiable, -1: Not Relevant	Harmonized
SOIL_TAKEN	SOIL (LC_LU_SO)	Soil taken	1: Yes, 2: Not possible, 3: No, already taken, 4: No sample required, -1: Not Relevant	Harmonized
EROSION_CANDO	SOIL (LC_LU_SO)	Erosion taken	1: Yes, 2: No, -1: Not Relevant	Primary
BIO_SAMPLE	SOIL (LC_LU_SO)	Sample bio soil module	0: FALSE, 1: TRUE	Primary
SOIL_BIO_TAKEN	SOIL (LC_LU_SO)	Bio soil taken	0: FALSE, 1: TRUE, -1: Not Relevant	Primary
BULK0_10_SAMPLE	SOIL (LC_LU_SO)	Sample bulk 0–10 module	0: FALSE, 1: TRUE	Primary
SOIL_BLK_0_10_TAKEN	SOIL (LC_LU_SO)	Bulk 0–10 taken	1: Yes, 2: No, -1: Not Relevant	Primary
BULK10_20_SAMPLE	SOIL (LC_LU_SO)	Sample bulk 10–20 module	0: FALSE, 1: TRUE	Primary
SOIL_BLK_10_20_TAKEN	SOIL (LC_LU_SO)	Bulk 10–20 taken	1: Yes, 2: No, -1: Not Relevant	Primary
BULK20_30_SAMPLE	SOIL (LC_LU_SO)	Sample bulk 20–30 module	0: FALSE, 1: TRUE	Primary
SOIL_BLK_20_30_TAKEN	SOIL (LC_LU_SO)	Bulk 20–30 taken	1: Yes, 2: No, -1: Not Relevant	Primary
STANDARD_SAMPLE	SOIL (LC_LU_SO)	Sample standard soil module	0: FALSE, 1: TRUE	Primary
SOIL_STD_TAKEN	SOIL (LC_LU_SO)	Standard soil taken	1: Yes, 2: No, -1: Not Relevant	Primary
ORGANIC_SAMPLE	SOIL (LC_LU_SO)	Sample organic soil module	0: FALSE, 1: TRUE	Primary
SOIL_ORG_DEPTH_CANDO	SOIL (LC_LU_SO)	Organic soil taken	1: Yes, 2: No, -1: Not Relevant	Primary
OFFICE_PI	DEFAULT	Sample photo interpreted in office	0: FALSE, 1: TRUE	Harmonized
PI_EXTENSION	DEFAULT	Point on extened part of survey (photo-interpreted)	0: FALSE, 1: TRUE	Primary
LNDMNG_PLOUGH	LAND USE (LC_LU, LC_LU_SO)	Signs of ploughing	1: Yes, 2: No, -1: Not Relevant	Primary
SPECIAL_STATUS	LAND USE (LC_LU, LC_LU_SO)	Special status	1: Protected, 2: Hunting, 3: Protected and hunting, 4: No special status, -1: Not Relevant	Primary
LC_LU_SPECIAL_REMARK	LAND COVER (LC_LU, LC_LU_SO)	Special remarks in LC/LU	1: Harvested field, 2: Tilled/sowed, 3: Clear cut, 4: Burnt area, 5: Fire break, 6: Nursey, 7: Dump site, 8: Temporary dry, 9: Temporary flooded, 10: No remark, -1: Not Relevant	Harmonized
SOIL_STONES_PERC	SOIL (LC_LU_SO)	Percentage of Stones on the surface	%, -1: Not Relevant	Harmonized
SOIL_STONES_PERC_CLS	SOIL (LC_LU_SO)	Percentage of Stones on the surface by codes	1: 5%, 2: 20%, 3: 40%, 4: 75%, -1: Not Relevant	New

Table A2. Cont.

Attribute	Group	Description	Units	Origin
PHOTO_POINT	LAND COVER (LC_LU, LC_LU_SO)	Photo point taken	1: Taken, 2: Not Taken, -1: Not Relevant	Primary
PHOTO_NORTH	LAND COVER (LC_LU, LC_LU_SO)	Photo north taken	1: Taken, 2: Not Taken, -1: Not Relevant	Primary
PHOTO_EAST	LAND COVER (LC_LU, LC_LU_SO)	Photo east taken	1: Taken, 2: Not Taken, -1: Not Relevant	Primary
PHOTO_SOUTH	LAND COVER (LC_LU, LC_LU_SO)	Photo south taken	1: Taken, 2: Not Taken, -1: Not Relevant	Primary
PHOTO_WEST	LAND COVER (LC_LU, LC_LU_SO)	Photo west taken	1: Taken, 2: Not Taken, -1: Not Relevant	Primary
CROP_RESIDUES	LAND COVER (LC_LU, LC_LU_SO)	Presence of crop residues	1: Yes, 2: No, -1: Not Relevant	Harmonized
TRANSECT	LAND COVER (LC_LU, LC_LU_SO)	Transect LC sequence		Primary
EX_ANTE	DEFAULT	Visited in the field	0: FALSE, 1: TRUE	Primary
SURVEY_YEAR	DEFAULT	Survey year		New
SURVEY_COUNT	SPACE-TIME	Number of visits		New
SURVEY_DIST	SPACE-TIME	Distance computed from representative location (GEOM) and measured GPS location (GEOM_GPS)	m	New
SURVEY_MAXDIST	SPACE-TIME	Maximum distance computed from representative location (GEOM) and measured GPS location (GEOM_GPS)	m	New

## References

- Akitsu, T.K.; Nasahara, K.N. In-Situ observations on a moderate resolution scale for validation of the Global Change Observation Mission-Climate ecological products: The uncertainty quantification in ecological reference data. *Int. J. Appl. Earth Obs. Geoinf.* **2022**, *107*, 102639. [\[CrossRef\]](#)
- Lee, J.G.; Kang, M. Geospatial Big Data: Challenges and Opportunities. *Big Data Res.* **2015**, *2*, 74–81. [\[CrossRef\]](#)
- Ishwarappa; Anuradha, J. A Brief Introduction on Big Data 5Vs Characteristics and Hadoop Technology. *Procedia Comput. Sci.* **2015**, *48*, 319–324. [\[CrossRef\]](#)
- Koubarakis, M.; Stamoulis, G.; Bilidas, D.; Ioannidis, T.; Pantazi, D.A.; Vlassov, V.; Payberah, A.H.; Wang, T.; Sheikholeslami, S.; Hagos, D.H.; et al. Artificial Intelligence and big data technologies for Copernicus data: The EXTREMEEARTH project. In Proceedings of the 2021 Conference on Big Data from Space, Virtual Event, 18–20 May 2021; pp. 9–12. [\[CrossRef\]](#)
- Overview—Land Cover/Use Statistics. Available online: <https://ec.europa.eu/eurostat/web/lucas> (accessed on 11 April 2022).
- Bettio, M.; Delincé, J.; Bruyas, P.; Croi, W.; Eiden, G. Area frame surveys: Aim, Principals and Operational Surveys. In *Building Agri-Environmental Indicators, Focussing on the European Area Frame Survey LUCAS*; Eurostat: Luxembourg, 2002; pp. 12–27. [\[CrossRef\]](#)
- Fritz, S.; McCallum, I.; Schill, C.; Perger, C.; Grillmayer, R.; Achard, F.; Kraxner, F.; Obersteiner, M. Geo-Wiki.Org: The Use of Crowdsourcing to Improve Global Land Cover. *Remote Sens.* **2009**, *1*, 345–354. [\[CrossRef\]](#)
- Defourny, P.; Mayaux, P.; Herold, M.; Bontemps, S. Global land-cover map validation experiences: Toward the characterization of quantitative uncertainty. In *Remote Sensing of Land Use and Land Cover*; Giri, C.P., Ed.; CRC Press: Boca Raton, FL, USA, 2016; pp. 207–223. [\[CrossRef\]](#)
- Close, O.; Benjamin, B.; Petit, S.; Fripiat, X.; Hallot, E. Use of Sentinel-2 and LUCAS Database for the Inventory of Land Use, Land Use Change, and Forestry in Wallonia, Belgium. *Land* **2018**, *7*, 154. [\[CrossRef\]](#)
- Weigand, M.; Staab, J.; Wurm, M.; Taubenböck, H. Spatial and semantic effects of LUCAS samples on fully automated land use/land cover classification in high-resolution Sentinel-2 data. *Int. J. Appl. Earth Obs. Geoinf.* **2020**, *88*, 102065. [\[CrossRef\]](#)
- Pflugmacher, D.; Rabe, A.; Peters, M.; Hostert, P. Mapping pan-European land cover using Landsat spectral-temporal metrics and the European LUCAS survey. *Remote Sens. Environ.* **2019**, *221*, 583–595. [\[CrossRef\]](#)
- Gao, Y.; Liu, L.; Zhang, X.; Chen, X.; Mi, J.; Xie, S. Consistency Analysis and Accuracy Assessment of Three Global 30-m Land-Cover Products over the European Union using the LUCAS Dataset. *Remote Sens.* **2020**, *12*, 3479. [\[CrossRef\]](#)
- d’Andrimont, R.; Yordanov, M.; Martinez-Sanchez, L.; Eiselt, B.; Palmieri, A.; Dominici, P.; Gallego, J.; Reuter, H.I.; Joebges, C.; Lemoine, G.; et al. Harmonised LUCAS In-Situ land cover and use database for field surveys from 2006 to 2018 in the European Union. *Sci. Data* **2020**, *7*, 352. [\[CrossRef\]](#)

14. Borrelli, P.; Poesen, J.; Vanmaercke, M.; Ballabio, C.; Hervás, J.; Maerker, M.; Scarpa, S.; Panagos, P. Monitoring gully erosion in the European Union: A novel approach based on the Land Use/Cover Area frame survey (LUCAS). *Int. Soil Water Conserv. Res.* **2022**, *10*, 17–28. [CrossRef]
15. Jeppesen, J.H.; Ebeid, E.; Jacobsen, R.H.; Toftegaard, T.S. Open geospatial infrastructure for data management and analytics in interdisciplinary research. *Comput. Electron. Agric.* **2018**, *145*, 130–141. [CrossRef]
16. Wiemann, S.; Brauner, J.; Karrasch, P.; Henzen, D.; Bernard, L. Design and prototype of an interoperable online air quality information system. *Environ. Model. Softw.* **2016**, *79*, 354–366. [CrossRef]
17. Li, W.; Wang, S.; Bhatia, V. PolarHub: A large-scale web crawling engine for OGC service discovery in cyberinfrastructure. *Comput. Environ. Urban Syst.* **2016**, *59*, 195–207. [CrossRef]
18. Klug, H.; Knoch, A. A SMART groundwater portal: An OGC web services orchestration framework for hydrology to improve data access and visualisation in New Zealand. *Comput. Geosci.* **2014**, *69*, 78–86. [CrossRef]
19. Best, B.D.; Halpin, P.N.; Fujioka, E.; Read, A.J.; Qian, S.S.; Hazen, L.J.; Schick, R.S. Geospatial web services within a scientific workflow: Predicting marine mammal habitats in a dynamic environment. *Ecol. Inform.* **2007**, *2*, 210–223. [CrossRef]
20. Rosatti, G.; Zorzi, N.; Zugliani, D.; Piffer, S.; Rizzi, A. A Web Service ecosystem for high-quality, cost-effective debris-flow hazard assessment. *Environ. Model. Softw.* **2018**, *100*, 33–47. [CrossRef]
21. Rautenbach, V.; Coetzee, S.; Iwaniak, A. Orchestrating OGC web services to produce thematic maps in a spatial information infrastructure. *Comput. Environ. Urban Syst.* **2013**, *37*, 107–120. [CrossRef]
22. Web Feature Service | OGC. Available online: <https://www.ogc.org/standards/wfs> (accessed on 11 April 2022).
23. OGC API—Features. Available online: <https://ogcapi.ogc.org/features/> (accessed on 11 April 2022).
24. Giuliani, G.; Ray, N.; Lehmann, A. Grid-enabled Spatial Data Infrastructure for environmental sciences: Challenges and opportunities. *Future Gener. Comput. Syst.* **2011**, *27*, 292–303. [CrossRef]
25. Blauth, D.A.; Ducati, J.R. A Web-based system for vineyards management, relating inventory data, vectors and images. *Comput. Electron. Agric.* **2010**, *71*, 182–188. [CrossRef]
26. Zioti, F.; Ferreira, K.R.; Queiroz, G.R.; Neves, A.K.; Carlos, F.M.; Souza, F.C.; Santos, L.A.; Simoes, R.E.O. A platform for land use and land cover data integration and trajectory analysis. *Int. J. Appl. Earth Obs. Geoinf.* **2022**, *106*, 102655. [CrossRef]
27. Data—Land Cover/Use Statistics. Available online: <https://ec.europa.eu/eurostat/web/lucas/data> (accessed on 11 April 2022).
28. Eurostat Regional Yearbook 2021. Available online: <https://ec.europa.eu/statistical-atlas/viewer/> (accessed on 11 April 2022).
29. Witjes, M.; Parente, L.; van Diemen, C.; Hengl, T.; Landa, M.; Brodsky, L.; Halounova, L.; Krizan, J.; Antonic, L.; Ilie, C.; et al. A spatiotemporal ensemble machine learning framework for generating land use/land cover time-series maps for Europe (2000–2019) based on LUCAS, CORINE and GLAD Landsat. *PeerJ-Life Environ.* **2022**, *accepted*. [CrossRef]
30. LUCAS Primary Data 2006. Available online: <https://ec.europa.eu/eurostat/en/web/lucas/data/primary-data/2006> (accessed on 8 April 2022).
31. LUCAS Primary Data 2009. Available online: <https://ec.europa.eu/eurostat/en/web/lucas/data/primary-data/2009> (accessed on 8 April 2022).
32. LUCAS Primary Data 2012. Available online: <https://ec.europa.eu/eurostat/en/web/lucas/data/primary-data/2012> (accessed on 8 April 2022).
33. LUCAS Primary Data 2015. Available online: <https://ec.europa.eu/eurostat/en/web/lucas/data/primary-data/2015> (accessed on 8 April 2022).
34. LUCAS Primary Data 2018. Available online: <https://ec.europa.eu/eurostat/en/web/lucas/data/primary-data/2018> (accessed on 8 April 2022).
35. LUCAS Grid—Land Cover/Use Statistics. Available online: <https://ec.europa.eu/eurostat/web/lucas/data/lucas-grid> (accessed on 8 April 2022).
36. LUCAS 2009. Technical Reference Document C3 Classification (Land Cover & Land Use). Available online: [https://ec.europa.eu/eurostat/documents/205002/208938/LUCAS2009\\_C3-Classification\\_20121004.pdf](https://ec.europa.eu/eurostat/documents/205002/208938/LUCAS2009_C3-Classification_20121004.pdf) (accessed on 27 May 2020).
37. LUCAS 2012. Technical Reference Document C3 Classification (Land Cover & Land Use). Available online: [https://ec.europa.eu/eurostat/documents/205002/208012/LUCAS\\_2012\\_C3-Classification\\_20131004\\_0.pdf](https://ec.europa.eu/eurostat/documents/205002/208012/LUCAS_2012_C3-Classification_20131004_0.pdf) (accessed on 27 May 2020).
38. LUCAS 2015. Technical Reference Document C3 Classification (Land Cover & Land Use). Available online: [https://ec.europa.eu/eurostat/documents/205002/6786255/LUCAS2015\\_C3-Classification\\_20160729.pdf](https://ec.europa.eu/eurostat/documents/205002/6786255/LUCAS2015_C3-Classification_20160729.pdf) (accessed on 27 May 2020).
39. LUCAS 2018. Technical Reference Document C3 Classification (Land Cover & Land Use). Available online: <https://ec.europa.eu/eurostat/documents/205002/8072634/LUCAS2018-C3-Classification.pdf> (accessed on 27 May 2020).
40. Contents of the 2006 Lucas Primary Data. Available online: [https://ec.europa.eu/eurostat/documents/205002/209869/Contents\\_LUCAS\\_2006\\_primary\\_data.xls](https://ec.europa.eu/eurostat/documents/205002/209869/Contents_LUCAS_2006_primary_data.xls) (accessed on 27 May 2020).
41. LUCAS Survey 2009 Technical Reference Document c-1: Instructions for Surveyors. Available online: <https://ec.europa.eu/eurostat/documents/205002/208938/LUCAS+2009+Instructions> (accessed on 27 May 2020).
42. LUCAS Survey 2012 Technical Reference Document c-1: Instructions for Surveyors. Available online: [https://ec.europa.eu/eurostat/documents/205002/208012/LUCAS2012\\_C1-InstructionsRevised\\_20130110b.pdf](https://ec.europa.eu/eurostat/documents/205002/208012/LUCAS2012_C1-InstructionsRevised_20130110b.pdf) (accessed on 27 May 2020).
43. LUCAS Survey 2015 Web CSV Record Descriptor. Available online: [https://ec.europa.eu/eurostat/documents/205002/6786255/WebCsv\\_RecordDescriptor20161006.pdf](https://ec.europa.eu/eurostat/documents/205002/6786255/WebCsv_RecordDescriptor20161006.pdf) (accessed on 27 May 2020).

44. LUCAS Survey 2018 Web CSV Record Descriptor. Available online: <https://ec.europa.eu/eurostat/documents/205002/8072634/LUCAS2018-RecordDescriptor-190611.pdf> (accessed on 27 May 2020).
45. PostGIS Documentation—ST\_GeometricMedian. Available online: [https://postgis.net/docs/ST\\_GeometricMedian.html](https://postgis.net/docs/ST_GeometricMedian.html) (accessed on 11 April 2022).
46. Weiszfeld, E.; Plastria, F. On the point for which the sum of the distances to n given points is minimum. *Ann. Oper. Res.* **2009**, *167*, 7–41. [[CrossRef](#)]
47. Vale, T.; Crnkovic, I.; de Almeida, E.S.; da Mota Silveira Neto, P.A.; Cavalcanti, Y.C.; de Lemos Meira, S.R. Twenty-eight years of component-based software engineering. *J. Syst. Softw.* **2016**, *111*, 128–148. [[CrossRef](#)]
48. Nierstrasz, O.; Meijler, T.D. Research Directions in Software Composition. *ACM Comput. Surv.* **1995**, *27*, 262–264. [[CrossRef](#)]
49. Pytest: Helps You Write Better Programs—Pytest Documentation. Available online: <https://docs.pytest.org/en/7.1.x/> (accessed on 13 April 2022).
50. Merkel, D. Docker: Lightweight Linux Containers for Consistent Development and Deployment. *Linux J.* **2014**, *2014*, 2. Available online: <https://dl.acm.org/doi/10.5555/2600239.2600241> (accessed on 13 April 2022).
51. PostgreSQL: The World’s Most Advanced Open-Source Database. Available online: <https://www.postgresql.org/> (accessed on 13 April 2022).
52. PostGIS Documentation. Available online: <https://postgis.net/> (accessed on 13 April 2022).
53. GDAL Documentation. Available online: <https://gdal.org/> (accessed on 13 April 2022).
54. GeoServer Documentation. Available online: <https://geoserver.org/> (accessed on 13 April 2022).
55. GeoPandas Documentation. Available online: <https://geopandas.org/en/stable/> (accessed on 13 April 2022).
56. Perkel, J.M. BY Jupyter, it all makes sense. *Nature* **2018**, *563*, 145–146. [[CrossRef](#)]
57. Reference Data—GISCO. Available online: <https://gisco-services.ec.europa.eu/lucas/photos/> (accessed on 11 April 2022).
58. Land Parcel Identification System—LPIS. Available online: <https://eagri.cz/public/app/lpisext/lpis/verejny2/plpis/> (accessed on 1 February 2022).