

Table S1. Primer sequences used for qRT-PCR analysis.

| Primer name | Primer sequence | Reference |
|-----------------|--------------------------|----------------------|
| NRT2.1-F | GCTGCTTGTGGTGCTGTAT | Wang et al., 2020 |
| NRT2.1-R | GAGCGACGGGTAATGTGC | |
| NRT2.2-F | CCTCACTAAGCGTGACATCAGC | Wang et al., 2020 |
| NRT2.2-R | CGAGGAGGTCGCACACCAC | |
| NRT2.3-F | GCTGGGTCTTCGTGCTCCT | Wang et al., 2020 |
| NRT2.3-R | GGCGATGACATTGTCCGGTG | |
| NRT2.4-F | CTGCCCCGTCGCCTTCAT | Wang et al., 2020 |
| NRT2.4-R | CGGTCCACTCCCCGTTGTAG | |
| NRT2.5-F | CGGTGGACTCCGACAACAA | Wang et al., 2020 |
| NRT2.5-R | GAGAACCACGAGAGGTGGAAG | |
| NRT2.6-F | TTCGGCATCGTTCCTTC | Wang et al., 2020 |
| NRT2.6-R | TCCTCCACCGTGTACCTGG | |
| NAR2.1-F | CACGGTCGGAGCTGGTCA | Wang et al., 2020 |
| NAR2.1-R | TGGCTCGCGTTCAGGG | |
| NAR2.2-F | GGCGGCAGGGTCGAGTA | Wang et al., 2020 |
| NAR2.2-R | GGTGACGCCCGTGATGC | |
| NAR2.3-F | GGCTCGCCAAGGCATGGTC | Wang et al., 2020 |
| NAR2.3-R | AGGTCGTCTGGGGTCTTGC | |
| NPF2.1-F | CGGTGGTACAGGTTCAAGAAGTC | Buchner et al., 2014 |
| NPF2.1-R | CGGAGTATAKGAAGCCCCAAAAG | |
| NPF2.2-F | GTGGACCTRTTCTACCTTGTAC | Buchner et al., 2014 |
| NPF2.2-R | AGCCTTCTTGGGGCTTTCATCG | |
| NPF6.1-F | CAATCGGACGGCCTTGATTTCTTC | Buchner et al., 2014 |
| NPF6.1-R | ACGTAGGCACACGTGGACGAC | |
| NPF6.2-F | ATCGATGCATGCTACTTGCSTTC | Buchner et al., 2014 |
| NPF6.2-R | GCCACCGAATACACAAAAACAAG | |
| NPF7.1-F | CTACAAGACCTGCGCCATCTTC | He et al., 2015 |
| NPF7.1-R | GATGAGGTATAGCCGCGAGGAG | |
| NPF7.2-F | CCCAGCAGTCAAAGCAAACACTG | He et al., 2015 |
| NPF7.2-R | GGAGGAACACCACCAGGTTAC | |
| AMT6-F | GGACTACCTGTGCAACCAGTTC | Bajgain et al., 2018 |
| AMT6-R | CGAAGCCGAAGAGGTAGTAGAA | |
| AMT49-F | ATGGCTAACACAGAAGCATCAA | Bajgain et al., 2018 |
| AMT49-R | TTTCTTTCGTGGCCTCTACTTC | |
| AMT52-F | CAGTATGGCAATGGAGAGTTCA | Bajgain et al., 2018 |
| AMT52-R | CACCGATATATGCATTTTGTGC | |
| EF- α -F | GGTTAAGATGATTCCCACCAAGCC | Wang et al., 2020 |
| EF- α -R | GACAACACCAACAGCAACAGTCTG | |
| Actin3 like-F | GACGCACAACAGGTATCGTGTTG | Buchner et al., 2014 |
| Actin3 like-R | AGCGAGGTCAAGACGAAGGATG | |

References

- Bajgain, P.; Russell, B.; Mohammadi, M. Phylogenetic analyses and in-seedling expression of ammonium and nitrate transporters in wheat. *Sci. Rep.* **2018**, *8*, 7082.
- Buchner, P.; Hawkesford, M.J. Complex phylogeny and gene expression patterns of members of the NITRATE TRANSPORTER 1/PEPTIDE TRANSPORTER family (NPF) in wheat. *J. Exp. Bot.* **2017**, *65*, 5697-5710.
- He, X.; Qu, B.; Li, W.; Zhao, X.; Teng, W.; Ma, W.; Ren, Y.; Li, B.; Li, Z.; Tong, Y. The nitrate-inducible NAC transcription factor TaNAC2-5A controls nitrate response and increases wheat yield. *Plant Physiol.* **2015**, *169*, 1991-2005.
- Wang, M.; Zhang, P.; Liu, Q.; Li, G.; Di, D.; Xia, G.; Kronzucker, H.J.; Fang, S.; Chu, J.; Shi, W. TaANR1-TaBG1 and TaWabi5-TaNRT2s/NARs link ABA metabolism and nitrate acquisition in wheat roots. *Plant Physiol.* **2020**, *182*, 1440-1453.