

Biosynthesized Silicon Dioxide Nanoparticles Mitigate Wheat Damping-Off, Caused by *Rhizoctonia solani*, via the Simultaneous Activation of a Multilayered Defense System

Abdelrazek S. Abdelrhim ¹, Yasser S. A. Mazrou ^{2,3}, Yasser Nehela ^{4,5,*}, Osama O. Atallah ⁶, Ranya M. El-Ashmoni ¹ and Mona F. A. Dawood ⁷

¹ Department of Plant Pathology, Faculty of Agriculture, Minia University, El-Minya 61512, Egypt; Abdelrazek.sharawy@mu.edu.eg (A.S.A); ranya.elashmoni@mu.edu.eg (R.M.E.-A.)

² Business Administration Department, Community college, King Khalid University, Guraiger, Abha 62529, Saudi Arabia; ymazrou@kku.edu.sa

³ Department of Agriculture Economic, Faculty of Agriculture, Tanta University, Tanta 31527, Egypt; yasser.mazroua@agr.tanta.edu.eg

⁴ Department of Agricultural Botany, Faculty of Agriculture, Tanta University, Tanta 31511, Egypt; yasser.nehela@ufl.edu

⁵ Department of Plant Pathology, Citrus Research and Education Center, University of Florida, 700 Experiment Station Rd., Lake Alfred, FL 33850, USA

⁶ Department of Plant Pathology, Zagazig University, Zagazig 44519, Egypt; osamaoatall1h@ufl.edu

⁷ Botany and Microbiology Department, Faculty of Science, Assiut University, Assiut 71516, Egypt; mo_fa87@aun.edu.eg

* Correspondence: yasser.nehela@ufl.edu




Figure S1. Energy-dispersive X-ray spectroscopy (EDS) elemental mapping of Si K α 1 (**A**) and O K α 1 (**B**).