

Supplementary Material

Involvement of Polyamine Metabolism in the Response of *Medicago truncatula* Genotypes to Salt Stress

Chrystalla Antoniou ¹, Xavier Zarza ², Gholamreza Gohari ³, Sima Panahirad ⁴, Panagiota Filippou ^{1,†}, Antonio F. Tiburcio ² and Vasileios Fotopoulos ^{1,*}

Table 1. Oligonucleotides used as primers for real-time RT-qPCR.

Gene	Primer	Nucleotide sequence	T _a (°C)	Reference
<i>Act11</i>	Actin11-F	5'-ATG CCA TCC TTC GTC TTG A-3'	53	[Filippou et al., 2013]
	Actin11-R	5'-GCT GGT CCT GGC TGT CTC-3'		
<i>ADC</i>	ADC-F	5'-CTG GCC ATT TTG GTT CAA CT-3'	56	[Filippou et al., 2013]
	ADC-R	5'-ACC CAA ACG AAG CAA TTC AC- 3'		
<i>SPDS</i>	SPDS-F	5'-AAG GGA TGA GTG TGC GTA CC-	56	[Filippou et al., 2013]
	SPDS-R	3' 5'-TTT GGA GAC ATC GAC AAC CA- 3'		
<i>SPMS</i>	SPMS-F	5'-GCC AGT GAA GAA AAG GGT CA-3'	60	[Filippou et al., 2013]
	SPMS-R	5'-AGA ACC ACC CAG AAA CAA CG-3'		
<i>PAO</i>	PAO-F	5'-TTT TGG CAG CAC ATG GAT AA- 3'	60	[Filippou et al., 2013]
	PAO-R	5'-TTA TTC CAC CAG CAG GGA AC- 3'		
<i>DAO</i>	DAO-F	5'-TGC AAT CCC AGA TGA AGT GA- 3'	60	[Filippou et al., 2013]
	DAO-R	5'-CAG CTA GCA ATG TGC CAT GT- 3'		
<i>SAMD</i> <i>C</i>	SAMD-C-F	5'-CAA CGG TGG CGT AGA AAA AT- 3'	56	[Filippou et al., 2013]
	SAMD-C-R	5'-GCC TTC AAA ACC GAT AGC TG - 3'		

References

Filippou, P.; Antoniou, C.; Fotopoulos, V. The nitric oxide donor sodium nitroprusside regulates proline and polyamine biosynthesis in *Medicago truncatula* plants. *Free Rad. Biol. Med.* **2013**, *56*, 172–183