

SUPPLEMENTARY MATERIAL

Triterpenoids from *Vitellaria paradoxa* Stem Barks Reduce Nitrite Levels in LPS-Stimulated Macrophages

Carmina Sirignano ¹, Pascal Nadembega ², Ferruccio Poli ³, Barbara Romano ¹,
Giuseppe Lucariello ¹, Daniela Rigano ^{1,*}, and Orazio Taglialatela-Scafati ¹

¹ Department of Pharmacy, School of Medicine and Surgery, University of Naples Federico II, Via Domenico Montesano 49, 80131 Naples, Italy.; carmina.sirignano@unina.it (C.S.); barbara.romano@unina.it (B.R.); giuseppe.lucariello@unina.it (G.L.); drigano@unina.it (D.R.); scatagli@unina.it (O.T.S.)

² University of Ouagadougou UFR/SVT, 03 BP, 848 Ouagadougou 03, Burkina Faso; pascal.nadembega@gmail.com

³ Department of Pharmacy and Biotechnology, University of Bologna, Via Imerio, 42, 40126, Bologna, Italy; ferruccio.poli@unibo.it




Figure S1. Negative ion MS of lupeol (**1**)




Figure S2. Positive ion MS of lupeol acetate (**2**)




Figure S3. Positive ion MS of lupeol cinnamate (3)




Figure S4. Negative ion MS of lupenone (4)




Figure S5. Negative ion MS of betulinic acid (5)




Figure S6. Positive ion MS of α -amyrin cinnamate (6)




Figure S7. Positive ion MS of ursaldehyde cinnamate (7)




Figure S8. Negative ion MS of ursolic acid (8)




Figure S9. Positive ion MS of β -amyrin acetate (**9**)




Figure S10. Negative ion MS of 11-hydroxy- β -amyrin cinnamate (**10**)