

Review

Mechanistic Insight into Antimicrobial and Antioxidant Potential of *Jasminum* Species: A Herbal Approach for Disease Management

Acharya Balkrishna ¹, Akansha Rohela ¹, Abhishek Kumar ¹, Ashwani Kumar ^{1,*}, Vedpriya Arya ¹, Pallavi Thakur ¹, Patrik Oleksak ², Ondrej Krejcar ³, Rachna Verma ^{4,*}, Dinesh Kumar ⁵ and Kamil Kuca ^{2,6,*}

- ¹ Patanjali Herbal Research Department, Patanjali Research Institute, Haridwar 249405, India; acharya.balkrishnapri@prft.in (A.B.); akansha.rohela@prft.co.in (A.R.); abhishekkumar@prft.co.in (A.K.); vedpriya.arya@prft.in (V.A.); pallavi.thakur@prft.co.in (P.T.)
- ² Department of Chemistry, Faculty of Science, University of Hradec Kralove, 50003 Hradec Kralove, Czech Republic; patrik.oleksak@uhk.cz
- ³ Center for Basic and Applied Science, Faculty of Informatics and Management, University of Hradec Kralove, 50003 Hradec Kralove, Czech Republic; ondrej.krejcar@uhk.cz
- ⁴ School of Biological and Environmental Sciences, Shoolini University of Biotechnology and Management Sciences, Solan 173229, India
- ⁵ School of Bioengineering and Food Technology, Shoolini University of Biotechnology and Management Sciences, Solan 173229, India; chatantadk@yahoo.com
- ⁶ Biomedical Research Center, University Hospital in Hradec Kralove, Sokolska 581, 50005 Hradec Kralove, Czech Republic
- * Correspondence: dr.ashwanikumar@prft.co.in (A.K.); rachnac83@gmail.com (R.V.); kamil.kuca@uhk.cz (K.K.); Tel.: +420-603-289-166 (K.K.)

Citation: Balkrishna, A.; Rohela, A.; Kumar, A.; Kumar, A.; Arya, V.; Thakur, P.; Oleksak, P.; Krejcar, O.; Verma, R.; Kumar, D.; et al. Mechanistic Insight into Antimicrobial and Antioxidant Potential of *Jasminum* Species: A Herbal Approach for Disease Management. *Plants* **2021**, *10*, 1089. <https://doi.org/10.3390/plants10061089>

Academic Editors: Yun-Soo Seo and Joong-Sun Kim

Received: 25 April 2021

Accepted: 25 May 2021

Published: 28 May 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Drug resistance among microbial pathogens and oxidative stress caused by reactive oxygen species are two of the most challenging global issues. Firstly, drug-resistant pathogens cause several fatalities every year. Secondly aging and a variety of diseases, such as cardiovascular disease and cancer, are associated with free radical generated oxidative stress. The treatments currently available are limited, ineffective, or less efficient, so there is an immediate need to tackle these issues by looking for new therapies to resolve resistance and neutralize the harmful effects of free radicals. In the 21st century, the best way to save humans from them could be by using plants as well as their bioactive constituents. In this specific context, *Jasminum* is a major plant genus that is used in the Ayurvedic system of medicine to treat a variety of ailments. The information in this review was gathered from a variety of sources, including books, websites, and databases such as Science Direct, PubMed, and Google Scholar. In this review, a total of 14 species of *Jasminum* have been found to be efficient and effective against a wide variety of microbial pathogens. In addition, 14 species were found to be active free radical scavengers. The review is also focused on the disorders related to oxidative stress, and it was concluded that *Jasminum grandiflorum* and *J. sambac* normalized various parameters that were elevated by free radical generation. Alkaloids, flavonoids (rutoside), terpenes, phenols, and iridoid glucosides are among the main phytoconstituents found in various *Jasminum* species. Furthermore, this review also provides insight into the mechanistic basis of drug resistance, the generation of free radicals, and the role of *Jasminum* plants in combating resistance and neutralizing free radicals.

Keywords: *Jasminum* species; antimicrobial; antioxidants; mechanistic insight; reactive oxygen species

1. Introduction

Emerging infections and the rise in antibiotic resistance among pathogens have been the major challenges that will endanger society's health today. Worldwide, millions of deaths are attributed to microbial diseases. In 2013, a total of 9.2 million deaths due to microbial infections were reported [1,2]. The occurrence of drug resistance has resulted in a decline in the efficacy and ineffectiveness of existing antibiotics [3,4]. Recently, several studies have mainly focused primarily on seeking promising ways to solve these challenges, looking for new antimicrobials, as well as modification of existing ones. Interestingly, as the safest source to obtain various medicines, medicinal plants are preferred, as per the World Health Organization [5].

Plants and their bioactive compounds could be one of the most promising antimicrobial agents to fight resistance in the current scenario. They are pioneers in the discovery of new medicines, used directly or as a precursor for the synthesis of bioactive constituents. Phytochemicals have exerted antimicrobial action by different mechanisms against a wide variety of pathogens (susceptible and resistant). In plants, significant quantities of secondary metabolites are found, such as tannins, alkaloids, phenolics, and flavonoids [6,7]. A large number of medicinal plants have been shown to be effective in treating bacterial infections [8]. Researchers around the world have already investigated and continue to explore a variety of plants for their antimicrobial activity. For instance, *Osyris auriculata* and *Ficus auriculata* were found to be active towards *Escherichia coli* and *Salmonella typhimurium* [9].

On the other hand, under metabolic conditions, oxygen, an efficient molecule, can create free radicals; both are important and essential in the biochemical process and in redox reactions [10,11]. A number of oxygen free radicals are grouped under reactive oxygen species (ROS). However, ROS are highly unstable, attacking nucleic acids, lipids, and proteins, initiating a chain reaction, and playing a part in the etiology of major human diseases [12,13]. In addition, ROS generation is also associated with cancer, and cardiovascular disorders [14]. In regard to this, plant phytochemicals provide significant protection against free radicals as they neutralize their harmful effects. For instance, *Prunus domestica*, *Syzygium cumini*, *Rubus ellipticus*, and *Prunus armeniaca* fruits were found to be enriched with antioxidants [15]. Interestingly, medicinal plants could be seen as a precious gift from nature to humans [16].

Jasminum in the *Oleaceae* family are an important group of flowering plants, commonly cultivated for their aromatic flowers [17]. There are 197 taxonomically recognized species (spp.) of the genus *Jasminum* in the world [18]. *Jasminum* plants have been recommended against intestinal worms and venereal diseases [19,20]. Notably, in the aspect of medicine and pharmaceutical sectors, almost all parts of the plant are important [17]. Flowers are used for treating vesicles, ulcers, skin diseases, boils, and eye disorders, while leaves are used against breast tumors [20]. Traditionally, many members of the *Jasminum* genus have been used medicinally. For instance, *J. officinale* exhibits various therapeutic properties, viz., depurative, analgesic, diuretic, antiseptic, expectorant, anti-depressant, and sedative [21]. It is used by various tribals for treating gastrointestinal disorders, cough, pyrexia, eye inflammation, and also against irregular menstruation [22]. Likewise, *J. grandiflorum* is recommended against cough, hysteria, uterine ailments, and partum problems [23]. Subsequently, *J. sambac* is expectorant, analgesic, antiseptic, aphrodisiac, anti-depressant, and sedative [24]. Its flowers are taken orally by the Meitei community to treat piles [25]. Furthermore, a decoction of its roots (3g) mixed with honey (10g) is taken twice daily in the treatment of amenorrhoea [26]. This review focuses on evaluating the antimicrobial and antioxidant ability of *Jasminum* spp., keeping in mind the emergence of antibiotic resistance and oxidative stress induced by ROS. The first half of the review sheds light on their antimicrobial potential and mechanistic viewpoint of drug resistance, whereas the second half demonstrates the antioxidant potential, the effectiveness of plants against disorders associated with oxidative stress, and the mechanistic basis for free radical neutralization by *Jasminum* spp.

2. Search Strategy

Different databases like Science Direct, PubMed, and Google Scholar have been explored in this review with various keywords such as *Jasminum* species, antimicrobial, antibacterial, antifungal, antioxidant activity, reactive oxygen species, oxidative stress, phytochemistry, bacterial and antifungal drug resistance. The compilation of literature was carried out between 1 December 2020 and 1 February 2021. In order to maintain quality, only full length, original, and English language papers from Web of Science and Scopus indexed peer-reviewed journals have been included in this study. Further, papers with more than 5 citations are also included in some cases. The current review was compiled based on 100 studies, 34 review papers, 4 website reports, and 5 books that were published between 1971 and 2021.

3. Distribution of Genus *Jasminum*

The native range of the genus *Jasminum* covers tropical, subtropical Old World to Central China and the Pacific region (Figure 1). Further, they have been introduced into Europe, the Caribbean region, South and Central America, and the U.S.A. [18].




Figure 1. Global distribution of genus *Jasminum* (Created with mapchart.net).

4. Antimicrobial Profile of *Jasminum* spp.

A total of 14 *Jasminum* spp. have been documented for their anti-microbial activity against Gram positive and negative bacterial strains, and fungal pathogens. In terms of efficacy against a wide variety of bacterial pathogens and minimal antifungal activity, all of the *Jasminum* plants are extremely encouraging.

A fraction of acetone extract from the leaves of *Jasminum azoricum* has shown anti-*Staphylococcus aureus* activity with the highest inhibition zone (30 mm at 30 mg/mL) among all the 14 species studied, whereas methanolic extract of *Jasminum syringifolium* leaves exhibited a 22.67-mm inhibition zone against *Shigella flexneri* (Table 1). Further, jatamansone extract from leaves of *Jasminum brevilobum* has shown the lowest minimum inhibitory concentration (MIC 0.05 µg/mL) against *Staphylococcus aureus* among all the studied species, whereas, it showed the highest MIC against *Escherichia coli* (MIC 0.07 µg/mL).

Table 1. Antimicrobial potential of *Jasminum* spp.

Botanical Name	Extract/Solvent (Conc.)	Microbes	ZOI (mm)/MIC (µg/mL)	References
<i>Jasminum abyssinicum</i> Hochst. ex DC.	Aerial parts extract/Methanol (250–2000 µg/mL)	<i>Staphylococcus aureus</i>	N.A.	[27]
		<i>Streptococcus pyogenes</i>	N.A.	
		<i>S. pneumonia</i>	N.A.	
		<i>Neisseria gonorrhoea</i>	Active	
		<i>Escherichia coli</i>	N.A.	
		<i>Bacillus cereus</i>	N.A.	
		<i>Shigella dysenteriae</i>	N.A.	
		<i>S. flexneri</i>	N.A.	
		<i>Salmonella typhi</i>	N.A.	
		<i>S. typhimurium</i>	N.A.	
		<i>Aspergillus flavus</i>	N.A.	
		<i>A. niger</i>	N.A.	
		<i>Candida albicans</i>	N.A.	
		<i>Trichophyton mentagrophytes</i>	N.A.	
		<i>T. violaceum</i>	N.A.	
		<i>Cryptococcus neoformans</i>	N.A.	
	Positive control	(Tetracycline, Co-trimoxazole, Gentamycin, Chloroamphenicol, Sulphadiazine, Cephalotin)	D.N.S.	[28]
	Leaves extract/Ethanol	<i>Bacillus cereus</i> ,	MIC 512	
		<i>Clostridium perfringens</i> ,	N.A.	
		<i>Listeria monocytogenes</i>	512	
		<i>Staphylococcus epidermidis</i>	512	
		<i>Enterococcus faecalis</i>	N.A.	
		<i>Staphylococcus aureus</i>	N.A.	
		<i>Streptococcus pyogenes</i>	256	
		<i>Bacteroides fragilis</i>	N.A.	
		<i>Escherichia coli</i> ,	N.A.	
		<i>Pseudomonas aeruginosa</i>	N.A.	
		<i>Salmonella enteritidis</i>	N.A.	
		<i>Candida albicans</i>	N.A.	
	Positive control	(Ciprofloxacin, Tioconazole, Penicillin)	0.015–8	
<i>Jasminum angustifolium</i> (L.) Willd.	Flower extract / Methanol (500 ppm)	<i>Bacillus</i> sp.	ZOI N.A.	[29]
		<i>Escherichia coli</i>	N.A.	
		<i>Staphylococcus</i> sp.	N.A.	
		<i>Klebsiella pneumoniae</i>	N.A.	
		<i>Lactobacillus</i> sp.	6	
		<i>Yersinia</i> sp.	5.5	
		<i>Enterococcus</i> sp.	5	
		<i>Pseudomonas</i> sp.	6	
	Positive control		D.N.S.	
<i>Jasminum angustifolium</i> var. <i>sessiliflorum</i> (Vahl) P.S.Green	Leaves and Stem extracts/Ethanol (25 mg/50 µL)	<i>Escherichia coli</i>	ZOI S: N.A; L: N.A	[30]
		<i>Pseudomonas aeruginosa</i>	S: 22; L: 17	
		<i>Staphylococcus aureus</i>	S: 12; L: N.A	
		<i>Enterococcus faecalis</i>	S: 14; L: 11	
		<i>Bacillus cereus</i>	S: 12; L: 11	

	Positive control	<i>Candida albicans</i> [Chloramphenicol (30 µg/well)]	S: 13; L: 15 24–30 except <i>Pseudomonas aeruginosa</i>	
	Leaves extract / Methanol (50 mg/mL)		ZOI	
<i>Jasminum arborescens</i> Roxb.		<i>Escherichia coli</i>	2.8	[31]
		<i>Pseudomonas aeruginosa</i>	3.1	
		<i>Staphylococcus aureus</i>	3.7	
		<i>Bacillus subtilis</i>	3.6	
	Positive control	[Streptomycin (1mg/mL)]	2.8–3.6	
	Leaves extract /Ethanol		MIC	
<i>Jasminum auriculatum</i> Vahl		<i>Bacillus subtilis</i> ,	1560	[32]
		<i>Staphylococcus aureus</i>	6250	
		<i>Pseudomonas aeruginosa</i>	780	
		<i>Micrococcus luteus</i>	3125	
		<i>Escherichia coli</i>	12500	
		<i>Aspergillus niger</i>	N.A.	
		<i>Candida albicans</i>	N.A.	
	Positive control	(Ciprofloxacin against bacterial strains Fluconazole against fungi)	1.25–2.5 2.5	
	Leaves extract / Acetone (30 mg/mL)		ZOI	
		<i>Staphylococcus aureus</i>	20	[33]
		<i>Bacillus cereus</i>	24	
		<i>B. subtilis</i>	9	
		<i>Escherichia coli</i>	14	
		<i>Pseudomonas sp.</i>	17	
	Positive control		D.N.S.	
<i>Jasminum azoricum</i> L.	Flowers extract/Butanol (500 mg/mL)	<i>Salmonella typhi</i>	22	[34]
		<i>Staphylococcus aureus</i>	15	
		<i>Pseudomonas sp.</i>	20	
		<i>Vibrio cholerae</i>	18	
		<i>Streptococcus sp.</i>	17	
		<i>Corynebacterium sp.</i>	14	
		<i>Enterobacter aerogenes</i>	N.A.	
		<i>Proteus vulgaris</i>	18	
		<i>Escherichia coli</i>	21	
	Positive control	(Ampicillin)	D.N.S.	
	Leaves extract/Acetone, Water, Methanol, Petroleum ether, Jatamansone		MIC	
<i>Jasminum breviloobum</i> DC.		<i>Staphylococcus aureus</i>	D: 0.44; E: 0.92; F: 1.17; G: 1.56; H: 0.05	[35]
			D: 0.42; E: 0.62; F: 1.04; G: 1.36; H: 0.15	
		<i>Bacillus subtilis</i>	D: 0.89; E: 1.24; F: 1.09; G: 1.08; H: 0.07	
		<i>Escherichia coli</i>	D: 0.54; E: 0.66; F: 0.95; G: 1.00; H: 0.14	
		<i>Klebsiella pneumoniae</i>	D: 0.49; E: 0.51; F: 0.60; G: 0.92; H: 0.09	
		<i>Proteus mirabilis</i>		
		Positive control		
			D.N.S.	

<i>Jasminum fluminense</i> Vell.	Root extracts / Methanol	<i>Candida albicans</i>	MIC 3100
		<i>Gardnerella vaginalis</i>	<12500
		<i>Neisseria gonorrhoeae</i>	6300
		<i>Oligella ureolytica</i>	3100
	Positive control	(Ciprofloxacin)	10–< 10
<i>Jasminum grandiflorum</i> L.	Leaves extract/Aqueous and Ethanol (hot solvent)	<i>Streptococcus mutans</i>	MIC J: 6.25 E: 50
		<i>Lactobacillus acidophilus</i>	J: 25 E: 50
	Positive control	(Ciprofloxacin)	10–< 10
	Plant extract/Ethanol (500 µg/µL)	<i>Enterococcus faecalis</i>	ZOI N.A.
		<i>Hafnia alvei</i>	N.A.
		<i>Pseudomonas aeruginosa</i>	7
		<i>Proteus vulgaris</i>	N.A.
		<i>Plesiomonas shigelloides</i>	N.A.
		<i>Staphylococcus epidermidis</i>	15
		<i>S. aureus</i> ,	7
		<i>S. saprophyticus</i>	7
		<i>S. pyogenes</i>	7
		<i>Salmonella typhi</i>	N.A.
		<i>Shigella flexneri</i>	10
		<i>S. sonnei</i>	7
		<i>S. boydii</i>	7
		<i>S. dysenteriae</i>	6
	Positive control		D.N.S.
<i>Jasminum grandiflorum</i> subsp. <i>floribundum</i> (R.Br. ex Fresen.) P.S.Green	Plant extract/Methanol (10 mg/mL)	<i>Escherichia coli</i>	ZOI 14
		<i>Proteus vulgaris</i>	12
		<i>Pseudomonas aeruginosa</i>	22
		<i>Staphylococcus aureus</i>	20
		<i>Sarcina lutea</i>	20
		<i>Bacillus subtilis</i>	15
		<i>Mycobacterium phlei</i>	N.A.
		<i>Candida albicans</i>	22
	Positive control	(Ofloxacin, Amphotericin B)	D.N.S.
<i>Jasminum nervosum</i> Lour. (Synonym <i>Jasminum subtriplinerve</i> Blume)	Stem and leaves extract/petroleum ether, ethyl acetate, ethanol, methanol and water	<i>Escherichia coli</i>	MIC F: 200 AE: N.A.
		<i>Pseudomonas aeruginosa</i>	G: 100
		<i>Bacillus subtilis</i>	I and J: 200
		<i>Staphylococcus aureus</i>	
		<i>Aspergillus Niger</i>	AE: N.A.
		<i>Fusarium oxysporum</i>	AE: N.A.
		<i>Candida albicans</i>	AE: N.A.
		<i>Saccharomyces cerevisiae</i>	AE: N.A.
	Positive control		D.N.S.

	Leaves extract / Methanol [80% methanol at a ratio of 1:5 (v/v, dry plant material/solvent)] Positive control	<i>Fusarium solani</i> <i>F. oxysporum</i> <i>Rhizoctonia solani</i>	N.A. N.A. Active	[41]
			D.N.S.	
<i>Jasminum officinale</i> L.	Essential oil from flowers extract Positive control	<i>Trichosporon ovoides</i> [Imidazole (50µg/disc) Nystatin B (100µg/disc)]	MIC 3.1 12.5 6.2	[42]
			ZOI	
<i>Jasminum polyanthum</i> Franch.	Flower and leaf extracts / water extract (2 g flowers as well as leaves used for extract preparation)	<i>Escherichia coli</i> <i>Klebsiella pneumoniae</i> <i>Staphylococcus aureus</i> <i>Pseudomonas aeruginosa</i> <i>Aspergillus flavus</i> <i>A. niger</i>	Fl: 8; L: 7 Fl: 9; L: 8 Fl: 13; L: 11 Fl: 13; L: 12 Fl: 8; L: 10 Fl: N.A; L: N.A	[20]
	Positive control	(Gentamicin for bacterial strains)	10	
			ZOI	
<i>Jasminum</i> <i>syringifolium</i> Wall. ex G.Don	Leaves extract / Methanol (100 g leaves in 95% methanol)	<i>Escherichia coli</i> <i>Pseudomonas aeruginosa</i> <i>Staphylococcus aureus</i> <i>Bacillus cereus</i> <i>Staphylococcus epidermidis</i> <i>Vibrio cholerae</i> <i>Proteus mirabilis</i> <i>Shigella flexneri</i> <i>Salmonella enterica typhi</i> <i>Klebsiella pneumoniae</i> <i>Aspergillus niger</i> <i>Candida albicans</i>	21.33 16.67 21.67 22.33 16.33 18.67 15.33 22.67 19.33 18.33 17.33 15.33	[43]
	Positive control	(Gentamycin for bacterial strains Nystatin for fungi)	12.67–22.67 17.67–21.33	

S: Stem; L: Leaves; Fl: Flower; D: Acetone; E: Water; F: Methanol; G: Petroleum ether; H: Jatamansone; I: Ethyl acetate; J: Ethanol; AE: All extracts (petroleum ether, ethyl acetate, ethanol, methanol and water extracts), N.A.: Not active; ZOI: Zone of inhibition; MIC: Minimum inhibitory concentration; D.N.S.: Data not shown.

Compared to Gram negative ones, the impact is more obvious in the case of Gram positive pathogens. The difference in susceptibility between Gram negative positive strains is due to structural dissimilarities and composition of membranes [44,45]. Thus, from Table 1, it was concluded that, among other species, *Jasminum azoricum* and *Jasminum brevilobum* were found to be the most active species.

However, most of the plant species displayed no fungal activity except for a few species. Methanol extract of *Jasminum grandiflorum* subsp. *floribundum* has demonstrated anti-fungal activity against *Candida albicans* with a zone of inhibition of 22 mm. Meanwhile, essential oil from flowers extract of *Jasminum officinale* has shown anti-fungal activity against the *Trichosporon ovoides* with MIC 3.1µg/mL.

5. Role of *Jasminum* Plants in Combating Resistance

In the 21st century, antibiotic resistance has become a serious public health concern. Bacterial and fungal strains constantly develop new ways through various unknown/un-described mechanisms to adapt and withstand the biostatic or lethal effects of antibiotics [46,47]. Key factors leading to resistance include misuse and abuse of antimicrobials over decades, inadequate laboratory resources, and poor surveillance. In particular, their introduction to human as well as veterinary medicine contributed a lot in this regard [48–51].

5.1. Bacterial Antibiotic Resistance

The antibacterial drug's mechanism usually involves degradation of the bacterial membrane, and an inhibitory effect on biosynthesis of the cell wall and synthesis of nucleic acid [52–54]. Bacterial strains have number of well-differentiated mechanisms by which they survive and develop antibiotic resistance [50,55,56]. The mechanistic basis of resistance includes numerous paths, such as molecular target alteration, efflux pumps' overexpression, formation of biofilm, antibiotic degradation or modification, enzyme mediated destruction, and modification of bacterial target structures [49,50,57–62]. These mechanisms are shown in Figure 2. The mechanisms mentioned above assist bacteria to withstand pressure of antibiotic selection. Hydrolysis, functional group transfer, and structural modifications of antibiotics can be caused by a wide range of bacterial enzymes, thereby limiting their effectiveness. The standard process of making the β -lactam class of antibiotics ineffective is hydrolysis [63–65]. In bacterial strains, efflux pumps constitute the major resistance mechanism as their hyperactivity in resistant strains efflux antimicrobials outside the cell, reducing their concentration and thereby rendering them inefficient [59,66]. These mechanisms might be innately encoded within bacterial chromosome and through random mutations in chromosomal genes [56]. In addition, plasmids containing resistance genes can confer antimicrobial resistance [46].




Figure 2. Major antibiotic resistance mechanisms evolved among bacterial strains and impact of *Jasminum* spp. against them (Created using Biorender.com).

5.2. Antibiotic Resistance in Fungi

Fungal infections seem to be a critical threat in clinical research over the past few decades, with immune-compromised individuals becoming readily susceptible. In particular, fungal infections are commonly linked to higher mortality [67]. Moreover, in healthy populations with an increased occurrence of fungal pathogens such as *Aspergillus fumigatus*, there are increasing indications of fungal infections, rendering fungi a potent threat. *Candida auris*, for example, made headlines as an emerging pandrug-resistant microorganism to effective antifungal drugs [67]. Four major groups, namely azoles, polyenes, echinocandins, and allylamines constitute currently available antifungal drugs. Ergosterol (polyenes), its biosynthetic route (allylamines and azoles), and β -glucan synthesis (echinocandins) are among the targets of many antifungal drugs; unfortunately, these protein molecules can trigger the emergence of resistance [68]. Polyenes and azoles hinder biosynthesis of ergosterol, a sterol found in fungal membranes. However, their use in humans is confined because of their toxicity that affects mammalian cholesterol which has structural similarities with ergosterol of fungal strains [68]. Considering that drug options are minimal, studies examining resistance mechanisms to current antifungals are valuable.

In this study, azoles resistance mechanism of *Candida albicans* and *Aspergillus fumigatus* have been reviewed (Figure 3). Azole resistance may develop through a variety of mechanisms, including drug target overexpression or modification, drug transporter up-regulation, or cellular changes that minimise drug toxicity or allow tolerance to drug-induced stress [69,70].

The resistance mechanism of *Candida albicans* involves mutations in lanosterol 14- α demethylase (ERG11); transcription factor UPC2, TAC1 (transcriptional activator of CDR genes), and MRR1 (transcription factor); heterozygosity loss; and overexpression of Cdr1, Cdr2 (*Candida albicans* multidrug resistant protein), and Mdr1 (multidrug resistant protein 1). A number of stress-response pathway mediators (Hsp90 (Heat shock protein-90), Sgt1, calcineurin, KDACs (lysine deacetylases), PKC (protein kinase C) can also contribute to resistance. On the other hand, *Aspergillus fumigatus* resistance is related to mutation at Gly54 (position) in the cyp51A gene, while overexpression of cyp51A is due to TR/L98H mutations. Additionally, there is overexpression of ATP Binding Cassette (ABC) transporter gene (AtrF), Mdr3, and Mdr4. Similar to *Candida albicans*, Hsp90 is major contributing stress-response pathway mediator [70–73].




Figure 3. Markers of antifungal resistance in *Aspergillus fumigatus* and *Candida albicans* and protective role of *Jasminum* plants against the same. Overexpression (O↑); Lanosterol 14- α demethylase (ERG11); Transcription factor (UPC2, MRR1),

Hsp90 (Heat shock protein-90), Cytochrome P450 14 α -sterol demethylases (*Cyp51*); Multidrug resistant (MDR); Complementarity-determining regions (CDRs), Lysine deacetylases (*KDAC*); Protein kinase C (PKC), ATP Binding Cassette transporter gene (*AtrF*); Transcriptional activator of CDR genes (*TAC1*).

5.3. Protective Role of *Jasminum* Species

The activity of plants is related to their bioactive composition. Although the mechanisms of action of plant bioactive substances (PBS) are not clear, they are assumed to intervene with cell membrane organisation, leading to decreased membrane potential and lower levels of synthesis of ATP. The addition of PBS to the medium induces cellular membrane permeability, chelation of metal ions, and disruption of membrane-bound ATPase activity that alters the bacteria's physiological state and leads to the death of the bacterial strain [74,75]. PBS is capable of acting on many bacterial resistance production targets. They play an important role as drug-inactivating enzyme inhibitors, as well as being involved in inhibition of efflux pump over-expression. In addition, they inhibit synthesis of protein and DNA and also exhibit anti-biofilm activity (Figure 2). Several other researchers have expressed similar views. [59,60].

There are reports that carvacrol, thymol, as well as eugenol and catechins are reported for ATP depletion through membrane structure degradation leading to discharge of cellular components [76–78]. In addition, tea tree oil, consisting of monoterpenes, terpenes, sesquiterpenes, 1,8-cineol, alpha-terpineol, and terpinen-4-ol, is capable of interfering with the permeability of the membrane, destroying the cell membrane and obstructing cell development, causing cell death in resistant microbes (*Staphylococcus aureus*, *Escherichia coli*, and *Candida albicans* [79]).

A brief overview of the different phytoconstituents of *Jasminum* spp. is compiled here. In continuation, root bark of *Jasminum abyssinicum* contains flavonoids, glycosides, saponins, secoiridoid glucosides, terpenoids, and triterpenes [80,81], while leaves of *Jasminum angustifolium* contain alkaloids, flavonoids, phenolics, saponins, sterols, tannins, and terpenoids [82]. Similarly, *Jasminum angustifolium* var. *sessiliflorum* contains alkaloids, glycosides, phenols, quinones (anthraquinones), saponins, steroids, terpenoids, and tannins [83,84]. Moreover, leaves as well as flowers of *Jasminum arborescens* contain alkaloids (quinines), flavonoids, phenols, saponins, and terpenoids [20].

Jasminum auriculatum contains flavonoids, phenolics, and terpenoids [85]. Its flowers contain alkaloids, essential oil, flavonoids, glycosides, phenolic acid (salicylic acid), sterols, and tannins [86]. Leaves of *Jasminum azoricum* contain alkaloids, coumarins, flavonoids (kaempferol, quercetin, rutoside), glycosides, iridoid glucosides (azoricin, sambacin), polyphenols, quinones, sterols, steroids, tannins, triterpenes (amyrins), and terpenoids [33,87]. *Jasminum brevilobum*'s leaves contain sesquiterpenoids (jatamansone) [35]. Leaves and flowers of *Jasminum fluminense* contain alkaloids, flavonoids, glycosides, phenols, saponins, triterpenes (squalene), and tannins [88]. *Jasminum grandiflorum* contains flavonoids (rutoside), monoterpenoids (geraniol, iridoids, secoiridoids), phenols (cresol), phenylpropanoid (eugenol), sesquiterpene alcohol (farnesol), phenolic acids, tannins, and terpenes [19,38,89]. *Jasminum nervosum* leaves contain alkaloids and flavonoids [89].

Jasminum officinale contains secoiridoid glycosides (aucubin, jasgranoside B, ligstroside, loganin, oleoside, oleuropein, and 8-dehydroxy shanzhiside), and flavonoid glycosides (sulfurein) [90–92], with leaves of *Jasminum officinale* containing phenols, phenylethanoids, flavonoids, and polyphenols [93,94]. Further, its stem contains sesquiterpenoids [95]. *Jasminum polyanthum*'s leaves and flowers contain alkaloids, phenols, quinines, saponins, and terpenoids [20]. Finally, leaves of *Jasminum syringifolium* contain alkaloids, flavonoids, triterpenoids, steroids, and tannins [43].

6. Antioxidant Potential of *Jasminum* spp.

In Table 2, the antioxidant potential of *Jasminum* plants is shown. The ethanol extract from the leaves of *J. abyssinicum* possessed strong antioxidant activity with IC₅₀ 26.3 μ g/mL, which was higher than the standard Trolox (IC₅₀ 5.8 μ g/mL) as per DPPH assay,

whereas it showed an ORAC value of 1023.7 $\mu\text{g TE/mg}$ extract. Moreover, a moderate amount of total phenolic content (401.3 $\mu\text{g GAE/mg}$) was also observed in the *J. abyssinicum* leaves extract by using the total phenolic content assay [96]. The study conducted by Moe et al. [30] demonstrated the antioxidant potential of ethanolic extract from *J. sessiliflorum* leaves and stems (0.5 mg/mL) by using DPPH, NO, and superoxide radical-scavenging assays as well as by measuring total phenolic content (TPC). In this study, ascorbic acid was used as a standard for DPPH (84.78 %) and NO (78.96%) assays, and Gallic acid was used against superoxide (83.24%) radicals. The study revealed that the extract from leaves showed 11.12%, 51.49%, and 51.29% inhibition of DPPH, NO, and superoxide radical-scavenging activity, respectively, while the extract from stems showed only superoxide radical-scavenging activity with a 53.93% inhibitory rate. Also, the total phenolic content observed by leaves and stem extract was 2.09 and 23.23 mg GAE/g, respectively. Dose-dependent (25–400 $\mu\text{g/mL}$) antioxidant activity of ethanol, chloroform, and petroleum ether leaves extract of *J. arborescens* was observed by Bhagath et al. [97] with DPPH inhibition ranging from 40–90% and reducing power activity ranging from 0.2–0.45 absorbance at 700 nm. The maximum effect was found in ethanol extract, preceded by chloroform, and petroleum ether extract.

In a subsequent study, the ethanol extract of *J. auriculatum* leaves showed DPPH scavenging activity with an IC_{50} value of 33.39 $\mu\text{g/mL}$ and total phenolic content of 8.47 mg GAE/g, whereas the standard, ascorbic acid, showed an IC_{50} value of 35.41 $\mu\text{g/mL}$ in DPPH scavenging assay [32]. Boiling water (BWE), and hydromethanolic (HME) extracts of *J. grandiflorum* flower buds revealed an antioxidant effect, as evaluated using DPPH, superoxide, nitric oxide, and hydroxyl peroxide scavenging activity. The standard ascorbic acid exhibited IC_{50} > 6.93, 372.85, 248.25, and > 24.78 $\mu\text{g/mL}$, respectively. It was found that BWE displayed IC_{50} values of 327.89 and 38.27 $\mu\text{g/mL}$ in superoxide and nitric oxide assays, respectively, whereas HME showed IC_{50} values of 1354.30 and 225.51 $\mu\text{g/mL}$. Similarly, both extracts, viz. HME (IC_{50} 189.93 $\mu\text{g/mL}$) and BWE (IC_{50} > 150.57 $\mu\text{g/mL}$) in DPPH assay were found to be less active than ascorbic (IC_{50} 6.93 $\mu\text{g/mL}$). However, BME showed an IC_{50} value of 397.09 $\mu\text{g/mL}$ in hydroxyl peroxide radical scavenging activity which was almost similar to HME (IC_{50} 403.31 $\mu\text{g/mL}$ [98]). Also, the ethanolic extract (JGLE) from leaves of *J. grandiflorum* displayed potent DPPH scavenging ability (IC_{50} 15 $\mu\text{g/mL}$) which was equivalent to ascorbic acid (IC_{50} 12 $\mu\text{g/mL}$). Moreover, JGLE also showed nitric oxide radical scavenging ability with IC_{50} 98 $\mu\text{g/mL}$ compared to standard, curcumin (IC_{50} 92 $\mu\text{g/mL}$). Furthermore, JGLE increased reducing power with IC_{50} 19.5 $\mu\text{g/mL}$, where the IC_{50} value for standard quercetin was 15.5 $\mu\text{g/mL}$. In the superoxide anion assay, reduction of nitro blue tetrazolium (NBT) was found to rise in a dose-dependent pattern [99].

Likewise, Chaturvedi and Tripathi [100] inferred that the methanolic leaves extract of *J. grandiflorum* have strong antioxidant potential as evaluated by using iron-induced lipid peroxidation, reducing power, and trapped $\text{ABTS}^{\bullet+}$, superoxide, and superoxide radicals scavenging assays. The results showed that the extract showed $\text{ABTS}^{\bullet+}$ and superoxide scavenging activity with EC_{50} 222.50 and 207 $\mu\text{g/mL}$, respectively, where vitamin C (EC_{50} 36.72 $\mu\text{g/mL}$) was used as a standard for $\text{ABTS}^{\bullet+}$ assay. Moreover, the extract exhibited lower reducing capabilities at a concentration of 71.42 $\mu\text{g/mL}$ compared to standard BHT (63.29 $\mu\text{g/mL}$) at 700 nm absorbance (Optical density 0.1). Concurrently, the extract inhibited iron-induced lipid peroxidation with EC_{50} 667.53 $\mu\text{g/mL}$, whereas the standards, BHT & quercetin, showed lipid peroxidation inhibition with EC_{50} 0.75 and 0.21 $\mu\text{g/mL}$, respectively. In the hydroxyl scavenging assay, the extract in the presence of EDTA scavenged hydroxyl radicals (non-site-specific reaction) with EC_{50} 288.19 $\mu\text{g/mL}$, while in the absence of EDTA (site-specific reaction) it showed EC_{50} 102.16 $\mu\text{g/mL}$. Along with this, the standard drug, BHT, showed EC_{50} 0.22 $\mu\text{g/mL}$ for site-specific reaction and at 0.58 $\mu\text{g/mL}$ for a non-site-specific reaction. Aqueous extract (500, 1000, 1500, and 2000 $\mu\text{g/mL}$) of *J. malabaricum* leaves, roots, and bark showed 7%, 22.2%, 44.4%, and 66.6% hydrogen peroxide scavenging activity, respectively, when compared with standard ascorbic acid (86%) [101].

The 90% methanolic and aqueous extracts of *J. mesnyi* leaves showed DPPH scavenging ability with IC_{50} 25.27 and 71.84 $\mu\text{g/mL}$, respectively, whereas the standard ascorbic acid and rutoside showed IC_{50} 8.84 and 3.78 $\mu\text{g/mL}$, respectively. Moreover, a concentration-dependent increase in reducing power was observed with both extracts (methanolic and aqueous) in the FRAP method. In addition, methanol extract, aqueous extract, and BHT (standard) displayed lipid peroxidation inhibitory activity with IC_{50} 84.69, 145.62, and 48.89 $\mu\text{g/mL}$, respectively [102].

The ethyl acetate and n-butanol fractions of methanolic leaves extract from *J. mesnyi* showed anti-oxidant potential using DPPH, nitric oxide (NO), and reducing power assay where standards, rutoside and ascorbic acid, were used. The n-butanol fraction showed high DPPH radical scavenging ability with IC_{50} 6.22 $\mu\text{g/mL}$ compared to ascorbic acid (IC_{50} 6.54 $\mu\text{g/mL}$) and rutoside (IC_{50} 5.44 $\mu\text{g/mL}$), while the ethyl acetate fraction revealed an IC_{50} value of 153.45 $\mu\text{g/mL}$. Moreover, n-butanol and ethyl acetate fractions inhibited NO scavenging with IC_{50} 35.12 and 141.54 $\mu\text{g/mL}$, respectively, as compared to rutoside (IC_{50} 29.93 $\mu\text{g/mL}$) and ascorbic acid (IC_{50} 21.06 $\mu\text{g/mL}$). Furthermore, n-butanol at a concentration range of 25–400 $\mu\text{g/mL}$ showed reductive capability with an absorbance range of 0.07–2.76 compared to ethyl acetate at 25–400 $\mu\text{g/mL}$ (0.05–1.11) when compared to rutoside (0.06–2.76) and ascorbic acid with absorbance ranges between 0.07–2.82 [103].

Subsequently, a recent study of the 80% methanolic leaves extract from *J. multiflorum*, *J. azoricum*, *J. humile*, *J. officinale*, and *J. sambac* from two different locations (Arabian nights and Grand Duke of Tuskany) possessed DPPH radical scavenging activity with IC_{50} 34.8, 199.2, 94.6, 76.6, 130.7, and 155.5 $\mu\text{g/mL}$, respectively, while showing total phenolic amounts of 167.3, 56.9, 88.0, 133.4, 47.3, and 50.2 $\mu\text{g GAE/mg}$, respectively. Moreover, extract from *J. azoricum*, *J. officinale*, *J. multiflorum*, *J. humile*, and *J. sambac* (Arabian nights and Grand Duke of Tuskany) had total flavonoid contents of 46.3, 34.7, 44.4, 38.3, 39.2, and 40.5 $\mu\text{g QE/mg}$, respectively [94]. Guo et al. [104] showed that the isolated compounds Jasnervosides A–H isolated from stems of *J. nervosum* showed DPPH radical scavenging activity with inhibitory percentage ranges of 18.44 to 82.6%. Among the tested compounds, Jasnervosides A, B, D, and G exerted strong antioxidant activity with IC_{50} 0.22, 0.09, 0.19, and 1.21 $\mu\text{g/mL}$, respectively, whereas ascorbic acid showed IC_{50} 0.88 $\mu\text{g/mL}$.

Jasminum nudiflorum water-soluble and fat-soluble flower fractions showed ferric-reducing antioxidant power (FRAP) activity of 11.05 and 3.71 $\mu\text{mol Fe(II)/g}$, respectively, with total phenolic content of 2.42 and 0.66 mg GAE/g, respectively. Moreover, water soluble and fat-soluble flower fraction revealed trolox equivalent antioxidant capacity of 3.85 and 0.79 $\mu\text{mol trolox/g}$, respectively [105]. In another study, aqueous extract of *J. officinale* leaves displayed antioxidant potential with IC_{50} 41.16, 30.29, 20.19, and 29.48 $\mu\text{g/mL}$ by using DPPH, nitric oxide, superoxide, and ABTS•+ radical scavenging assays, respectively, with ascorbic acid as a standard (IC_{50} 42.79, 36.74, 38.22, and 45.57 $\mu\text{g/mL}$, respectively). Moreover, the aqueous extract and ascorbic acid both showed a concentration-dependent reducing power (200–1000 $\mu\text{g/mL}$) as the absorbance increased with an increase in concentration by using reducing power assay [93]. Also, the 80% methanolic extract of *J. officinale* leaves displayed DPPH radical scavenging activity with IC_{50} value of 76.6 $\mu\text{g/mL}$ [94], whereas *J. multiflorum* flower methanolic extract showed DPPH radical scavenging activity with IC_{50} value of 81 $\mu\text{g/mL}$ [106]. Interestingly, the extract of *J. grandiflorum* dried flower buds proved to be a beneficial neuroprotective agent by acting on monoamine oxidase A (MAO-A), which catalyzes the reaction of monoamine deamination. Compared to the reference standard, clorgyline, *J. grandiflorum* extracts showed a higher MAO-A inhibiting activity, thereby supporting its antioxidant potential to alleviate symptoms of depression and lower cell oxidative injury [98].

Table 2. In vitro antioxidant activity of *Jasminum* spp.

Botanical Name	Part Used	Solvent/Compound/Conc.	Method Used and Major findings (IC ₅₀ and EC ₅₀ - µg/mL)	References
<i>Jasminum abyssinicum</i> Hochst. ex DC.	L	E	DPPH (IC ₅₀) = 26.3 ORAC = 1023.7 µg TE/mg	[96]
<i>Jasminum angustifolium</i> var. <i>sessiliflorum</i> (Vahl) P.S. Green (Synonym: <i>Jasminum sessiliflorum</i>)	L	E (0.5 mg/mL)	DPPH = 11.12% NO = 51.49% O ₂ ⁻ = 51.29%	[30]
	S	E (0.5 mg/mL)	O ₂ ⁻ = 53.93%	
<i>Jasminum arborescens</i> Roxb.	L	E, CH and PE (0.025–0.4 mg/mL)	DPPH = 40–90% Fe ⁺³ reducing power (absorbance at 700 nm) = 0.2 to 0.45	[97]
<i>Jasminum auriculatum</i> Vahl	L	E	DPPH (IC ₅₀) = 33.39	[32]
<i>Jasminum azoricum</i> L.	L	80% M	DPPH (IC ₅₀) = 199.2	[94]
<i>Jasminum grandiflorum</i> L.	F	BWE	DPPH (IC ₅₀) = 150.57 O ₂ ⁻ (IC ₅₀) = 327.89 NO (IC ₅₀) = 38.27 H ₂ O ₂ (IC ₅₀) = 397.09 DPPH (IC ₅₀) = 189.93 O ₂ ⁻ (IC ₅₀) = 1354.30 NO (IC ₅₀) = 225.51 H ₂ O ₂ (IC ₅₀) = 403.31	[98]
		HME		
	L	E	DPPH (IC ₅₀) = 15 Reducing power (IC ₅₀) = 19.5 NO (IC ₅₀) = 98	[99]
	L	M	Iron-induced lipid peroxidation (EC ₅₀) = 667.53 ABTS•+ (EC ₅₀) = 222.50 O ₂ ⁻ (EC ₅₀) = 207 OH (EC ₅₀) = 288.19 (+ EDTA) and 102.16 (- EDTA)	
				[100]
<i>Jasminum humile</i> L.	L	80% M	DPPH (IC ₅₀) = 94.6	[94]
<i>Jasminum malabaricum</i> Wight	L, R, B	Aq (500, 1000, 1500 and 2000 µg/mL)	H ₂ O ₂ = 7, 22.2, 44.4, and 66.6%	[101]
<i>Jasminum mesnyi</i> Hance	L	EA (25–400 µg/mL)	DPPH (IC ₅₀) = 153.45 NO (IC ₅₀) = 141.54 FRAP= concentration-dependent Reducing power (absorbance range) = 0.05–1.11	[103]
		n-but (25–400 µg/mL)	DPPH (IC ₅₀) = 6.22 NO (IC ₅₀) = 35.12 FRAP= concentration-dependent Reducing power (absorbance range) = 0.07–2.76	
		M	DPPH (IC ₅₀) = 25.27 Lipid peroxidation assay (IC ₅₀) = 84.69	
	L	Aq	DPPH (IC ₅₀) = 71.84 Lipid peroxidation assay (IC ₅₀) = 145.62	[102]
<i>Jasminum multiflorum</i>	L	M	DPPH (IC ₅₀) = 34.8	[94]

(Burm.f.) Andrews				
	F	M	DPPH (IC ₅₀) = 81	[106]
<i>Jasminum nervosum</i> Lour.	S	Jasnervosides A *	DPPH (IC ₅₀) = 0.22	[104]
		Jasnervoside B *	DPPH (IC ₅₀) = 0.09	
		Jasnervoside D *	DPPH (IC ₅₀) = 0.19	
		Jasnervoside G *	DPPH (IC ₅₀) = 1.21	
<i>Jasminum nudiflorum</i> Lindl.	F	Water-soluble (tetrahydrofuran)	FRAP = 11.05 µmol Fe(II)/g TEAC = 3.85 µmol trolox/g	[105]
		Fat soluble (methanol–acetic acid–water mixture; 0:3.7:46.3)	FRAP = 3.71 µmol Fe(II)/g TEAC = 0.79 µmol trolox/g	
<i>Jasminum officinale</i> L.	L	Aq	DPPH (IC ₅₀) = 41.16 NO (IC ₅₀) = 30.29 O ₂ ^{•−} (IC ₅₀) = 20.19 ABTS•+ (IC ₅₀) = 29.48	[93]
<i>Jasminum sambac</i> (L.) Aiton	L	80% M	DPPH (IC ₅₀) = 76.6	[94]
	F	M	DPPH (IC ₅₀) = 208	[106]
	L (Arabian nights)	80% M	DPPH (IC ₅₀) = 130.7	[94]
	L (Grand Duke of Tuskany)	80% M	DPPH (IC ₅₀) = 155.5	

L: Leaves; R: Roots; B: Bark; S: Stem; F: Flower; WP: Whole plant; *: compound; E: Ethanol; Aq: Aqueous; M: Methanol; DPPH: DPPH radical scavenging activity; H₂O₂: Hydrogen peroxide radical scavenging activity; TEAC: Trolox equivalent antioxidant capacity; ABTS•+: ABTS radical scavenging activity; OH: Hydroxyl radical scavenging activity; NO: Nitric oxide radical scavenging activity; FRAP: Ferric reducing antioxidant power; O₂^{•−}: Superoxide radical scavenging activity; ORAC: Oxygen radical absorbance capacity; CH: Chloroform; PE: Petroleum ether; BWE: Boiling water extract; EA: Ethyl acetate; n-but: n-butanol; HME: Hydromethanolic; +EDTA: In presence of Ethylenediaminetetraacetic acid (EDTA); - EDTA: In absence of EDTA, IC₅₀: Half maximal inhibitory concentration; %: Percent scavenging.

Antioxidant activity is challenging to distinguish on basis of a single test model. Several in vitro methods that are used to assess the antioxidant effect of the desired samples such as DPPH radical scavenging assay, Hydroxyl scavenging assay, ABTS scavenging assay, Oxygen radical absorbance capacity (ORAC) LPO inhibition capacity (LPIC) assay, β-carotene–linoleic acid (linoleate) assay, and so forth. These test techniques differ from one another based on cost, accessibility, etc.

It is evident from Figure 4 that in in vitro study, four methods that are most frequently used are DPPH > Nitric oxide > superoxide radical > hydrogen-peroxide radical scavenging assay. On the basis of the most used method, phenylpropanoid glycoside (Jasnervoside B), isolated from the stems of *J. nervosum*, exhibited strong antioxidant potential with IC₅₀ 0.09 µg/mL. Considerably, DPPH is considered the quickest, simplest, and rational approach out of all the in vitro methods, and thus it is used mainly for a sample's antioxidant activity assessment. Further, Figure 4 shows that *Jasminum* spp. leaves have the highest frequency of plant parts used, followed by flower, stems, whole plant, and LRB (leaves, roots, and bark).




Figure 4. Frequency of method employed and plant parts used in antioxidant studies.

7. Oxidative Stress Related Diseases

Oxidative stress caused by ROS damages biomolecules (lipids, proteins, or DNA) thus contributing to cell survival regulation, inflammation, and stress responses [107–109]. Prolonged oxidative stress results in damage of body organs, which can potentially lead to the progression of chronic diseases like myocardial infarction, rheumatoid arthritis, diabetes, inflammatory diseases, cancer, vascular diseases, neurodegenerative diseases, and other metabolic diseases [110,111]. Enzymatic and non-enzymatic antioxidants provide a defense mechanism against free radicals by quenching or scavenging them from having harmful effects on the body. Catalase, thioredoxin, coenzyme Q, glutathione peroxidase, beta carotenoids, superoxide dismutase, polyphenols, glutathione, glutathione transferase, and glutathione reductase are widely evaluated antioxidants in the treatment of oxidative damage related diseases [112–115].

8. Impact of *Jasminum* Plants against Oxidative Stress In Vivo

The role of *Jasminum* spp. in combating oxidative stress related disorders is highlighted in Table 3. The anti-lipid peroxidative potential and chemopreventive efficacy of ethanolic extract (JgEt) from flowers of *J. grandiflorum* was evaluated on 7,12-enz(a)-anthracene (DMBA; 25 mg, s.c.)-induced Wistar albino rat mammary carcinogenesis. The extract (300 mg/kg p.o.) completely prevented the occurrence of tumours, while preneoplastic lesions that were mild to moderate (hyperplasia, dysplasia, and keratosis) were found in histopathological evaluation of extract-treated rats. Moreover, JgEt significantly ($p < 0.05$) downregulated the levels of TBARS and improved the antioxidant status when compared with the DMBA-treated group. In addition, the extract markedly incremented ($p < 0.05$) in vitamin C level (in plasma), vitamin E level (in plasma and erythrocytes), and reduced glutathione level (in plasma and erythrocytes) with respect to the DMBA group. Also, superoxide dismutase (SOD), glutathione peroxidase (GPx), and catalase (CAT)

levels were increased in mammary plasma, erythrocytes, and tissues of DMBA-treated rats, however, levels of vitamin E, glutathione peroxidase, and reduced glutathione were lowered ($p < 0.05$) in mammary tissue of experimental animals as compared to DMBA-treated rats. It was concluded that the extract showed chemopreventive efficacy in experimental mammary carcinogenesis [116]. Also, the hydromethanolic (HME) and boiling water (BWE) extracts of *J. grandiflorum* flower buds (dried) were assessed for in vitro efficacy towards central nervous system (CNS) disorders by measuring acetylcholinesterase (AChE), monoamine oxidase A (MAO-A), and butyrylcholinesterase (BuChE) inhibitory activity. It was observed that, both the extracts displayed MAO-A inhibitory activity with IC_{50} values of 603.16 $\mu\text{g/mL}$ (HME) and 699.74 $\mu\text{g/mL}$ (BWE) whereas the reference compound (clorgyline) showed an IC_{50} value $> 0.012 \mu\text{g/mL}$. Moreover, BWE, HME, and galantamine (reference compound) exhibited AChE inhibition with IC_{25} 1731.08, 1913.06, and 0.79 $\mu\text{g/mL}$, respectively. In addition, BuChE was highly inhibited by HME (IC_{50} 2610.87 $\mu\text{g/mL}$) followed by BWE (IC_{50} 5175.75 $\mu\text{g/mL}$) but weaker than reference compound, galantamine (IC_{50} 4.71 $\mu\text{g/mL}$). It was concluded from the study that the dried flower buds from *J. grandiflorum* can be used in treating psychiatric disorders and this activity is associated with antioxidant protection [98].

Similarly, Chaturvedi and Tripathi [100] showed that the methanolic leaves (100–800 $\mu\text{g/mL}$) extract of *J. grandiflorum* significantly ($p < 0.001$) inhibited LPS (20 ng/mL)-induced NO production in peritoneum fluid isolated macrophages from normal healthy Charles Foster (CF) strain albino rats in a concentration-dependent manner (300–800 $\mu\text{g/mL}$) with an inhibitory range of 9.5 to 4.41 $\mu\text{M}/1 \times 10^5$ cells as compared with an experimental control value (14.15 $\mu\text{M}/1 \times 10^5$ cells). Additionally, the wound healing effect of ointment (2% and 4%, topically) prepared using *J. grandiflorum* leaves methanolic extract was evaluated on cutaneous wound healing in diabetic Charles Foster (CF) strain albino rats. The extract (2% and 4%) significantly contracted wounds by 76.35% ($p < 0.05$) and 96.12% ($p < 0.01$), respectively, on day 12 as compared to the diabetic control group (62.94%). Moreover, the levels of total hydroxyl proline, hexosamine, protein, and DNA were all substantially higher ($p < 0.01$) at 4% ointment treatment, and a little less significant ($p < 0.05$) at 2% ointment treatment when compared to the diabetic control group. Furthermore, 4% of ointment-treated wounds displayed highly significant ($p < 0.01$) tensile strength on day 10 in comparison to the untreated wounds. Histopathological analysis revealed that ointment in a concentration-dependent manner (2% and 4%) increased fibrous tissue, collagen, and blood vessels. Also, ointment at 4% significantly led to the rise of SOD ($p < 0.05$), catalase ($p < 0.001$), and GSH ($p < 0.05$) content, whereas it decreased lipid peroxidation level ($p < 0.05$) in wound tissue with respect to the diabetic control group. Additionally, this ointment at 2% significantly ($p < 0.05$) effected only catalase level. Thus, it was concluded that the wound healing effect of ointment from *J. grandiflorum* leaves was through antioxidants [117]. The anti-inflammatory potential of the ethanol root extract (EJS; 100, 200, and 400 mg/kg, p.o.) of *J. sambac* was investigated using acute (carrageenan-induced paw edema), and sub-chronic (cotton pellet-induced granuloma) inflammation model of Charles Foster albino rats using diclofenac (10 mg/kg p.o.) as a reference standard. It was found that, EJS (400 mg/kg) and standard significantly ($p < 0.001$) inhibited rat paw edema after 3, 4, and 6 h of treatment as compared to the untreated control. Moreover, EJS inhibited granuloma formation by 3.7%, 5.93%, and 33.58% at 100, 200, and 400 mg/kg, while diclofenac showed 43.40% inhibition in granuloma formation. In addition, EJS extract decreased AST ($p < 0.05$ and $p < 0.05$), ALT ($p < 0.05$ and $p < 0.05$), and lipid peroxidation ($p < 0.05$ and $p < 0.01$) levels, whereas it increased SOD ($p < 0.05$ and $p < 0.01$) and catalase ($p < 0.001$) in rats edematous tissue after acute and sub-chronic inflammation exposure, respectively in comparison to respective model groups. Also, EJS extract decreased lipid peroxidation ($p < 0.001$) levels whereas enhanced SOD ($p < 0.05$ and $p < 0.05$) and catalase ($p < 0.01$) in the serum of acute and sub-chronic inflammation model, respectively, with respect to their model group. Likewise, the standard, diclofenac, markedly ($p < 0.05$) reversed the altered parameters in serum and edematous tissue by both the models [118].

In another study, the anti-aging effect of *J. sambac* flower extract fermented with *Lactobacillus rhamnosus* (F-FEJS, 0.1%, 0.25%, 0.5%, 1.0%, 2.5%) was assessed in UVB (40 mJ/cm²) or H₂O₂ (200 µM) -induced aging in HS68 dermal fibroblast cells. It was observed that the extract (2.5%) significantly ($p < 0.001$) reduced intracellular reactive oxygen species production (ROS) stimulated by UVB or H₂O₂ as revealed by fluorescence microscopy and flow cytometry analysis. Moreover, the extract (2.5%) markedly decreased p53, p21 and p16 levels in H₂O₂ ($p < 0.001$, $p < 0.001$ & $p < 0.001$, respectively) and UVB treated HS68 cells when compared to H₂O₂ and UVB treated cells. In addition, the extract led to MMP-1 inhibition, SA-β-Gal positive cells, p-JNK, p-P38, p-ERK, and p-c-jun protein levels whereas efficiently upregulated collagen synthesis-related pathway components (p-smad2/3, TGF-β, COL3A1, and COL1A1), p-Nrf2 and antioxidant gene expression (HO-1) levels with respect to H₂O₂/UVB treated cells. Likewise, the extract enhanced p-Nrf2 nuclear translocation while down-regulated p-c-jun in the nuclear fractions. Further, it was confirmed that *J. sambac* flower extract attenuated H₂O₂/UVB-induced aging, ROS production, and degradation of collagen in HS68 cells through smad2/3, Nrf2, and c-jun pathways [119].

Table 3. Effect of *Jasminum* spp. against oxidative stress-related diseases.

Botanical Name	Plant Part	Solvent/Dose	Activity	Model	Biomarkers Affected	References
<i>Jasminum grandiflorum</i> L.	F	E (300 mg/kg p.o.)	Chemo preventive	7,12-dimethylbenz(a)anthracene (DMBA)-induced rat mammary carcinogenesis	↑ vitamin E (plasma and erythrocytes) ↑ vitamin C (plasma) ↑ reduced glutathione (plasma and erythrocytes) ↑ SOD, CAT (plasma, erythrocytes and mammary tissues) ↑ glutathione peroxidase (plasma, erythrocytes) ↓ TBARS ↓ reduced glutathione (tissue) ↓ glutathione peroxidase (tissue)	[116]
	L	M (100–800 µg/mL)	Anti-inflammatory	LPS (20 ng/mL)-induced nitric oxide in rat peritoneal macrophage	↓ NO production (13.26 µM/1 × 10 ⁵ cells to 4.41 µM/1 × 10 ⁵ cells)	[100]
	L	O	Wound healing	Cutaneous wound healing in diabetic rats	↑ wound contraction ↑ total hydroxyl proline, ↑ hexosamine ↑ protein ↑ DNA content ↑ Tensile strength ↑ collagen & fibrous tissue ↑ number of blood vessels ↑ SOD, CAT and GSH content ↓ lipid peroxidation	[117]
<i>Jasminum sambac</i> (L.) Aiton	R	E	Anti-inflammatory	Carrageenan-induced rat paw edema model and cotton pellet-induced granuloma in rats	↓ paw edema ↓ granuloma formation ↓ AST, ALT, LPO, ↑ SOD and CAT	[118]
	F	LFE	Anti-aging	UVB (40 mJ/cm ²) or H ₂ O ₂ (200 µM) -induced HS68 dermal fibroblast cell	↓ ROS production ↓ aging markers, such as p16, p21, and p53, ↓ MMP-1	[119]

↓ SA-β-Gal -positive cells
 ↓ p-ERK, p-JNK, p-P38, and p-c-jun protein levels
 ↑ p-smad2/3 in the nuclear fraction
 ↑ TGFβ, p-smad2/3, COL1A1, and COL3A1 protein levels
 ↑ phospho-Nuclear respiratory factor 2 and antioxidant gene expression (HO-1)

F: Flower, R: Roots, L: Leaves, E: Ethanol, M: Methanol, BWE: a boiling water extract, HME: Hydromethanolic extract, LFE: *Lactobacillus rhamnosus* fermented extract, O: ointment, LPS: Lipopolysaccharide, UVB: Ultraviolet B, H₂O₂: Hydrogen peroxide, SOD: Superoxide Dismutase, CAT: Catalase, TBARS: Thiobarbituric acid reactive substances (lipid peroxidation), NO: Nitric Oxide, GSH: Glutathione, AST: Aspartate transaminase, ALT: Alanine transaminase, LPO: lipid peroxidation, ROS: Reactive Oxygen Species, MMP-1: Matrix metalloproteinase-1, SA-β-Gal: Senescence-associated beta-galactosidase; COL1A1: Collagen Type I Alpha 1; COL3A1: Collagen Type III Alpha 1, and TGF-β: Transforming growth factor-beta.

With a detailed understanding of the role of diseases generated by oxidative stress, *Jasminum* provided a useful approach in relation to their respective criteria for possible interventions in diseases related to oxidative stress (Table 3). Sengar et al. [118] scientifically validated the anti-inflammatory effect of *J. sambac* ethanol root extract against acute and chronic inflammation models with respect to their reference standard, diclofenac. The results suggested that the diclofenac illustrated efficient restoration of altered biochemical parameters in both acute and sub-chronic models' edematous and granulomatous tissues than the plant extract. This plant has been used since ancient times as an anti-inflammatory, anti-pyretic, and anti-nociceptive agent [120,121]. Also, *J. sambac* leaves have been studied for their anti-inflammatory [122] and analgesic properties [123]. Due to the obvious negative effects of non-steroidal anti-inflammatory medicines (NSAIDs) and opioids, there is a strong demand for new products with minimal or no side effects and medicinal plants such as *Jasminum* will play a crucial role in this context.

Jasminum plants demonstrated protective activity against oxidative stress (triggered by free radicals) and its associated disorders. These plants significantly raise the proportion of antioxidant enzymes such as SOD, Gpx, and CAT and their effect is related to a broad range of bioactive constituents (Table 3). In the role of plants in combating the resistance section, the bioactive constituents of some *Jasminum* spp. reviewed in this study have already been discussed. In addition, antioxidant activity has been shown by some other plants, and their bioactive components are listed below. Leaves, bark, and roots from *Jasminum malabaricum* contain alkaloids, phenolics, glycosides, flavonoids, steroids, saponins, tannins, and terpenoids [101]. *Jasminum mesnyi* contains caffeic glycoside esters, flavonoids, sterols, secoiridoid glucosides (jasmoside, jasminoside), and triterpenes (amyrin) [124,125]. Similarly, leaves and flowers from *Jasminum multiflorum* contain alkaloids, cardiac glycosides, flavonoids, phenols, sterols, terpenoids, and tannins [126]. *Jasminum nudiflorum*'s stems contain secoiridoid glucosides (jasnudiflosides and nudiflosides) [127], while its flowers contain phenols (epicatechin, gallic acid, and protocatechuic acid) [105]. Additionally, *Jasminum sambac* contain anthranils, sesquiterpenes (farnesol), alkaloids, flavonoids (hesperidin), terpenoids (oleanoic acid, geraniol), phenols, tannins, steroidal saponins (daucosterol) and sterols, monoterpenoids (iridoid glucosides (jasminin, sambacin, sambacoside A-G), geraniol), phenylpropanoid (eugenol), sesquiterpene alcohol (farnesol) [24,118,121,123,128,129].

9. Mechanistic Basis of ROS Neutralization

Adenosine triphosphates (ATPs), the energy currencies of the cell, are generated by mitochondria. Some low-energy electrons are released near the nucleus during the energy conversion, which are disposed of by the reduction of molecular oxygen to water, whilst a few of them escape and lead to the formation of superoxide radicals ($O_2^{\cdot-}$) [130]. In a biological system, there are several different types of free radicals, but those generated from oxygen like superoxide anion ($O_2^{\cdot-}$), singlet oxygen ($O=O$), are commonly referred to as reactive oxygen species (ROS). This superoxide anion can lead to the development of a variety of other reactive species like nitrosoperoxycarbonate, hydroxyl radical, peroxynitrite, and hydrogen peroxide through multiple chain reactions or pathways. However, certain key enzymes like glutathione peroxidase (GPx) and catalase (CAT), superoxide dismutase (SOD) assist to break down these free radicals into harmless and less active molecules (hydrogen peroxide/alcohol and O_2) [130,131–135]. Among all, SOD is a key player for radical neutralization (Figure 5).




Figure 5. First line anti-oxidant defense mechanism mediated by *Jasminum* spp. to neutralize free radicals. Glutathione Peroxidase (GPx); catalase (CAT); Hydrogen peroxide (H_2O_2); Lipid Radical (LR^{\cdot}); Water (H_2O); Oxygen (O_2); Superoxide Dismutase (SOD); Lipid Peroxide radical (LPR^{\cdot}); Hydroxyl radical (OH^{\cdot}); Fenton reaction (Fe^{2+}); Nitric Oxide (NO); Carbon dioxide (CO_2); Nicotinamide adenine dinucleotide phosphate oxidase (NADPH oxidase); Nitrosoperoxycarbonate ($ONOOCO_2^{\cdot-}$); Peroxynitrate ($ONOO^-$); Carbonate ($CO_3^{\cdot-}$); Nitrogen dioxide radical (NO_2^{\cdot}).

SOD is the first line defensive enzyme that assists in the dismutation of superoxide radicals into oxygen and hydrogen peroxide. In contrast, some of the hydrogen peroxide molecules in the presence of reduced iron (Fe^{2+}), in what is called a Fenton reaction, are reduced into the deleterious (OH^{\cdot}) hydroxyl radical [135]. The hydroxyl radical is one of the highly reactive radicals which can result in cell toxicity [136,137]. CAT, GPx, and other enzymes prevent the formation of hydroxyl radical (OH^{\cdot}), by degrading the hydrogen peroxide (H_2O_2) into O_2 and H_2O . Sometimes, the hydroxyl radical facilitates the formation of lipid radicals (LR^{\cdot}) by acting on the lipid membrane, which further leads to the formation of lipid peroxy radical (LPR^{\cdot}) in combination with oxygen. This can result in attenuation of membrane-bound enzymes activity [138], dysfunction of membrane receptors [139], altered membrane permeability [140], as well as enhancing the rigidity of the membrane while lowering its fluidity [141]. On the other hand, multifaceted antioxidant enzyme, GPx is known to act in the removal of peroxynitrite anion, hydrogen-peroxides, and lipid-peroxides [142]. In the presence of NADPH-oxidase enzyme and arginine,

superoxide anion and nitric oxide leads to the generation of peroxynitrite anion which is a powerful tissue-damaging oxidant. Furthermore, the peroxynitrite anion may often react with carbon dioxide, resulting in the formation of nitrosoperoxycarbonate, which gradually disintegrates to form nitrogen dioxide and carbonate radicals [143].

SOD, GPx, and CAT are imperative antioxidant enzymes that are crucial against the protection of the bio-system from free radicals. Hence, the proposed protective mechanisms of *Jasminum* plants explained in Figure 5 against free radicals might be that *Jasminum* plants would up-regulate the levels of the antioxidant enzymes which would further block the formation of the peroxynitrite anion or disintegrate hydrogen peroxide into water and oxygen.

10. Conclusions and Perspectives

In conclusion, most bio-activities were determined by the researchers, primarily for crude undefined extracts and the majority of the tests were conducted in vitro. In addition, for oxidative stress associated disorders, only 2 *Jasminum* species (*Jasminum grandiflorum* L. *Jasminum sambac* (L.) Aiton) have been studied (in vivo), yet they are highly successful in normalising various elevated parameters. The chemical profile of the *Jasminum* species revealed the presence of alkaloids, flavonoids, tannins, sterols, phenols, terpenoids, cardiac glycosides, terpenes, and secoiridoid glucosides. No studies have investigated the integrated role of *Jasminum* with standard drugs. It is expected that the data collected will serve as a useful protocol for researchers of herbal drug industry worldwide to explore various *Jasminum* species and their active components against human disorders. This review will provide more insight into the development of an effective drug candidate against diseases associated with oxidative stress and also against microbial diseases.

Authors Contribution: A.B. supervised first draft. A.R., A.K. (Abhishek Kumar) and A.K. (Ashwani Kumar) wrote the first draft of manuscript. A.R. and A.K. (Ashwani Kumar) contributed in making figures and providing literature. A.K. (Ashwani Kumar), V.A., P.T., R.V., D.K., P.O., O.K. and K.K. revised and improved the first draft. All authors have read and agreed to the published version of the manuscript.

Funding: Supported by the projects VT2019-2021 and Excellence FIM UHK (K.K., P.O., O.K.). Also supported by MH CZ - DRO (UHHK, 00179906) (K.K.).

Acknowledgments: The authors (A.B., A.R., A.K., A.K. (Ashwani Kumar), V.A. and P.T.) are grateful to Param Pujya Swami Ramdev Ji, Patanjali Yogpeeth, Haridwar and Patanjali Research Foundation Trust for providing all the necessary facilities. Also supported by the projects VT2019-2021 and Excellence FIM UHK (K.K., P.O., O.K.).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. WHO. *WHO Methods and Data Sources for Global Burden of Disease Estimates 2000–2011*; WHO: Geneva, Switzerland, 2013.
2. Gupta, M.; Sharma, R.; Kumar, A. Comparative potential of Simvastatin, Rosuvastatin and Fluvastatin against bacterial infection: An in silico and in vitro study. *Orient. Pharm. Exp. Med.* **2019**, *19*, 259–275, doi:10.1007/s13596-019-00359-z.
3. WHO. *WHO Antimicrobial Resistance: Global Report on Surveillance*; WHO: Geneva, Switzerland, 2014.
4. Baym, M.; Stone, L.K.; Kishony, R. Multidrug evolutionary strategies to reverse antibiotic resistance. *Science* **2016**, *351*, 1–21, doi:10.1126/science.aad3292.
5. WHO. *WHO Traditional Medicine Strategy*; WHO: Geneva, Switzerland, 2002.
6. Djeussi, D.E.; Noumedem, J.A.K.; Seukep, J.A.; Fankam, A.G.; Voukeng, I.K.; Tankeo, S.B.; Nkuete, A.H.L.; Kuete, V. Antibacterial activities of selected edible plants extracts against multidrug-resistant Gram-negative bacteria. *BMC Complement. Altern. Med.* **2013**, *13*, 1–8, doi:10.1186/1472-6882-13-164.
7. Duraipandiyan, V.; Ayyanar, M.; Ignacimuthu, S. Antimicrobial activity of some ethnomedicinal plants used by Paliyar tribe from Tamil Nadu, India. *BMC Complement. Altern. Med.* **2006**, *6*, 1–7, doi:10.1186/1472-6882-6-35.
8. Iwu, M.W.; Duncan, A.R.; Okunji, C.O. New antimicrobials of plant origin. In *Perspectives on New Crops and New Uses*; Janick, J. Ed.; ASHS Press: Alexandria, VA, USA, 1999; pp. 457–462.
9. Kumari, A.; Verma, R.; Sharma, M.; Chauhan, P.; Kumar, A. Evaluation of phytochemical, antioxidant, antibacterial and anticancerous activity of *Ficus auriculata* Lour. and *Osyris wightiana* Wall. ex Wight. *Bull. Environ. Pharmacol. Life Sci.* **2018**, *7*, 64–70.

10. Phaniendra, A.; Jestadi, D.B.; Periyasamy, L. Free Radicals: Properties, Sources, Targets, and Their Implication in Various Diseases. *Ind. J. Clin. Biochem.* **2015**, *30*, 11–26, doi:10.1007/s12291-014-0446-0.
11. Banerjee, J.; Das, A.; Sinha, M.; Saha, S. Biological Efficacy of Medicinal Plant Extracts in Preventing Oxidative Damage. *Oxidative Med. Cell. Longev.* **2018**, *2018*, 1–2, doi:10.1155/2018/7904349.
12. Halliwell, B. Antioxidants and Human Disease: A General Introduction. *Nutr. Rev.* **1997**, *55*, S44–S49, doi:10.1111/j.1753-4887.1997.tb06100.x.
13. Khan, F.; Garg, V.K.; Singh, A.K.; Kumar, T. Role of free radicals and certain antioxidants in the management of huntington's disease: A review. *J. Anal. Pharm. Res.* **2018**, *7*, 386–392, doi:10.15406/japlr.2018.07.00256.
14. Parmar, J.; Sharma, P.; Verma, P.; Goyal, P.K. Chemopreventive action of *Syzygium cumini* on DMBA-induced skin papillomagenesis in mice. *Asian Pac. J. Cancer Prev.* **2010**, *11*, 261–265.
15. Kumar, A.; Kumar, D. Development of antioxidant rich fruit supplemented probiotic yogurts using free and microencapsulated *Lactobacillus rhamnosus* culture. *J. Food Sci. Technol.* **2016**, *53*, 667–675, doi:10.1007/s13197-015-1997-7.
16. Raghuvanshi, D.; Dhalaria, R.; Sharma, A.; Kumar, D.; Kumar, H.; Valis, M.; Kuča, K.; Verma, R.; Puri, S. Ethnomedicinal Plants Traditionally Used for the Treatment of Jaundice (Icterus) in Himachal Pradesh in Western Himalaya—A Review. *Plants* **2021**, *10*, 232, doi:10.3390/plants10020232.
17. Priya, J.; Raja, D.P. Anti-bacterial activity studies of *Jasminum grandiflorum* and *Jasminum sambac*. *Ethnobot. Leaflet.* **2008**, *12*, 481–483.
18. KewScience-Plants of the World Online Home Page. Available online: <http://www.plantsoftheworldonline.org/> (accessed on 10 January 2021).
19. Ali, S.T.; Ayub, A.; Ali, S.N. Antibacterial activity of methanolic extracts from some selected medicinal plants. *FUUAST J. Biol.* **2017**, *7*, 123–125.
20. Prakkash, M.J.; Ragunathan, R.; Josteena, J. Evaluation of bioactive compounds from *Jasminum polyanthum* and its medicinal properties. *J. Drug Deliv. Ther.* **2019**, *9*, 303–310, doi:10.22270/jddt.v9i2.2413.
21. Rani, B.; Yadav, M.; Pachauri, G. Awesome medicinal benefits of jasmine plant. *J. Biol. Chem. Res.* **2017**, *34*, 918–922.
22. Upaganlawar, A.B.; Bhagat, A.; Tenpe, C.R.; Yeole, P.G. Effect of *Jasminum sambac* leaves extracts on serum glucose and lipid profile rats treated with alloxan. *Pharmacologyonline* **2003**, *1*, 1–6.
23. Lis-Balchin, M.; Hart, S.; Lo, W.H. Jasmine absolute (*Jasminum grandiflora* L.) and its mode of action on guinea pig ileum in vitro. *Phytother. Res.* **2002**, *16*, 437–439.
24. Kunhachan, P.; Banchonglikitkul, C.; Kajsongkram, T. Chemical composition, toxicity and vasodilatation effect of the flowers extract of *Jasminum sambac* (L.) Ait. “G. Duke of Tuscany”. *Evid. Based Complement. Alter. Med.* **2012**, *2012*, 1–7.
25. Khan, M.H.; Yadava, P.S. *Ethno Medicinal Plants of Manipur, North-East India (Thoubal District)*; Bishen Singh Mahendra Pal Singh: Dehradun, India, 2014; pp. 242–261.
26. Bhutya, R.K. *Ayurvedic Medicinal Plant of India*; Scientific Publishers: Jodhpur, India, 2011; Volume 1, pp. 253–254.
27. Geyid, A.; Abebe, D.; Debella, A.; Makonnen, Z.; Abera, F.; Tekla, F.; Kebede, T.; Urga, K.; Yersaw, K.; Biza, T.; et al. Screening of some medicinal plants of Ethiopia for their anti-microbial properties and chemical profiles. *J. Ethnopharmacol.* **2005**, *97*, 421–427, doi:10.1016/j.jep.2004.08.021.
28. Lulekal, E.; Rondevaldova, J.; Bernaskova, E.; Cepkova, J.; Asfaw, Z.; Kelbessa, E.; Kokoska, L.; van Damme, P. Antimicrobial activity of traditional medicinal plants from Ankober District, North Shewa Zone, Amhara Region, Ethiopia. *Pharm. Biol.* **2014**, *52*, 614–620, doi:10.3109/13880209.2013.858362.
29. Ramya, V.; Dhayan, V.D.; Umamaheswari, S. In vitro studies on antibacterial activity and separation of active compounds of selected flower extracts by HPTLC. *J. Chem. Pharm. Res.* **2010**, *2*, 86–91.
30. Moe, T.S.; Win, H.H.; Hlaing, T.T.; Lwin, W.W.; Htet, Z.M.; Mya, K.M. Evaluation of in vitro antioxidant, antiglycation and antimicrobial potential of indigenous Myanmar medicinal plants. *J. Integr. Med.* **2018**, *16*, 358–366, doi:10.1016/j.joim.2018.08.001.
31. Abhipsa, V.; Manasa, M.; Poornima, G.; Rekha, C.; Kekuda, T.R. In vitro antibacterial efficacy of selected plant extracts, streptomycin and their combination. *Asian J. Res. Chem.* **2012**, *5*, 791–793.
32. Mittal, A.; Satish, S.; Anima, P. Evaluation of wound healing, antioxidant and antimicrobial efficacy of *Jasminum auriculatum* Vahl. leaves. *Avicenna J. Phytomed.* **2016**, *6*, 295–304.
33. Thiruvengadam, S.; Nivedha, S.; Pujita, V.; Romauld, S.I. Detection of Antioxidant and Antimicrobial Activity of Leaf Extract of *Jasminum azoricum*. *Res. J. Pharm. Technol.* **2018**, *11*, 3629–3632, doi:10.5958/0974-360x.2018.00668.6.
34. SyamSree, K.; Anudeep, M.; Ramana, C.V.; Bhaskar, C. Screening of antimicrobial activity of flower extracts on human bacterial pathogens. *J. Pharmacog. Phytochem.* **2015**, *3*, 153–156.
35. Anoopkumar, A.N.; Aneesh, E.M.; Sudhikumar, A.V. Exploring the mode of action of isolated bioactive compounds by induced reactive oxygen species generation in *Aedes aegypti*: A microbes based double-edged weapon to fight against Arboviral diseases. *Int. J. Trop. Insect Sci.* **2020**, *40*, 573–585, doi:10.1007/s42690-020-00104-z.
36. Mamba, P.; Adebayo, S.A.; Tshikalange, T.E. Anti-microbial, anti-inflammatory and HIV-1 reverse transcriptase activity of selected South African plants used to treat sexually transmitted diseases. *Int. J. Pharmacog. Phytochem. Res.* **2016**, *8*, 1870–1876.
37. Nagarajappa, R.; Batra, M.; Sharda, A.J.; Asawa, K.; Sanadhya, S.; Daryani, H.; Ramesh, G. Antimicrobial Effect of *Jasminum grandiflorum* L. and *Hibiscus rosa-sinensis* L. Extracts Against Pathogenic Oral Microorganisms—An In Vitro Comparative Study. *Oral Health Prev. Dent.* **2013**, *13*, 441–448.

38. Rahman, M.; Khatun, A.; Khan, S.; Hossain, F.; Khan, A.A. Phytochemical, cytotoxic and antibacterial activity of two medicinal plants of Bangladesh. *Pharmacologyonline* **2014**, *1*, 3–10.
39. Abdel-Sattar, E.; Harraz, F.M.; El-Gayed, S.H. Antimicrobial Activity of Extracts of some Plants Collected from the Kingdom of Saudi Arabia. *JKAU Med. Sci.* **2008**, *15*, 25–33, doi:10.4197/med.15-1.3.
40. Ngan, D.H.; Hoai, H.T.C.; Huong, L.M.; Hansen, P.E.; Vang, O. Bioactivities and chemical constituents of a Vietnamese medicinal plant Che Vang, *Jasminum subtriplinerne* Blume (Oleaceae). *Nat. Prod. Res.* **2008**, *22*, 942–949, doi:10.1080/14786410701647119.
41. Nguyen, D.M.C.; Seo, D.-J.; Park, R.-D.; Jung, W.-J. Antifungal, Nematicidal and Antioxidant Activity of the Methanol Extracts Obtained from Medicinal Plants. *J. Appl. Biol. Chem.* **2013**, *56*, 199–204, doi:10.3839/jabc.2013.032.
42. Saxena, S.; Uniyal, V.; Bhatt, R.P. Inhibitory effect of essential oils against *Trichosporon ovoides* causing Piedra Hair Infection. *Braz. J. Microbiol.* **2012**, *43*, 1347–1354, doi:10.1590/s1517-83822012000400016.
43. Chander, M.P.; Pillai, C.R.; Sunish, I.P.; Vijayachari, P. Antimicrobial and antimalarial properties of medicinal plants used by the indigenous tribes of Andaman and Nicobar Islands, India. *Microb. Pathog.* **2016**, *96*, 85–98, doi:10.1016/j.micpath.2016.04.017.
44. Lambert, P.A. Cellular impermeability and uptake of biocides and antibiotics in Gram-positive bacteria and mycobacteria. *J. Appl. Microbiol.* **2002**, *92*, 46S–54S.
45. Kumar, A.; Singh, S.; Kumar, D. Evaluation of antimicrobial potential of cadmium sulphide nanoparticles against bacterial pathogens. *Int. J. Pharm. Sci. Rev. Res.* **2014**, *24*, 202–207.
46. Brown, E.D.; Wright, G.D. Antibacterial drug discovery in the resistance era. *Nature* **2016**, *529*, 336–343, doi:10.1038/nature17042.
47. Sekyere, J.O.; Asante, J. Emerging mechanisms of antimicrobial resistance in bacteria and fungi: Advances in the era of genomics. *Futur. Microbiol.* **2018**, *13*, 241–262, doi:10.2217/fmb-2017-0172.
48. Sekyere, J.O. Current State of Resistance to Antibiotics of Last-Resort in South Africa: A Review from a Public Health Perspective. *Front. Public Health* **2016**, *4*, 1–11, doi:10.3389/fpubh.2016.00209.
49. Sekyere, J.O.; Govinden, U.; Bester, L.A.; Essack, S.Y. Colistin and tigecycline resistance in carbapenemase-producing Gram-negative bacteria: Emerging resistance mechanisms and detection methods. *J. Appl. Microbiol.* **2016**, *121*, 601–617, doi:10.1111/jam.13169.
50. Sekyere, J.O.; Govinden, U.; Essack, S. The molecular epidemiology and genetic environment of carbapenemases detected in Africa. *Microb. Drug Resist.* **2016**, *22*, 59–68.
51. Voss-Rech, D.; Potter, L.; Vaz, C.S.L.; Pereira, D.I.B.; Sangioni, L.A.; Vargas, A.C.; de Avila Botton, S. Antimicrobial resistance in non-typhoidal *Salmonella* isolated from human and poultry-related samples in Brazil: 20-year meta-analysis. *Foodborne Pathog. Dis.* **2017**, *14*, 116–124.
52. Maxwell, A. DNA gyrase as a drug target. *Trends Microbiol.* **1997**, *5*, 102–109, doi:10.1016/s0966-842x(96)10085-8.
53. Tenover, F.C. Mechanisms of Antimicrobial Resistance in Bacteria. *Am. J. Med.* **2006**, *119*, S3–S10, doi:10.1016/j.amjmed.2006.03.011.
54. Schneider, T.; Sahl, H.-G. An oldie but a goodie—Cell wall biosynthesis as antibiotic target pathway. *Int. J. Med. Microbiol.* **2010**, *300*, 161–169, doi:10.1016/j.ijmm.2009.10.005.
55. Sekyere, J.O.; Amoako, D.G. Carbonyl Cyanide m-Chlorophenylhydrazine (CCCP) Reverses Resistance to Colistin, but Not to Carbapenems and Tigecycline in Multidrug-Resistant Enterobacteriaceae. *Front. Microbiol.* **2017**, *8*, 1–9, doi:10.3389/fmicb.2017.00228.
56. Sekyere, J.O.; Amoako, D.G. Genomic and phenotypic characterisation of fluoroquinolone resistance mechanisms in Enterobacteriaceae in Durban, South Africa. *PLoS ONE* **2017**, *12*, 1–14, doi:10.1371/journal.pone.0178888.
57. Levy, S.B. Active efflux mechanisms for antimicrobial resistance. *Antimicrob. Agents Chemother.* **1992**, *36*, 695–703, doi:10.1128/aac.36.4.695.
58. Paulsen, I.T.; Brown, M.H.; Skurray, R.A. Proton-dependent multidrug efflux systems. *Microbiol. Rev.* **1996**, *60*, 575–608.
59. Khameneh, B.; Diab, R.; Ghazvini, K.; Bazzaz, B.S.F. Breakthroughs in bacterial resistance mechanisms and the potential ways to combat them. *Microb. Pathog.* **2016**, *95*, 32–42, doi:10.1016/j.micpath.2016.02.009.
60. Kongkham, B.; Prabakaran, D.; Puttaswamy, H. Opportunities and challenges in managing antibiotic resistance in bacteria using plant secondary metabolites. *Fitoterapia* **2020**, *147*, 104762, doi:10.1016/j.fitote.2020.104762.
61. Walsh, C. Molecular mechanisms that confer antibacterial drug resistance. *Nature* **2000**, *406*, 775–781.
62. Savjani, A.K.G.A.K.T.S.J.K.; Gajjar, A.K.; Savjani, K.T. Mechanisms of Resistance: Useful Tool to Design Antibacterial Agents for Drug—Resistant Bacteria. *Mini Rev. Med. Chem.* **2009**, *9*, 194–205, doi:10.2174/138955709787316038.
63. Fink, A.L. The molecular basis of β -lactamase catalysis and inhibition. *Pharm. Res.* **1985**, *2*, 55–61.
64. Blair, J.M.; Webber, M.A.; Baylay, A.J.; Ogbolu, D.O.; Piddock, L.J.V. Molecular mechanisms of antibiotic resistance. *Nat. Rev. Microbiol.* **2015**, *13*, 42–51, doi:10.1038/nrmicro3380.
65. Olsen, I. New promising β -lactamase inhibitors for clinical use. *Eur. J. Clin. Microbiol. Infect. Dis.* **2015**, *34*, 1303–1308, doi:10.1007/s10096-015-2375-0.
66. Stavri, M.; Piddock, L.J.V.; Gibbons, S. Bacterial efflux pump inhibitors from natural sources. *J. Antimicrob. Chemother.* **2007**, *59*, 1247–1260, doi:10.1093/jac/dkl460.
67. Berkow, E.L.; Angulo, D.; Lockhart, S.R. In Vitro Activity of a Novel Glucan Synthase Inhibitor, SCY-078, against Clinical Isolates of *Candida auris*. *Antimicrob. Agents Chemother.* **2017**, *61*, 1–2, doi:10.1128/aac.00435-17.

68. He, X.; Li, S.; Kaminskyj, S.G.W. Using *Aspergillus nidulans* To Identify Antifungal Drug Resistance Mutations. *Eukaryot. Cell* **2014**, *13*, 288–294, doi:10.1128/ec.00334-13.
69. Cowen, L.; Sanglard, D.; Howard, S.J.; Rogers, P.D.; Perlin, D.S. Mechanisms of Antifungal Drug Resistance. *Cold Spring Harb. Perspect. Med.* **2015**, *5*, 1–2, doi:10.1101/cshperspect.a019752.
70. Cowen, L.E. The evolution of fungal drug resistance: Modulating the trajectory from genotype to phenotype. *Nat. Rev. Microbiol.* **2008**, *6*, 187–198, doi:10.1038/nrmicro1835.
71. Cowen, L.E.; Steinbach, W.J. Stress, Drugs, and Evolution: The Role of Cellular Signaling in Fungal Drug Resistance. *Eukaryot. Cell* **2008**, *7*, 747–764, doi:10.1128/ec.00041-08.
72. Cowen, L.E. Hsp90 Orchestrates Stress Response Signaling Governing Fungal Drug Resistance. *PLOS Pathog.* **2009**, *5*, 1–3, doi:10.1371/journal.ppat.1000471.
73. Cowen, L.E. The fungal Achilles' heel: Targeting Hsp90 to cripple fungal pathogens. *Curr. Opin. Microbiol.* **2013**, *16*, 377–384, doi:10.1016/j.mib.2013.03.005.
74. Naidu, A.; Davidson, P.M. Phyto-phenols. In *Natural Food Antimicrobial Systems*; Naidu, A.S., Ed.; CRC Press: Boca Raton, FL, USA, 2000; pp. 278–307.
75. Burt, S. Essential oils: Their antibacterial properties and potential applications in foods—A review. *Int. J. Food Microbiol.* **2004**, *94*, 223–253, doi:10.1016/j.ijfoodmicro.2004.03.022.
76. Gill, A.O.; Holley, R.A. Mechanisms of Bactericidal Action of Cinnamaldehyde against *Listeria monocytogenes* and of Eugenol against *L. monocytogenes* and *Lactobacillus sakei*. *Appl. Environ. Microbiol.* **2004**, *70*, 5750–5755, doi:10.1128/aem.70.10.5750-5755.2004.
77. Gill, A.O.; Holley, R.A. Disruption of *Escherichia coli*, *Listeria monocytogenes* and *Lactobacillus sakei* cellular membranes by plant oil aromatics. *Int. J. Food Microbiol.* **2006**, *108*, 1–9, doi:10.1016/j.ijfoodmicro.2005.10.009.
78. Negi, P.S. Plant extracts for the control of bacterial growth: Efficacy, stability and safety issues for food application. *Int. J. Food Microbiol.* **2012**, *156*, 7–17, doi:10.1016/j.ijfoodmicro.2012.03.006.
79. Ozfenerci, M.; Caliskan, U.K. Tea tree oil and its use in aromatherapy. *Curr. Pers. Maps.* **2018**, *2*, 90–102.
80. Gallo, F.R.; Palazzino, G.; Federici, E.; Iurilli, R.; Monache, F.D.; Chifundera, K.; Galeffi, C. Oligomeric secoiridoid glucosides from *Jasminum abyssinicum*. *Phytochemistry* **2006**, *67*, 504–510, doi:10.1016/j.phytochem.2005.11.007.
81. Tadiwos, Y.; Nedi, T.; Engidawork, E. Analgesic and anti-inflammatory activities of 80% methanol root extract of *Jasminum abyssinicum* Hochst. ex. Dc. (Oleaceae) in mice. *J. Ethnopharmacol.* **2017**, *202*, 281–289, doi:10.1016/j.jep.2017.02.036.
82. Lakshmanan, P.; Gabriel, J.J. Comparative qualitative analysis of callus extracts of in-vitro and in-vivo plants of *Jasminum angustifolium*, a wild and medicinal plant. *World J. Pharm. Sci.* **2015**, *3*, 1421–1425.
83. Kathiresan, K.; Philip, R. Phytochemical screening and in vitro antioxidant activity of extracts of *Jasminum sessiliflorum*. *Int. J. Pharmacol. Clin. Res.* **2018**, *2*, 117–123.
84. Philip, R.; Krishnasamy, K.; Abraham, E. Evaluation of anti-inflammatory activity of *Jasminum sessiliflorum* extracts. *Int. J. Res. Pharm. Sci.* **2019**, *10*, 2515–2518, doi:10.26452/ijrps.v10i3.1501.
85. Gupta, A.; Chaphalkar, S.R. Use of flow cytometry to measure the immunostimulatory activity of aqueous extract of *Jasminum auriculatum*. *Int. J. Curr. Adv. Res.* **2015**, *4*, 87–91.
86. Bahuguna, Y.; Juyal, V.; Rawat, M.S.M.; Jalalpure, S. Diuretic activity of flowers of *Jasminum auriculatum* Vahl. *J. Pharm. Res.* **2009**, *2*, 215–216.
87. Rastogi, R.P.; Mehrotra, B.N.; Sinha, S.; Pant, P.; Seth, R. *Compendium of Indian Medicinal Plants*; Central Drug Research Institute: Lucknow, India, 2001; pp. 395–396.
88. Arivoli, S.; Divya, S.; Arumugam, B.; Meeran, M.; Jayakumar, M.; Raveen, R.; Samuel, T. Phytochemical constituents of *Jasminum fluminense* Linnaeus (Oleaceae): An additional tool in the ecofriendly management of mosquitoes. *J. Pharmacog. Phytochem.* **2018**, *7*, 548–556.
89. Prajapati, N.D.; Purohit, S.S.; Sharma, A.K.; Kumar, T. *A Handbook of Medicinal Plants: A Complete Source Book*; Agrobios: Jodhpur, India, 2003; pp. 554–554.
90. Zhao, G.-Q.; Yin, Z.-F.; Liu, Y.-C.; Li, H.-B. Iridoid glycosides from buds of *Jasminum officinale* L. var. *grandiflorum*. *Yao Xue Xue Bao Acta Pharm. Sin.* **2011**, *46*, 1221–1224.
91. Zhao, G.-Q.; Xia, J.-J.; Dong, J.-X. Glycosides from flowers of *Jasminum officinale* L. var. *grandiflorum*. *Yao Xue Xue Bao Acta Pharm. Sin.* **2007**, *42*, 1066–1069.
92. Singh, B.; Sharma, R.A. *Secondary Metabolites of Medicinal Plants, 4 Volume Set: Ethnopharmacological Properties, Biological Activity and Production Strategies*; John Wiley & Sons: Hoboken, NJ, USA, 2020; pp. 574–584.
93. Dubey, P.; Tiwari, A.; Gupta, S.K.; Watal, G. Phytochemical and biochemical studies of *Jasminum officinale* leaves. *Int. J. Pharm. Sci. Res.* **2016**, *7*, 2632–2640.
94. El-Hawary, S.S.; El-Hefnawy, H.M.; Osman, S.M.; El-Raey, M.A.; Ali, F.A.M. Phenolic profiling of different *Jasminum* species cultivated in Egypt and their antioxidant activity. *Nat. Prod. Res.* **2019**, 1–6, doi:10.1080/14786419.2019.1700508.
95. Lu, Y.; Han, Z.-Z.; Zhang, C.-G.; Ye, Z.; Wu, L.-L.; Xu, H. Four new sesquiterpenoids with anti-inflammatory activity from the stems of *Jasminum officinale*. *Fitoterapia* **2019**, *135*, 22–26, doi:10.1016/j.fitote.2019.03.029.
96. Tauchen, J.; Dösköc, I.; Caffi, C.; Lulekal, E.; Marsik, P.; Havlik, J.; van Damme, P.; Kokoska, L. In vitro antioxidant and anti-proliferative activity of Ethiopian medicinal plant extracts. *Ind. Crop. Prod.* **2015**, *74*, 671–679, doi:10.1016/j.indcrop.2015.05.068.

97. Bhagath, K.; Kekuda, P.T.R.; Raghavendra, H.L.; Swarnalatha, S.P.; Preethi, H.R.; Surabhi, K.S. In vitro antioxidant and anthelmintic activity of extracts of *Jasminum arborescens* (Roxb.). *Int. J. Drug. Dev. Res.* **2010**, *2*, 89–95.
98. Ferreres, F.; Grosso, A.C.; Gil-Izquierdo, A.; Valentão, P.; Andrade, P.B. Assessing *Jasminum grandiflorum* L. authenticity by HPLC-DAD-ESI/MSn and effects on physiological enzymes and oxidative species. *J. Pharm. Biomed. Anal.* **2014**, *88*, 157–161, doi:10.1016/j.jpba.2013.08.040.
99. Umamaheswari, M.; Asokkumar, K.; Rathidevi, R.; Sivashanmugam, A.T.; Subhadradevi, V.; Ravi, T.K. Antiulcer and in vitro antioxidant activities of *Jasminum grandiflorum* L. *J. Ethnopharmacol.* **2007**, *110*, 464–470, doi:10.1016/j.jep.2006.10.017.
100. Chaturvedi, A.P.; Tripathi, Y.B. Methanolic extract of leaves of *Jasminum grandiflorum* Linn modulates oxidative stress and inflammatory mediators. *Inflammopharmacology* **2011**, *19*, 273–281, doi:10.1007/s10787-011-0087-3.
101. Dessai, P.; Sawant, R.P. In-vitro pharmacological activities of *Jasminum malabaricum* Wight. *J. Glob. Trends Pharm. Sci.* **2018**, *9*, 5076–5082.
102. Poonia, P.; Niazi, J.; Chaudhary, G.; Kalia, A.N. In vitro antioxidant potential of *Jasminum mesnyi* Hance (Leaves) extracts. *Res. J. Pharm. Biol. Chem. Sci.* **2011**, *2*, 348–357.
103. Borar, S.; Punia, P.; Kalia, A.N. Antioxidant potential of n-butanol fraction from extract of *Jasminum mesnyi* Hance leaves. *Indian J. Exp. Boil.* **2011**, *49*, 39–43.
104. Guo, Z.-Y.; Li, P.; Huang, W.; Wang, J.-J.; Liu, Y.-J.; Liu, B.; Wang, Y.-L.; Wu, S.-B.; Kennelly, E.J.; Long, C.-L. Antioxidant and anti-inflammatory caffeoyl phenylpropanoid and secoiridoid glycosides from *Jasminum nervosum* stems, a Chinese folk medicine. *Phytochemistry* **2014**, *106*, 124–133, doi:10.1016/j.phytochem.2014.07.011.
105. Li, A.-N.; Li, S.; Li, H.-B.; Xu, D.-P.; Xu, X.-R.; Chen, F. Total phenolic contents and antioxidant capacities of 51 edible and wild flowers. *J. Funct. Foods* **2014**, *6*, 319–330, doi:10.1016/j.jff.2013.10.022.
106. Khidzir, K.M.; Cheng, S.-F.; Chuah, C.-H. Interspecies variation of chemical constituents and antioxidant capacity of extracts from *Jasminum sambac* and *Jasminum multiflorum* grown in Malaysia. *Ind. Crop. Prod.* **2015**, *74*, 635–641, doi:10.1016/j.indcrop.2015.05.053.
107. He, F.; Zuo, L. Redox Roles of Reactive Oxygen Species in Cardiovascular Diseases. *Int. J. Mol. Sci.* **2015**, *16*, 27770–27780, doi:10.3390/ijms161126059.
108. Dias, V.; Junn, E.; Mouradian, M.M. The role of oxidative stress in Parkinson's disease. *J. Parkinson's Dis.* **2013**, *3*, 461–491.
109. Zuo, L.; Zhou, T.; Pannell, B.K.; Ziegler, A.C.; Best, T.M. Biological and physiological role of reactive oxygen species—The good, the bad and the ugly. *Acta Physiol.* **2015**, *214*, 329–348, doi:10.1111/apha.12515.
110. Tan, B.L.; Norhaizan, M.E.; Huynh, K.; Heshu, S.R.; Yeap, S.K.; Hazilawati, H.; Roselina, K. Water extract of brewers' rice induces apoptosis in human colorectal cancer cells via activation of caspase-3 and caspase-8 and downregulates the Wnt/ β -catenin downstream signaling pathway in brewers' rice-treated rats with azoxymethane-induced colon carcinogenesis. *BMC Complement. Altern. Med.* **2015**, *15*, 1–14, doi:10.1186/s12906-015-0730-4.
111. Liu, Z.; Zhou, T.; Ziegler, A.C.; Dimitrion, P.; Zuo, L. Oxidative Stress in Neurodegenerative Diseases: From Molecular Mechanisms to Clinical Applications. *Oxidative Med. Cell. Longev.* **2017**, *2017*, 1–11, doi:10.1155/2017/2525967.
112. Hercberg, S.; Galan, P.; Preziosi, P.; Bertrais, S.; Mennen, L.; Malvy, D.; Roussel, A.M.; Favier, A.; Briançon, S. The SU. VI. MAX Study: A randomized, placebo-controlled trial of the health effects of antioxidant vitamins and minerals. *Arch. Intern. Med.* **2004**, *164*, 2335–2342.
113. Halliwell, B.; Rafter, J.; Jenner, A. Health promotion by flavonoids, tocopherols, tocotrienols, and other phenols: Direct or indirect effects? Antioxidant or not? *Am. J. Clin. Nutr.* **2005**, *81*, 268S–276S, doi:10.1093/ajcn/81.1.268s.
114. Dysken, M.W.; Sano, M.; Asthana, S.; Vertrees, J.E.; Pallaki, M.; Llorente, M.; Love, S.; Schellenberg, G.D.; McCarten, J.R.; Malphurs, J.; et al. Effect of vitamin E and memantine on functional decline in Alzheimer disease: The TEAM-AD VA cooperative randomized trial. *Jama* **2014**, *311*, 33–44.
115. Yuan, G.; Sun, B.; Yuan, J.; Wang, Q. Effect of 1-methylcyclopropene on shelf life, visual quality, antioxidant enzymes and health-promoting compounds in broccoli florets. *Food Chem.* **2010**, *118*, 774–781, doi:10.1016/j.foodchem.2009.05.062.
116. Kolanjiappan, K.; Manoharan, S. Chemopreventive efficacy and anti-lipid peroxidative potential of *Jasminum grandiflorum* Linn. on 7,12-dimethylbenz(a)anthracene-induced rat mammary carcinogenesis. *Fundam. Clin. Pharmacol.* **2005**, *19*, 687–693, doi:10.1111/j.1472-8206.2005.00376.x.
117. Chaturvedi, A.P.; Kumar, M.; Tripathi, Y.B. Efficacy of *Jasminum grandiflorum* L. leaf extract on dermal wound healing in rats. *Int. Wound J.* **2012**, *10*, 675–682, doi:10.1111/j.1742-481x.2012.01043.x.
118. Sengar, N.; Joshi, A.; Prasad, S.K.; Hemalatha, S. Anti-inflammatory, analgesic and anti-pyretic activities of standardized root extract of *Jasminum sambac*. *J. Ethnopharmacol.* **2015**, *160*, 140–148, doi:10.1016/j.jep.2014.11.039.
119. Ho, C.C.; Ng, S.C.; Chuang, H.L.; Wen, S.Y.; Kuo, C.H.; Mahalakshmi, B.; Huang, C.Y.; Kuo, W.W. Extracts of *Jasminum sambac* flowers fermented by *Lactobacillus rhamnosus* inhibit H₂O₂- and UVB-induced aging in human dermal fibroblasts. *Environ. Toxicol.* **2020**, *36*, 607–619.
120. Abdoul-Latif, F.; Edou, P.; Eba, F.; Mohamed, N.; Ali, A.; Djama, S.; Obame, L.; Bassolé, I.; Dicko, M. Antimicrobial and antioxidant activities of essential oil and methanol extract of *Jasminum sambac* from Djibouti. *Afr. J. Plant Sci.* **2010**, *4*, 38–43.
121. AlRashdi, A.S.; Salama, S.M.; Alkiyumi, S.S.; Abdulla, M.A.; Hadi, A.H.A.; Abdelwahab, S.I.; Taha, M.M.; Hussiani, J.; Asykin, N. Mechanisms of gastroprotective effects of ethanolic leaf extract of *Jasminum sambac* against HCl/ethanol-induced gastric mucosal injury in rats. *Evid. Based Compl. Alt. Med.* **2012**, *2012*, 1–15.

122. Bhagat, A.D.; Khairnar, A.U.; Tenpe, C.R.; Upaganalwar, A.B.; Yeole, P.G. Anti-inflammatory activity of *Jasminum sambac* leaf extract against carrageenan induced rat paw edema. *Indian J. Nat. Prod.* **2007**, *23*, 25–28.
123. Rahman, M.A.; Hasan, M.; Hossain, S.M.A.; Biswas, N.N. Analgesic and cytotoxic activities of *Jasminum sambac* (L.) Aiton. *Pharmacologyonline* **2011**, *1*, 124–131.
124. Kumar, M.; Randhava, N.K. *Jasminum mesnyi* Hance: Review at a Glance. *J. Drug Deliv. Ther.* **2014**, *4*, 44–47, doi:10.22270/jddt.v4i5.935.
125. Bhushan, B.; Sardana, S.; Bansal, G. Phytochemical and pharmacognostical studies of leaves of *Jasminum mesnyi* Hance. *J. Chem. Pharma. Res.* **2015**, *7*, 922–926.
126. Kumaresan, M.; Kannan, M.; Sankari, A.; Chandrasekhar, C.N.; Vasanthi, D. Phytochemical screening and antioxidant activity of *Jasminum multiflorum* (pink Kakada) leaves and flowers. *J. Pharmacog. Phytochem.* **2019**, *8*, 1168–1173.
127. Jain, A.; Sharma, R.; Kumar, A.; Sharma, S. *Jasminum* species: An overview. *Int. J. Inst. Pharm. Life Sci.* **2011**, *1*, 251–266.
128. Shekhar, S.; Prasad, M.P. Evaluation of antimicrobial activity of *Jasminum* species using solvent extracts against clinical pathogens. *World J. Pharm. Pharm. Sci.* **2015**, *4*, 1247–1256.
129. Yuniarto, A.; Kurnia, I.; Ramadhan, M. Anti-obesity effect of ethanolic extract of jasmine flowers (*Jasminum sambac* (L.) Ait.) in high fat diet induced mice: Potent inhibitor of pancreatic lipase enzyme. *Int. J. Adv. Phar. Biol. Chem.* **2015**, *4*, 18–22.
130. Dhalaria, R.; Verma, R.; Kumar, D.; Puri, S.; Tapwal, A.; Kumar, V.; Nepovimova, E.; Kuca, K. Bioactive Compounds of Edible Fruits with Their Anti-Aging Properties: A Comprehensive Review to Prolong Human Life. *Antioxidants* **2020**, *9*, 1123, doi:10.3390/antiox9111123.
131. Nagmoti, D.M.; Khatri, D.K.; Juvekar, P.R.; Juvekar, A.R. Antioxidant activity free radical-scavenging potential of *Pithecellobium dulce* Benth seed extracts. *Free Radic. Antioxid.* **2012**, *2*, 37–43, doi:10.5530/ax.2012.2.2.7.
132. Sharma, A.; Bernatchez, P.N.; de Haan, J.B. Targeting Endothelial Dysfunction in Vascular Complications Associated with Diabetes. *Int. J. Vasc. Med.* **2012**, *2012*, 1–12, doi:10.1155/2012/750126.
133. Boora, F.; Chirisa, E.; Mukanganyama, S. Evaluation of Nitrite Radical Scavenging Properties of Selected Zimbabwean Plant Extracts and Their Phytoconstituents. *J. Food Process.* **2014**, *2014*, 1–7, doi:10.1155/2014/918018.
134. Tehrani, H.S.; Moosavi-Movahedi, A.A. Catalase and its mysteries. *Prog. Biophys. Mol. Biol.* **2018**, *140*, 5–12, doi:10.1016/j.pbiomolbio.2018.03.001.
135. Rakotoarisoa, M.; Angelov, B.; Espinoza, S.; Khakurel, K.; Bizien, T.; Angelova, A. Cubic Liquid Crystalline Nanostructures Involving Catalase and Curcumin: BioSAXS Study and Catalase Peroxidatic Function after Cubosomal Nanoparticle Treatment of Differentiated SH-SY5Y Cells. *Molecules* **2019**, *24*, 1–21, doi:10.3390/molecules24173058.
136. Lobo, V.; Patil, A.; Phatak, A.; Chandra, N. Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacogn. Rev.* **2010**, *4*, 118–126, doi:10.4103/0973-7847.70902.
137. Devasagayam, T.P.A.; Tilak, J.C.; Boloor, K.K.; Sane, K.S.; Ghaskadbi, S.S.; Lele, R.D. Free radicals and antioxidants in human health: Current status and future prospects. *J. Assoc. Physicians India* **2004**, *52*, 794–804.
138. Wills, E.D. Effects of lipid peroxidation on membrane-bound enzymes of the endoplasmic reticulum. *Biochem. J.* **1971**, *123*, 983–991, doi:10.1042/bj1230983.
139. Farooqui, A.A.; Horrocks, L.A. Lipid Peroxides in the Free Radical Pathophysiology of Brain Diseases. *Cell. Mol. Neurobiol.* **1998**, *18*, 599–608, doi:10.1023/a:1020625717298.
140. Cheeseman, K. Mechanisms and effects of lipid peroxidation. *Mol. Asp. Med.* **1993**, *14*, 191–197, doi:10.1016/0098-2997(93)90005-x.
141. Yu, B.P.; Suescun, E.A.; Yang, S.Y. Effect of age-related lipid peroxidation on membrane fluidity and phospholipase A2: Modulation by dietary restriction. *Mech. Ageing Dev.* **1992**, *65*, 17–33, doi:10.1016/0047-6374(92)90123-u.
142. Sies, H.; Sharov, V.S.; Klotz, L.O.; Briviba, K. Glutathione peroxidase protects against peroxynitrite-mediated oxidations: A new function for selenoproteins as peroxynitrite reductase. *J. Biol. Chem.* **1997**, *272*, 27812–27817.
143. Szabó, C.; Ischiropoulos, H.; Radi, R. Peroxynitrite: Biochemistry, pathophysiology and development of therapeutics. *Nat. Rev. Drug Discov.* **2007**, *6*, 662–680, doi:10.1038/nrd2222.