

Article

# Edible Flower Species as a Promising Source of Specialized Metabolites

Mia Dujmović <sup>1</sup>, Sanja Radman <sup>2,\*</sup>, Nevena Opačić <sup>2</sup>, Sanja Fabek Uher <sup>2</sup>, Vida Mikuličin <sup>1</sup>, Sandra Voća <sup>1</sup> and Jana Šic Žlabur <sup>1</sup>

<sup>1</sup> Department of Agricultural Technology, Storage and Transport, University of Zagreb Faculty of Agriculture, Svetosimunska cesta 25, 10000 Zagreb, Croatia

<sup>2</sup> Department of Vegetable Crops, University of Zagreb Faculty of Agriculture, Svetosimunska cesta 25, 10000 Zagreb, Croatia

\* Correspondence: sradman@agr.hr

**Table S1.** Flower species used in study

| Picture of flower species   | Scientific name                 | Common name and<br>synonyms                        | Family        |
|---|---------------------------------|--|---------------|
|    | <i>Calendula officinalis</i> L. | Common marigold<br>Pot marigold<br>Calendula       | Asteraceae    |
|    | <i>Tagetes erecta</i> L.        | African marigold<br>Aztec marigold<br>Big marigold | Asteraceae    |
|   | <i>Tropaeolum majus</i> L.      | Nasturtium<br>Indian cress<br>Monk's cress         | Tropaeolaceae |
|  | <i>Cucurbita pepo</i> L.        | Zucchini<br>Pumpkin                                | Cucurbitaceae |
|  | <i>Centaurea cyanus</i> L.      | Cornflower<br>Bachelor's button                    | Asteraceae    |