

Suppressive effects of flavonoids on macrophage-associated adipocyte inflammation in a differentiated murine preadipocyte 3T3-L1 cells co-cultured with a murine macrophage RAW264.7 cells

Dahae Lee ¹, Sukyong Hong ², Kiwon Jung ^{2,3}, Sung youl Choi ^{4,*}, Ki Sung Kang ^{1,*}

¹Department of Preventive Medicine, College of Korean Medicine, Gachon University, Seongnam 13120, Korea; pjsldh@gachon.ac.kr (D. L.); kkang@gachon.ac.kr (K. S. K.)

²College of Pharmacy, CHA University, Sunghnam 13844, Republic of Korea; hongsk@chauniv.ac.kr (S. H.)

³Oncobix Co., Ltd., Yongin-si 16950, Gyeonggi-do, Republic of Korea; pharmj@oncobix.com (K. J.)

⁴Department of Neuropsychiatry, College of Korean Medicine, Gachon University

*Correspondence: pc1075@gachon.ac.kr (S. C.), Tel.: +82-32-770-1343 (S. C.); kkang@gachon.ac.kr (K. S. K.), Tel.: +82-31-750-5402




Figure S1. Effect of the flavonoids on the viability of 3T3-L1 preadipocytes and RAW264.7 cells. (A and B) Effect of flavonoids compared with the control (0 μM) on the viability of 3T3-L1 preadipocytes for 24 h. Cell viability was determined by the MTT assay. In individual experiments, each treatment condition was set up in quadruplicate, and each experiment was repeated 3 times independently ($p < 0.05$ compared to untreated cells, Kruskal–Wallis non-parametric test). Data are represented as the mean \pm SEM.