

***Tetracera loureiri* extract regulates lipopolysaccharide-induced inflammatory response via nuclear factor-κB and mitogen activated protein kinase signaling pathway**

Jung A Lee^{1, †}, Ju Young Shin^{2, †}, Seong Su Hong¹, Young-Rak Cho¹, Ju-Hyoung Park², Dong-Wan Seo², Joa Sub Oh², Jae-Shin Kang³, Jae Ho Lee³ and Eun-Kyung Ahn^{1,*}

¹*Bio-Center, Gyeonggido Business & Science Accelerator (GBSA), Suwon 16229, Republic of Korea.*

²*College of Pharmacy, Dankook University, Cheonan 31116, Republic of Korea.*

³*Biological Genetic Resources Utilization Division, National Institute of Biological Resources, Incheon 22689, Republic of Korea.*

SUPPORTING INFORMATION

List of Supporting Information	Page
Figure S1. The ^1H NMR spectrum of 1	
S	3
Figure S2. The ^{13}C NMR spectrum of 1	
S	3
Figure S3. ESI-MS spectrum of 1	
S	4
Figure S4. The ^1H NMR spectrum of 2	
S	5
Figure S5. The ^{13}C NMR spectrum of 2	
S	5
Figure S6. ESI-MS spectrum of 2	
S	6

Figure S1.




Figure S2.




Figure S3.






Figure S4.**Figure S5.**

Figure S6.

