



Article In Vitro Studies on the Antimicrobial and Antioxidant Activities of Total Polyphenol Content of *Cynara humilis* from Moulay Yacoub Area (Morocco)

Mostafa El Khomsi ^{1,*}, Mohammed Kara ^{2,*}, Anouar Hmamou ³, Amine Assouguem ⁴, Omkulthom Al Kamaly ⁵, Asmaa Saleh ⁵, Sezai Ercisli ⁶, Hafize Fidan ⁷ and Driss Hmouni ¹

- ¹ Natural Resources and Sustainable Development Laboratory, Department of Biology, Faculty of Sciences, Ibn Tofail University, B.P.133, Kenitra 14000, Morocco; Hmouni.driss@uit.ac.ma
- ² Laboratory of Biotechnology, Conservation and Valorisation of Natural Resources (LBCVNR), Faculty of Sciences Dhar El Mehraz, Sidi Mohamed Ben Abdellah University, BP.1796 Atlas, Fez 30000, Morocco
- ³ Laboratory of Engineering, Molecular Organometallic Materials and Environment, Faculty of Sciences Dhar El Mehraz, Sidi Mohamed Ben Abdellah University, BP.1796 Atlas, Fez 30000, Morocco; anouar.hmamou@usmba.ac.ma
- ⁴ Laboratory of Functional Ecology and Environment, Faculty of Sciences and Technology, Sidi Mohamed Ben Abdellah University, B.O. Box 2202 Imouzzer Street, Fez 30000, Morocco; assougam@gmail.com
- ⁵ Department of Pharmaceutical Sciences, College of Pharmacy, Princess Nourah bint Abdulrahman University, P.O. Box 84428, Riyadh 11671, Saudi Arabia; omalkmali@pnu.edu.sa (O.A.K.); ASAli@pnu.edu.sa (A.S.)
- ⁶ Department of Horticulture, Atatürk University, 25240 Erzurum, Turkey; sercisli@gmail.com
- ⁷ Department of Tourism and Culinary Management, Faculty of Economics, University of Food Technologies, 4000 Plovdiv, Bulgaria; hfidan@abv.bg
- * Correspondence: elkhomsi.mostafa@uit.ac.ma (M.E.K.); mohammed.kara@usmba.ac.ma (M.K.)

Abstract: In Morocco, Cynara humilis L. is used in traditional medicine. The objective of this research was to research the antioxidant and antimicrobial properties of hydroethanolic extracts from the C. humilis plant's leaves and roots. The content of polyphenols and flavonoids was evaluated using Folin-Ciocalteu's and aluminum chloride assays. Two techniques were used to evaluate antioxidant properties: antioxidant capacity in total (TAC) and 2,2-diphenyl-1-picrylhdrazyl (DPPH). In antimicrobial assays, five pathogenic microbial strains were studied including two Escherichia coli, one coagulase-negative Staphylococcus and Klebsiella pneumoniae, and one Candida albicans, by two techniques: agar disk diffusion and microdilution. Leaves had a greater content of flavonoids 27.07 mg QE/g of extract and the polyphenols 38.84 mg GAE/g of extract than root 24.39 mg QE/g of extract and 29.39 mg GAE/g of extract, respectively. The TAC test value of the 0.77 mg AAE/g extract in the leaf extract was found to be significantly greater than that of the 0.60 mg EAA/g extract in the root extract. The DPPH antioxidant assay IC_{50} values of the root and leaf extract were 0.23 and 0.93 µg/mL, respectively. C. humilis extracts showed an antimicrobial effect against all tested strains, the inhibitory zone (DIZ) have values in the range between 12 and 15 mm. Moreover, the root extract showed the lowest minimum inhibitory concentration (MIC) against coagulase-negative *Staphylococcus* with an IC_{50} value of 6.25 mg/mL. The higher content of flavonoids and polyphenols in the hydroethanolic extracts of *C. humilis* leaves and roots demonstrates that they have a significant antimicrobial and antioxidant effect, as found in this study.

Keywords: *Cynara humilis* L.; antimicrobial activity; antioxidant activity; in vitro; total polyphenols; 2.2-diphenyl-1-picrylhdrazyl

1. Introduction

The therapeutic value of medicinal and aromatic plants has long been recognized, and the concept that plants contain pharmacologically active substances has been well defined. Plants have been used to cure disorders of microbial origin as well other disorders since



Citation: El Khomsi, M.; Kara, M.; Hmamou, A.; Assouguem, A.; Al Kamaly, O.; Saleh, A.; Ercisli, S.; Fidan, H.; Hmouni, D. In Vitro Studies on the Antimicrobial and Antioxidant Activities of Total Polyphenol Content of *Cynara humilis* from Moulay Yacoub Area (Morocco). *Plants* **2022**, *11*, 1200. https:// doi.org/10.3390/plants11091200

Academic Editor: Gloria Castellano

Received: 1 April 2022 Accepted: 20 April 2022 Published: 29 April 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). ancient times. The first healers were priests or doctors. Indeed, science and religion have been associated with illness and healing for more than 20,000 years, and their goal has always been to heal the soul and the body [1–5]. A large number of active ingredients used in the preparation of medicines are of plant origin [6,7]. Indeed, scientific research has documented that the extracts of medicinal and aromatic plants and essential oils coming from many plants have medicinal properties that can be used in the phytotherapy, especially antibacterial, antioxidant, antifungal and also pharmacological properties [8–10].

Antioxidants used to fight free radicals can also be used to cure of certain illnesses, including heart problems, diabetes, neurodegenerative diseases and also cancer [11,12]. The use of synthetic antioxidants can be the cause of many diseases, such as carcinogenicity, which makes peoples tend to use antioxidants of natural origin, especially of plant origin, which represents a very important source of natural antioxidants [13]. The antimicrobial activity of natural antioxidants is well reported in the literature, and many natural chemicals of plant origin have been shown to have antimicrobial activities [14–16]. The biological compounds responsible for these properties are the secondary metabolites present in plants, such as polyphenols, flavonoids, terpenes, coumarins, tannins, and alkaloids [17–19]. For this reason, scientific research must be applied to the search for new plants to study their phytochemical composition and also their biological and pharmacological activities [20].

Cynara humilis L. is a tiny plant from the Mediterranean region and belongs to the Asteraceae family [21]. Based on genetic and morphological data, *C. humilis* was part of the secondary gene pool of *C. cardunculus* [21–23]. In Morocco, the roots of *Cynara humilis* L are traditionally used to treat wounds and burns [1,24,25]. The healing effect of the plant is well demonstrated in previous research [26]. In addition, because of its choleretic activities, the plant is used in therapy to increase bile secretion [27]. Other studies have indicated that *C. humilis* has been used in cheese making [28,29]. Phytochemical studies of methanolic extracts of the *C. humilis* plant have documented the presence of phenolic compounds, flavonoids, coumarins, and aurone [27]. Other research has documented that the biological activities of the plant have not yet studied [26].

However, no study has documented the plant's antimicrobial and antioxidant activities to our knowledge, and scientific research on this plant is few and limited. Within this framework, our research is the first to delve into the antioxidant and antibacterial biological characteristics of *C. humilis* root and leaf extracts, and also to evaluate its total polyphenol and flavonoid composition.

2. Results and Discussion

2.1. Yield of Extraction

The yield of extraction of the roots and leaves of *C. humilis* showed that the extract of the leaves presented a yield 17.4% higher than that of the extract of the roots, which presented a yield of 12.8% (Table 1).

Table 1. Yield obtained from the extraction of roots and leaves of C. humilis.

Extract	Mass of Plant Powder Used (g)	Mass of the Crude Extract Obtained (g)	Yield of Extraction (%)		
Leaf	50	8.7	17.4		
Root	50	6.4	12.8		

2.2. Phenolic Compounds and Flavonoid Contents

Polyphenols and flavonoids are types of secondary metabolites found in almost all plants, and their quantity varies depending on geographical origin, species, and variety [30–32]. The characteristic structure of each of these aforementioned compounds indicates a specific behavior/action/role, as statistically demonstrated in [33]. It is difficult to judge the global composition of polyphenols in a single study carried out in spring on a single site. The results of the polyphenol assay are written in unit (mg GAE/g of extract) which means milligrams of gallic acid equivalent by gram of extract) (Figure 1). The two extracts tested, leaves and roots, showed a statistically significant difference (p = 0.04). Leaves represented a value of $38.84 \pm 5.15 \text{ mg GAE/g}$ extract, compared to roots with $29.56 \pm 1.44 \text{ mg GAE/g}$ extract. When compared to the findings of previous investigations, our results show that the quantity of polyphenols in *C. humilis* is higher than that of *C. cardunculus*, which varied between 6.96 and 14.79 mg GAE/g extract [34], and lower than that of *C. scolymus*, at 50 mg GAE/g extract [35]. Several environmental factors, including climatic conditions, geographical location, soil fertility, cultivar genotype, and experimental factors, including the part of the plant used, the time of harvesting, the extraction method, as well as the polarity of the solution being used, and extraction time, can influence phenolic compounds in plants [36–39]. HPLC chromatographic analyses have shown that the *C. humilis* plant contains numerous phenolic compounds including isochlorogenic acid, chlorogenic acid and caffeic acid [27].



Figure 1. Histogram of flavonoid contents (TFC) and polyphenol content (TPC) in the *C. humilis* extracts. The significant difference (p < 0.05) between the tested samples is shown by the two tetters a and b.

The quantity of flavonoids in plant extracts was evaluated using the quercetin calibration curve, y = 0.0062 + 0.4289 and $R^2 = 0.857$, flavonoid results presented by the unit (mg QE/g of extract) which means milligrams of quercetin equivalents by gram of extract. The flavonoid values in the leaf extract (27.07 ± 2.79 mg QE/g of extract) are superior to those found in the root extract (24.39 ± 2.87 mg QE/g extract). The analysis of these finding revealed no statistically significant differences (p = 0.309) (Figure 1). In the literature, the biological activities, especially antioxidant and antibacterial properties, have been widely documented [14,40]. A phytochemical study indicated the presence of the following flavonoids, luteolin, apigenin, quercetin, maritimen in the *C. humilis* plant [27].

2.3. Antioxidant Assay

2.3.1. Antioxidant Capacity Total (TAC)

An ascorbic acid calibration curve was created to determine the overall antioxidant capacity, and the value were represented by the unit (mg EAA/g of extract) which means milligrams of ascorbic acid equivalent by gram of extract. the leaf extract has 0.77 ± 0.07 mg AAE/g extract total antioxidant capacity, is more effective than the root extract, which had a TAC of 0.60 ± 0.04 mg AAE/g of extract, as per TAC test findings (Table 2). The difference of the two extracts analyzed was found to be significant (p = 0.023) in the results analysis.

Extract	ТРС	TFC	IC ₅₀	TAC	E. coli ATCC25922	<i>E. coli</i> CIP 53126	K. pneumoniae	Staphylococcus	C. albicans
Leaf	32.93	26.74	0.23	0.77	13	12	12	12	14
Leaf	42.33	30.01	0.21	0.84	13	13	12	12	14
Leaf	41.27	24.47	0.25	0.7	12	13	13	13	14
Root	27.94	26.66	0.94	0.63	12	15	12	15	14
Root	30.03	25.34	0.92	0.55	12	15	12	16	14
Root	30.71	21.16	0.95	0.62	12	15	12	14	16

Table 2. The raw data of the analyses carried out on the *C. humilis* plant with three repetitions.

2.3.2. DPPH Free Radical Scavenging Antioxidant Activity

Polyphenols and flavonoids are abundant in the *C. humilis* plant, and these compounds are renowned for their antioxidant properties, scavenging of free radical by inhibiting the traces of metal ions that participate in their formation by exchange of proton atoms [30]. The IC₅₀ value is the necessary concentration to lower DPPH by 50%. According to Table 3, the IC₅₀ values of the investigated substances differed by a significant amount (p < 0.05). The results obtained indicate that leaves and roots of the plant had an inhibitory concentration of 0.23 ± 0.02 and $0.93 \pm 0.01 \,\mu$ g/mL, respectively, which is higher than those of the reference antioxidants; quercetin, BHT, and ascorbic acid had a concentration of 0.05, 0.20, and $0.16 \pm 0.01 \,\mu$ g/mL, respectively. As a lower IC₅₀ value indicates a higher antioxidant capacity, this shows that both extracts had lower antioxidant activity than the reference antioxidants. The phenolic content among those sample explains the antioxidant properties of the plant, which intervenes in the neutralization of the free radicals of the oxidizing agents used [41,42]. The antioxidant action of flavonoids is well understood, flavonoids scavenging free radicals through chelation of trace metal ions, iron chelation, and oxidative stress reduction [30,43,44].

Extract	IC ₅₀ (µg/mL)
Leaf	0.23 $^{\mathrm{B}}\pm0.02$
Root	0.93 $^{ m A}\pm 0.01$
BHT	0.20 $^{ m C}\pm$ 0.00
Quercetin	0.05 $^{ m E}\pm$ 0.00
Ascorbic acid	$0.16 ^{\mathrm{D}} \pm 0.01$

Table 3. DPPH antioxidant test result represented by IC_{50} .

The letters A-E indicate that there is a statistical difference (p < 0.05) between the tested samples.

In comparison with other species of *Cynara* spp., our results show that the leaves of the *C. humilis* plant have a higher DPPH scavenging capacity than the leaves of *C. cardunculus*, which showed a value of 50 µg/mL [34]. The following compounds were found in the phytochemical analysis of the *C. humilis* plant: chlorogenic acid, caffeic acid, isochlorogenic acid, quercetin, luteolin, maritimen, apigenin [27]. Quercetin is a powerful antioxidant with a high scavenging capacity for DPPH free radicals [45]. Apigenin and luteolin, as well as their glycosides, have been seen to be potent antioxidants in other research [46]. The antioxidant activities of luteolin and luteolin 7-glucoside, have previously been reported against DPPH free radical scavenging [46,47].

Phenolic acids are known for their antioxidant effect, including chlorogenic acid. These antioxidant properties are almost the same as those o 4-O-caffeoylquinic acid and 3-O-caffeolquinicf when analyzing the radical scavenging of superoxide anionic radical and ability to inhibit of the oxidation of methyl linoleate [48].

2.4. Antimicrobial Activity Disc Inhibitory Assay

To study their antimicrobial action, five bacterial strains were tested using a solid medium disc. The inhibitory zone (DIZ) was between 12 and 15 mm in diameter (Table 4). The highest inhibition zone was recorded in two strains, E. coli CIP 53126 and coagulasenegative Staphylococcus, in the root extract of 15.00 mm, followed by C. albicans ATCC 10231 which recorded a DIZ of 14.67 \pm 1.15 mm in the root extract and 14.00 mm in the leaf extract. The lowest DIZ was recorded in the two strains E. coli ATCC 25922 and K. pneumoniae susceptible in the root extract of 12.00 mm. Statistical analysis of the inhibition zone diameter values of the two *C. humilis* extracts was significant (p < 0.05) against coagulase-negative *Staphylococcus* and *E. coli* CIP 53126, and not significant (p > 0.05) against the three remaining strains C. albicans ATCC 10231, K. pneumoniae, and E. coli ATCC 25922. Compared to the standard antibiotics used streptomycin, fluconazole and ampicillin, Coagulase negative *Staphylococcus* recorded a DIZ of 9.61 \pm 0.20 mm in the presence of streptomycin, while the other bacterial strains showed resistance to the streptomycin and ampicillin. For *C. albicans* recorded a zone of inhibition of 21.20 ± 4.20 mm in the presence of fluconazole. In comparison with other species of Cynara spp., Salmonella typhimurium, *Bacillus subtilis, S. epidermidis,* and *E. coli* were all found to have antibacterial properties in C. cardunculus. On the other hand, C. scolymus showed antimicrobial effect against S. aureus, B. subtilis, E. coli, Agrobacterium tumefaciens, Micrococcus luteus, Pseudomonas aeruginosa, and C. albicans [46,49].

The Zones of Inhibition's Diameter (mm) Strains of Bacteria Leaf Extract **Root Extract** Ampicillin Streptomycin Fluconazole $12.67 \text{ }^{\text{B}} \pm 0.57$ $15.00\ ^{\rm A}\pm 0.00$ E. coli CIP 53126 R R $12.67 \ ^{\mathrm{A}} \pm 0.57$ $12.00\ ^{\rm A}\pm 0.00$ E. coli ATCC 25922 R R Coagulase-negative $12.33 \ ^{B} \pm 0.57$ $15.00^{\text{A}} \pm 0.00$ R 9.61 ± 0.20 Staphylococcus $12.33\ ^{\rm A}\pm 0.57$ $12.00\ ^{\rm A} \pm 1.00$ K. pneumoniae sensible R R C. albicans ATCC 10231 $14.00\ ^{\rm A}\pm 0.00$ $14.67\ ^{\rm A} \pm 1.15$ 21.20 ± 4.20

Table 4. DIZ values of the two extracts of *C. humilis*.

ATCC: American Type Culture Collection; CIP: Institut Pasteur Collection; R: resistant; —: this antibiotic is ineffective against this strain; the difference in the two extracts is significant statistically (p < 0.05), as seen in ^{A,B}.

The extracts minimum inhibitory concentration (MIC) values were assessed against all of the microorganisms that were tested. Based on the finding in Table 5, the MIC recorded in this study was between 6.25 and 20 mg/mL. the MIC value needed to stop coagulase-negative *Staphylococcus* from growing was 6.25 mg/mL in the leaf extract, and 12.5 mg/mL in the root extract. *E. coli* CIP 53126 and E. coli ATCC 25922 were arrested at minimum concentration of 12.5 mg/mL in the root extract and 20 mg/mL in the leaf extract. At a minimum dose of 20 mg/mL, the other strains *K. pneumoniae* and *C. albicans* were suppressed.

Previous studies have shown that the highest antimicrobial effect could be related to the phenolic chemicals, in particular the presence of structural hydroxyl-phenolic groups, which increases the antimicrobial impact [50]. Further studies have confirmed the antimicrobial activity of polyphenols and flavonoids [51]. Indeed, the actuation mechanism of polyphenols in the antimicrobial activity is carried out by modifying the structure of the cell membrane, regulating its permeability, regulating cellular interactions by creating hydrogen bonds, and also decreasing the lipid content and finally preventing microbial development [52–56]. *C. humilis* is rich in polyphenols, especially chlorogenic acid, caffeic acid, isochlorogenic acid; in this sense, the antimicrobial effect of chlorogenic acid and caffeic acid has been well reported in the literature [6,49]

	Concentration (mg/mL)			
Strains of Bacteria	Leaf Extract	Root Extract		
E. coli CIP 53126	20	12.5		
E. coli ATCC 25922	20	12.5		
Coagulase-negative Staphylococcus	6.25	12.5		
K. pneumoniae sensible	20	20		
C. albicans ATCC 10231	20	20		

Table 5. C. humilis extracts and their minimum inhibitory concentration (MIC) values.

Luteolin, luteolin 7-rutinoside, and chlorogenic acid are components of *C. humilis*. Studies have shown compounds isolated from *C. cardunculus*. Among them, luteolin, luteolin 7-rutinoside, and chlorogenic acid exhibited more potent antimicrobial activities, and with an MIC varying between 0.03 and 0.10 mg/mL, the most active compound has been luteolin [46]. In another study, components extracted from the leaves of *C. scolymus*, among them luteolin 7-rutinoside and chlorogenic acid, were found to have a potential antibacterial action, with a MICs varying from 0.05–0.20 and 0.03–0.10 mg/mL [49]. Previous research has said that it has antibacterial properties of luteolin, apigenin, and other flavones [57,58]

2.5. Correlation between the Quality Parameters of C. humilis Extracts That Were Studied

According to the results displayed in Table 6, many of the factors examined in this study demonstrated a positive link between several of them. The IC₅₀ shows a substantial positive correlation with *E. coli* CIP 53126 and *Staphylococcus* strains (r = 0.960 and r = 0.890, respectively), whereas the TFC has a link correlation with *E. coli* ATCC 25922 strain (r = 0.700), and the TPC with TAC (r = 0.626). With *E. coli* ATCC 25922 strains, the TAC exhibits a positive association (r = 0.750). A high correlation exists between *E. coli* CIP 53126 and *Staphylococcus* (r = 0.891). The TAC, on the other hand, has a strong negative association with *Staphylococcus* (r = -0.925), and the TPC in turn has a strong negative correlation with IC₅₀ and *Staphylococcus* (r = -0.836 and r = -0.710, respectively). Additionally, a negative correlation was found for the TCF with *C. albicans* ATCC 10231 (r = -0.765) and the *E. coli* ATCC 25922 with *Staphylococcus* and *E. coli* CIP 53126 strains (r = -0.791 and r = -0.777, respectively).

Table	e 6. 1	Pearson	coefficients	of	corre	lation	between	С. і	humilis	s parameters
-------	--------	---------	--------------	----	-------	--------	---------	------	---------	--------------

Studied Parameters	ТРС	TFC	IC ₅₀	TAC	E. coli ATCC 25922	E. coli CIP53126	K. pneumoniae	Staph
TFC	0.407							
IC ₅₀	-0.836	-0.531						
TAC	0.626	0.318	-0.688					
E. coli ATCC 25922	0.435	0.700	-0.725	0.750				
E. coli CIP 53126	-0.655	-0.466	0.960	-0.663	-0.777			
K. pneumoniae	0.567	-0.211	-0.424	-0.024	-0.316	-0.307		
Staph	0.710	-0.384	0.890	-0.925	-0.791	0.891	-0.200	
C. albicans	0.280	-0.765	0.464	0.166	-0.316	0.430	-0.200	0.100

Overall, the considerable the link between polyphenols and flavonoids and free radical scavenging indicates that *C. humilis* possesses potent antioxidant properties. Our results confirm previous studies that polyphenols and flavonoids have a strong free radical scavenging capacity and exert antioxidant activity [39,59–61].

3. Materials and Methods

3.1. Reagents and Apparatus

The following compounds, Na₂CO₃, AlCl₃, Folin-Ciocalteu (RE00180250), potassium acetate, quercetin, sulfuric acid 0.6 M; sodium phosphate 28 mM, 4 mM ammonium molybdate, methanol, ethanol, 2,2-diphenyl-1-picrylhydrazyl, and NaCl were provided to Sigma-Aldrich, St. Louis, MO, États-Unis. Spectrophotometry (SOMESTIM, Rabat, Morocco) to determine the absorption capacity.

3.2. The Plant Used and Extraction Process

The study area is the region of Moulay Yacoub, the region is located 25 km northwest of the city fez and its area is estimated at 1700 km². the climate is characterized by high precipitation during the winter season, the annual average is estimated at 600 mm/year. For temperature, it is estimated between 6 to 36 °C, with a very hot summer. The region consists of a sedimentary series dominated by marly deposits [1,62]. The annual herb C. humilis pertains to the Asteraceae family (Figure 2), easily recognizable by its finely dissected leaves and its 4-sided cypselas with wing-like veins ending in a rigid yellowish spine. Unbranched taproot. Stem: has two rings of vascular bundles. Receptacle: slightly concave to slightly convex. Period of flowering: mainly from May to July. The plant is native to: Algeria, Canary Islands, Morocco, Portugal, and Spain [21,23]. The leaves and roots of C. humilis, which were employed in this study, were gathered in the Moulay Yacoub area, Morocco, from March to April 2020. The plant was identified (specimen number is MY002) by the botanist Lahcen Zidane, professor at the Faculty of Science of Kenitra, University Ibn Tofail, Kenitra, Morocco. Water was used to clean the plant's parts, dried at 35 °C, and powdered using a mortar. An amount of 50 g of dry powder plant weight was used for extraction, and the dry extract was obtained by maceration using a mixture of 70% ethanol and 30% distilled water, which means 10% of the plant powder (mass/volume). The macerate was filtered, then a rotary evaporator at 40 °C under partial vacuum was used to evaporate the solvent, finally, the acquired dry extract was utilized.



Figure 2. The plant studied, Cynara humilis.

The extraction yield of C. humilis roots and leaves was determined using the following report:

$$Y(\%) = (ME/MP) \times 100$$

Y: Yield of extraction (%); MP: Mass of plant powder used (g); ME: Mass of the crude extract obtained (g).

3.3. Determination of the Phenolic Content (TPC)

The Folin-Ciocalteu technique was used to estimate the phenolic compounds in the plant with modest alterations [30]. Plant extracts were mixed with 4 mL of 2 percent Na₂CO₃ and 200 μ L of Folin-Ciocalteu reagent in this experiment, the absorbance has

been determined spectrophotometrically at 670 nm after the solution had been left for thirty minutes. The equation is used to calculate the phenolic composition of the plant extracts was established from the calibration range of gallic acid, y = 5.4067x + 0.2213 and $R^2 = 0.9304$.

3.4. Determination of the Flavonoid Content (TFC)

The following procedure has been used to investigate the flavonoids composition [53]. An amount of 0.2 mL of AlCl₃ 10% and 2 mL of the plant extracts has been blended with 7.6 mL of ethanol and 0.2 mL of 1 M potassium acetate. After leaving the solution for forty min, at 430 nm, the absorbance has been measured spectrophotometrically. The equation used to calculate the content of flavonoids was established from quercetin's calibration curve, y = 0.0062x + 0.4289 and $R^2 = 0.857$.

3.5. Antioxidant Assay

3.5.1. Total Antioxidant Capacity (TAC)

To study the antioxidant assay TAC the following method was employed after a modest adjustment [63]. In this assay, 50 μ L of the plant extract was combined with 2 mL of a solution prepared from sulfuric acid 0.6 M, sodium phosphate 28 mM, and 4 mM ammonium molybdate. Afterwards, the reaction mixture has been placed in a water bath at 95 °C for thirty minutes. A spectrophotometer set at 695 nm has been used to measure the absorbance of the reaction medium, with a 50 μ L tube of methanol without extract serving as a negative control. A calibration curve utilizing ascorbic acid has been used to asses the total antioxidant capacity of the extract, the finding TAC values has been written by the unit (mg EAA/g of extract), which means milligrams of ascorbic acid equivalent by gram of extract.

3.5.2. Scavenging of the Free Radical DPPH

The following method [64] was employed to evaluate the DPPH antioxidant activity. In this assay, 0.75 mL of 2,2-diphenyl-1-picrylhydrazyl (0.004) has been mixed with 0.2 mL of each dilution series of the extracts studied. After leaving the solution for thirty minutes in the dark, the spectrophotometer has been determined at 517 nm.

The percentage of inhibition (PI) has been determined utilizing the report bellow:

PI (%) =
$$((A_0 - A)/A_0) \times 100$$

 A_0 : the absorbance value in the solution (2,2-diphenyl-1-picrylhydrazyl) which does not contain the extract.

A: the absorbance value in the solution (2,2-diphenyl-1-picrylhydrazyl) containing the extract.

3.6. Antimicrobial Activity of C. humilis

To study the antimicrobial test five bacterial strains were employed including two strains of *Escherichia coli*: *E. coli* CIP 53126, *E. coli* ATCC 25922, coagulase-negative *Staphylococcus*, *K. pneumoniae* susceptible, and a single fungal strain of *C. albicans* ATCC 10231. The microbial strains utilized in this assay were taken from the FMP-Fez microbiology laboratory, Morroco. Colonies were collected employing 24 h cultures to create bacterial suspensions, and the cultures were kept at 4 °C to preserve the cultures prepared in agar MH. The colonies that had been prepared were suspended and shook for 15 s in sterile solution (0.9% NaCL). The density was set to 0.5 turbidity (equivalent to $1-5 \times 10^8$ CFU/mL) [65].

3.6.1. Agar Disc Diffusion

The following methodology was used to examine the agar disk diffusion assay [64]. Bacterial suspensions were used to prepare MH agar plates (10⁸ CFU/mL), then inoculated by swabbing. After, Whatman paper discs (6 mm) were laid on the surface of the pre-

inoculated agar. Then, a volume of 20 μ L of 50 mg/mL of *C. humilis* was used to dip each Whatman paper disc. A disc of Whatman tempered in dimethyl sulfoxide was used as a negative control in the medium of each Petri dish, the plates were then incubated at 37 °C for 24 h, and the widths of the inhibitory zone (DIZ) were calculated. The bacteria have been categorized to use the latter (DIZ) [20].

- Non-sensitive: when the value of (DIZ) did not exceed 8 mm.
- Sensitive: when the value of (DIZ) was found in the interval 9 to 14 mm.
- Very sensitive: when the value of (DIZ) was found in the interval 15 to 19 mm.
- Extremely sensitive: when the value of (DIZ) exceeded 20 mm.

3.6.2. Minimum Inhibitory Concentration (MIC)

To investigate the test for the minimal inhibitory concentration, 96-well plates were employed in accordance with NCCLS guidelines [66], with minor adjustments. Using dimethyl sulfoxide, extracts from the leaves and roots of *C. humilis* were synthesized in sterile hemolysis tubes. The concentration of the extracts in the wells was prepared from successive 1:1 dilution in Mueller Hinton broth mixture. Arriving at values in the range of 0.039 to 20 mg/mL following that, 50 μ L of microbial suspensions were added to 5 μ L of Mueller Hinton broth, with various concentrations of the were applied to estimate MIC value. After that, the plates were kept at 37 °C for 18 h, after which 40 μ L of 0.5% triphenyl tetrazolium chloride was added to each well. The MIC value has been calculated utilizing the smallest concentration that won't result a red color [65].

3.7. Analytical Statistics

The means and standard deviations were used to present the results of this study's findings. The one-way ANOVA and student's test used to evaluate the statistical difference. Tukey's test is often used as a post hoc test to analyze the significant difference in TAC and DPPH assay, Minitab 19.1 is the software used in this work, p < 005 was utilized to designate the significant difference in all analyzed tests.

4. Conclusions

This study determined the chemical constitution and antioxidant and antimicrobial biological properties of the extracts of the roots and leaves of the plant C. humilis collected in spring 2020 in one locality in Morocco. The results indicate that leaves are richer in polyphenols (38.84 mg GAE/g extract) and flavonoids (27.07 mg QE/g extract) than the roots 29.56 mg GAE/g extract and 24.39 mg QE/g extract, respectively). The TAC antioxidant assay showed that the leaf extract (0.77 mg AAE/g extract) has higher antioxidant power than roots (0.60 mg AAE/g extract). In the DPPH antioxidant assay, the IC₅₀ values in the leaf and root extract were 0.23 and 0.93 μ g/mL, respectively. These extracts demonstrated antimicrobial effect on all strains studied, approved by a zone of inhibition with a diameter varying from 12 to 15 mm, and an MIC value of 6.25 mg/mL against coagulase-negative Staphylococcus in the root extract. These findings can be used in industry of pharmaceuticals to take care of microbial disorders and create medications to combat resistant bacteria, as well as in the food industry to preserve food using natural preservations rather than synthetic preservatives that have numerous adverse effects. Given that this is the first study on this plant, and only from one locality in Morocco, it is also difficult to judge the overall polyphenol content; therefore, additional research and repeated analyses in at least spring, summer, and autumn, during at least two growing seasons, and at several sites are needed to confirm these findings and identify the substances responsible for these biological activities.

Author Contributions: Conceptualization, M.E.K. and M.K.; methodology, M.E.K. and A.H.; software, M.K. and A.A.; investigation, M.E.K., M.K. and A.H.; data curation, M.K. and O.A.K.; writing—original draft preparation, M.E.K., H.F. and A.A.; writing—review and editing, M.E.K., M.K., O.A.K., S.E. and A.S.; supervision, D.H. All authors have read and agreed to the published version of the manuscript.

Funding: The paper is funded by Princess Nourah bint Abdulrahman University Researchers Supporting Project number (PNURSP2022R141), Princess Nourah bint Abdulrahman University, Riyadh, Saudi Arabia.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: The authors extend their appreciation to Princess Nourah bint Abdulrahman University Researchers Supporting Project number (PNURSP2022R141), Princess Nourah bint Abdulrahman University, Riyadh, Saudi Arabia.

Conflicts of Interest: The authors declare no conflict of interest.

References

- 1. El Khomsi, M.; Dandani, Y.; Chaachouay, N.; Hmouni, D. Ethnobotanical Study of Plants Used for Medicinal, Cosmetic, and Food Purposes in the Region of Moulay Yacoub, Northeast of Morocco. *J. Pharm. Pharmacogn. Res.* **2022**, *10*, 13–29.
- 2. Amin, H.I.M.; Hussain, F.H.S.; Gilardoni, G.; Thu, Z.M.; Clericuzio, M.; Vidari, G. Phytochemistry of *Verbascum* Species Growing in Iraqi Kurdistan and Bioactive Iridoids from the Flowers of *Verbascum calvum*. *Plants* **2020**, *9*, 1066. [CrossRef]
- 3. Ríos, J.L.; Recio, M.C. Medicinal Plants and Antimicrobial Activity. J. Ethnopharmacol. 2005, 100, 80–84. [CrossRef] [PubMed]
- Kraft, K. Complementary/Alternative Medicine in the Context of Prevention of Disease and Maintenance of Health. *Prev. Med.* 2009, 49, 88–92. [CrossRef] [PubMed]
- Bungau, S.G.; Popa, V.-C. Between religion and science: Some aspects: Concerning illness and healing in antiquity. *Transylv. Rev.* 2015, 26, 3–19.
- El Khomsi, M.; Imtara, H.; Kara, M.; Hmamou, A.; Assouguem, A.; Bourkhiss, B.; Tarayrah, M.; AlZain, M.N.; Alzamel, N.M.; Noman, O.; et al. Antimicrobial and Antioxidant Properties of Total Polyphenols of *Anchusa italica* Retz. *Molecules* 2022, 27, 416. [CrossRef] [PubMed]
- Hussain, F.H.S.; Osw, P.S. Isolation of Kaempferol 3-O-Rutinoside From Kurdish Plant Anchusa italica Retz. And Bioactivity of Some Extracts. Eur. J. Mol. Clin. Med. 2020, 7, 2465–2480.
- Mssillou, I.; Agour, A.; El Ghouizi, A.; Hamamouch, N.; Lyoussi, B.; Derwich, E. Chemical Composition, Antioxidant Activity, and Antifungal Effects of Essential Oil from *Laurus nobilis* L. Flowers Growing in Morocco. *J. Food Qual.* 2020, 2020, e8819311. [CrossRef]
- Pandey, B.P.; Thapa, R.; Upreti, A. Chemical Composition, Antioxidant and Antibacterial Activities of Essential Oil and Methanol Extract of *Artemisia vulgaris* and *Gaultheria fragrantissima* Collected from Nepal. *Asian Pac. J. Trop. Med.* 2017, 10, 952–959.
 [CrossRef]
- Sepahvand, R.; Delfan, B.; Ghanbarzadeh, S.; Rashidipour, M.; Veiskarami, G.H.; Ghasemian-Yadegari, J. Chemical Composition, Antioxidant Activity and Antibacterial Effect of Essential Oil of the Aerial Parts of *Salvia sclareoides*. *Asian Pac. J. Trop. Med.* 2014, 7, S491–S496. [CrossRef]
- 11. Nagai, T.; Myoda, T.; Nagashima, T. Antioxidative Activities of Water Extract and Ethanol Extract from Field Horsetail (Tsukushi) Equisetum arvense L. Food Chem. 2005, 91, 389–394. [CrossRef]
- 12. Bouzidi, L.E.; Larhsini, M.; Markouk, M.; Abbad, A.; Hassani, L.; Bekkouche, K. Antioxidant and Antimicrobial Activities of *Withania frutescens. Nat. Prod. Commun.* **2011**, *6*, 1447–1450. [CrossRef] [PubMed]
- 13. Gülçın, İ.; Oktay, M.; Kıreçcı, E.; Küfrevıoğlu, Ö.İ. Screening of Antioxidant and Antimicrobial Activities of Anise (*Pimpinella anisum* L.) Seed Extracts. *Food Chem.* **2003**, *83*, 371–382. [CrossRef]
- Caturla, N.; Vera-Samper, E.; Villalaín, J.; Mateo, C.R.; Micol, V. The Relationship between the Antioxidant and the Antibacterial Properties of Galloylated Catechins and the Structure of Phospholipid Model Membranes. *Free Radic. Biol. Med.* 2003, 34, 648–662. [CrossRef]
- 15. Lahkimi, A.; Louaste, B.; Nechad, I.; Chaouch, M.; Eloutassi, N. Antibacterial, Antifungal and Antioxidant Activity of *Lavandula angustifolia* of the Middle Atlas Central (Morocco). *Moroc. J. Chem.* **2020**, *8*, 8–918. [CrossRef]
- Jaloob, A.A.A. Antibacterial Activity of an Aqueous Extracts of *Alkanna tinctoria* Roots against Drug Resistant Aerobic Pathogenic Bacteria Isolated from Patients with Burns Infections. *Russ. Open Med. J.* 2018, 7, 104.
- 17. Michel, T. Nouvelles Méthodologies D'extraction, de Fractionnement et D'identification: Application aux Molécules Bioactives de L'argousier (*Hippophae rhamnoides*). Ph.D. Thesis, Université d'Orléans, Orléans, France, 2011.

- 18. Farzaei, M.H.; Rahimi, R.; Abbasabadi, Z.; Abdollahi, M. An Evidence-Based Review on Medicinal Plants Used for the Treatment of Peptic Ulcer in Traditional Iranian Medicine. *Int. J. Pharmacol.* **2013**, *9*, 108–124. [CrossRef]
- 19. Rahimi, R.; Abdollahi, M. Herbal Medicines for the Management of Irritable Bowel Syndrome: A Comprehensive Review. *World J. Gastroenterol.* **2012**, *18*, 589–600. [CrossRef]
- 20. El Moussaoui, A.; Jawhari, F.Z.; Almehdi, A.M.; Elmsellem, H.; Fikri Benbrahim, K.; Bousta, D.; Bari, A. Antibacterial, Antifungal and Antioxidant Activity of Total Polyphenols of *Withania frutescens* L. *Bioorganic Chem.* **2019**, *93*, 103337. [CrossRef]
- Curci, P.L.; Sonnante, G. The Complete Chloroplast Genome of *Cynara humilis*. *Mitochondrial DNA Part A* 2016, 27, 2345–2346. [CrossRef]
- 22. Rottenberg, A.; Zohary, D. The Wild Ancestry of the Cultivated Artichoke. Genet. Resour. Crop Evol. 1996, 43, 53–58. [CrossRef]
- Wiklund, A. The Genus *Cynara* L.(Asteraceae-Cardueae). *Bot. J. Linn. Soc.* 1992, 109, 75–123. [CrossRef]
 Salhi, N.; Bouyahya, A.; Fettach, S.; Zellou, A.; Cherrah, Y. Ethnopharmacological Study of Medicinal Plants Used in the Treatment
- Saini, N.; Bouyanya, A.; Fettach, S.; Zellou, A.; Cherran, Y. Ethnopharmacological Study of Medicinal Plants Used in the Treatment of Skin Burns in Occidental Morocco (Area of Rabat). S. Afr. J. Bot. 2019, 121, 128–142. [CrossRef]
 National Action of Skin Burns in Occidental Morocco (Area of Rabat). S. Afr. J. Bot. 2019, 121, 128–142. [CrossRef]
- Nassiri, L.; Zarkani, S.; Daoudi, A.; Bammou, M.; Bouiamrine, E.H.; Ibijbijen, J. Contribution à l'élaboration d'un Catalogue Ethnobotanique de La Commune Rurale d'Aguelmous (Province de Khénifra, Maroc)[Contribution to the Establishment of Ethno Botanical Catalog of Aguelmous (Khenifra, Morocco)]. *Int. J. Innov. Appl. Stud.* 2016, 17, 373–387.
- 26. Salhi, N.; Bouyahya, A.; Bounihi, A.; Masrar, A.; Bouabdellah, M.; Chabraoui, L.; Zengin, G.; Taghzouti, K.; Rouas, L.; Cherrah, Y. Investigation of Wound Healing Activity *Cynara humilis* of Root Extracts. *J. Cosmet. Dermatol.* **2022**, *21*, 1596–1609. [CrossRef]
- Chinou, I.; Harvala, C. Polyphenolic Constituents from the Leaves of Two Cynara Species Growing in Greece. *Planta Med.* 1997, 63, 469–470. [CrossRef]
- Reis, L.V.; Regina Tavares, M.; Palma, F.M.S.B.; Marcelo-Curto, M.J. Sesquiterpene Lactones from *Cynara humilis*. *Phytochemistry* 1992, 31, 1285–1287. [CrossRef]
- 29. Silva, S.V.; Malcata, F.X. Action of Cardosin A from *Cynara humilis* on Ovine and Caprine Caseinates. *J. Dairy Res.* 2000, 67, 449–454. [CrossRef]
- 30. Kara, M.; Assouguem, A.; Abdou, R.Z.; Bahhou, J. Phytochemical Content and Antioxidant Activity of Vinegar Prepared from Four Apple Varieties by Different Methods. *Trop. J. Nat. Prod. Res.* **2021**, *5*, 1578–1585.
- 31. Minnocci, A.; Iacopini, P.; Martinelli, F.; Sebastiani, L. Micromorphological, Biochemical, and Genetic Characterization of Two Ancient, Late-Bearing Apple Varieties. *Eur. J. Hort. Sci.* 2010, 75, 1–7.
- 32. McGhie, T.K.; Hunt, M.; Barnett, L.E. Cultivar and Growing Region Determine the Antioxidant Polyphenolic Concentration and Composition of Apples Grown in New Zealand. *J. Agric. Food Chem.* **2005**, *53*, 3065–3070. [CrossRef] [PubMed]
- Glevitzky, I.; Dumitrel, G.A.; Glevitzky, M.; Pasca, B.; Otrisal, P.; Bungau, S.; Cioca, G.; Pantis, C.; Popa, M. Statistical Analysis of the Relationship between Antioxidant Activity and the Structure of Flavonoid Compounds. *Rev. Chim* 2019, 70, 3103–3107. [CrossRef]
- 34. Falleh, H.; Ksouri, R.; Chaieb, K.; Karray-Bouraoui, N.; Trabelsi, N.; Boulaaba, M.; Abdelly, C. Phenolic Composition of *Cynara cardunculus* L. Organs, and Their Biological Activities. *Comptes Rendus Biol.* **2008**, *331*, 372–379. [CrossRef] [PubMed]
- 35. Vamanu, E.; Vamanu, A.; Nita, S.; Colceriu, S. Antioxidant and Antimicrobial Activities of Ethanol Extracts of *Cynara scolymus* (*Cynarae* Folium, Asteraceae Family). *Trop. J. Pharm. Res.* **2011**, *10*, 777–783. [CrossRef]
- Miliauskas, G.; Venskutonis, P.R.; Van Beek, T.A. Screening of Radical Scavenging Activity of Some Medicinal and Aromatic Plant Extracts. *Food Chem.* 2004, 85, 231–237. [CrossRef]
- 37. Lee, C.E.; Petersen, C.H. Effects of Developmental Acclimation on Adult Salinity Tolerance in the Freshwater-Invading Copepod Eurytemora Affinis. *Physiol. Biochem. Zool.* **2003**, *76*, 296–301. [CrossRef]
- Deng, G.-F.; Lin, X.; Xu, X.-R.; Gao, L.-L.; Xie, J.-F.; Li, H.-B. Antioxidant Capacities and Total Phenolic Contents of 56 Vegetables. J. Funct. Foods 2013, 5, 260–266. [CrossRef]
- Hmamou, A.; Eloutassi, N.; Alshawwa, S.Z.; Al kamaly, O.; Kara, M.; Bendaoud, A.; El-Assri, E.-M.; Tlemcani, S.; El Khomsi, M.; Lahkimi, A. Total Phenolic Content and Antioxidant and Antimicrobial Activities of *Papaver rhoeas* L. Organ Extracts Growing in Taounate Region, Morocco. *Molecules* 2022, 27, 854. [CrossRef]
- 40. Sakanaka, S.; Tachibana, Y.; Okada, Y. Preparation and Antioxidant Properties of Extracts of *Japanese persimmon* Leaf Tea (Kakinoha-Cha). *Food Chem.* **2005**, *89*, 569–575. [CrossRef]
- 41. Benavente-García, O.; Castillo, J.; Marin, F.R.; Ortuño, A.; Del Río, J.A. Uses and Properties of Citrus Flavonoids. J. Agric. Food Chem. 1997, 45, 4505–4515. [CrossRef]
- Alonso, Á.M.; Castro, R.; Rodríguez, M.C.; Guillén, D.A.; Barroso, C.G. Study of the Antioxidant Power of Brandies and Vinegars Derived from Sherry Wines and Correlation with Their Content in Polyphenols. *Food Res. Int.* 2004, 37, 715–721. [CrossRef]
- 43. Trueba, G.P.; Sanchez, G.M. Los Flavonoides Como Antioxidantes Naturales. Acta Farm. Bonaer. 2001, 20, 297–306.
- Belščak-Cvitanović, A.; Durgo, K.; Huđek, A.; Bačun-Družina, V.; Komes, D. 1—Overview of Polyphenols and Their Properties. In *Polyphenols: Properties, Recovery, and Applications*; Galanakis, C.M., Ed.; Woodhead Publishing: Sawston, UK, 2018; pp. 3–44, ISBN 978-0-12-813572-3.
- 45. Ravichandran, R.; Rajendran, M.; Devapiriam, D. Antioxidant Study of Quercetin and Their Metal Complex and Determination of Stability Constant by Spectrophotometry Method. *Food Chem.* **2014**, *146*, 472–478. [CrossRef]
- Kukić, J.; Popović, V.; Petrović, S.; Mucaji, P.; Ćirić, A.; Stojković, D.; Soković, M. Antioxidant and Antimicrobial Activity of Cynara Cardunculus Extracts. *Food Chem.* 2008, 107, 861–868. [CrossRef]

- 47. Wang, M.; Li, J.; Rangarajan, M.; Shao, Y.; LaVoie, E.J.; Huang, T.-C.; Ho, C.-T. Antioxidative Phenolic Compounds from Sage (*Salvia officinalis*). J. Agric. Food Chem. **1998**, 46, 4869–4873. [CrossRef]
- 48. Takeoka, G.R.; Dao, L.T. Antioxidant Constituents of Almond [Prunus Dulcis (Mill.) D.A. Webb] Hulls. J. Agric. Food Chem. 2003, 51, 496–501. [CrossRef] [PubMed]
- 49. Zhu, X.; Zhang, H.; Lo, R. Phenolic Compounds from the Leaf Extract of Artichoke (*Cynara scolymus* L.) and Their Antimicrobial Activities. J. Agric. Food Chem. 2004, 52, 7272–7278. [CrossRef]
- Rota, M.C.; Herrera, A.; Martínez, R.M.; Sotomayor, J.A.; Jordán, M.J. Antimicrobial Activity and Chemical Composition of Thymus vulgaris, Thymus zygis and Thymus hyemalis Essential Oils. Food Control 2008, 19, 681–687. [CrossRef]
- 51. Rojas, A.; Hernandez, L.; Pereda-Miranda, R.; Mata, R. Screening for Antimicrobial Activity of Crude Drug Extracts and Pure Natural Products from Mexican Medicinal Plants. *J. Ethnopharmacol.* **1992**, *35*, 275–283. [CrossRef]
- 52. Bakour, M.; da Graça Campos, M.; Imtara, H.; Lyoussi, B. Antioxidant Content and Identification of Phenolic/Flavonoid Compounds in the Pollen of Fourteen Plants Using HPLC-DAD. J. Apic. Res. 2020, 59, 35–41. [CrossRef]
- 53. Yang, J.-F.; Yang, C.-H.; Wu, C.-C.; Chuang, L.-Y. Antioxidant and Antimicrobial Activities of the Extracts from *Sophora flavescens*. *J. Pharmacogn. Phytochem.* **2015**, *3*, 26–31.
- 54. Baghiani, A.; Boussoualim, N.; Trabsa, H.; Aouachria, S.; Arrar, L. In Vivo Free Radical Scavenging, Antihemolytic Activity and Antibacterial Effects of Anchusa Azurea Extracts. *Int. J. Med. Med. Sci.* **2013**, *46*, 1113–1118.
- 55. Ikigai, H.; Nakae, T.; Hara, Y.; Shimamura, T. Bactericidal Catechins Damage the Lipid Bilayer. *Biochim. Biophys. Acta BBA Biomembr.* **1993**, 1147, 132–136. [CrossRef]
- Helal, G.A.; Sarhan, M.M.; Abu Shahla, A.N.K.; Abou El-Khair, E.K. Effects of Cymbopogon Citratus L. Essential Oil on the Growth, Lipid Content and Morphogenesis of *Aspergillus niger* ML2-Strain. J. Basic Microbiol. 2006, 46, 456–469. [CrossRef] [PubMed]
- 57. Aljančić, I.; Vajs, V.; Menković, N.; Karadžić, I.; Juranić, N.; Milosavljević, S.; Macura, S. Flavones and Sesquiterpene Lactones from *Achillea atrata* Subsp. Multifida: Antimicrobial Activity. *J. Nat. Prod.* **1999**, *62*, 909–911. [CrossRef]
- 58. Tshikalange, T.E.; Meyer, J.J.M.; Hussein, A.A. Antimicrobial Activity, Toxicity and the Isolation of a Bioactive Compound from Plants Used to Treat Sexually Transmitted Diseases. *J. Ethnopharmacol.* **2005**, *96*, 515–519. [CrossRef]
- Maisuthisakul, P.; Suttajit, M.; Pongsawatmanit, R. Assessment of Phenolic Content and Free Radical-Scavenging Capacity of Some Thai Indigenous Plants. *Food Chem.* 2007, 100, 1409–1418. [CrossRef]
- 60. Kiselova, Y.; Ivanova, D.; Chervenkov, T.; Gerova, D.; Galunska, B.; Yankova, T. Correlation between the In Vitro Antioxidant Activity and Polyphenol Content of Aqueous Extracts from Bulgarian Herbs. *Phytother. Res.* **2006**, *20*, 961–965. [CrossRef]
- Ahmed, S.; Jubair, A.; Hossain, M.A.; Hossain, M.M.; Azam, M.S.; Biswas, M. Free Radical-Scavenging Capacity and HPLC-DAD Screening of Phenolic Compounds from Pulp and Seed of *Syzygium claviflorum* Fruit. J. Agric. Food Res. 2021, 6, 100203. [CrossRef]
- Lakhdar, A.; Ntarmouchant, A.; Ribeiro, M.L.; Beqqali, M.; El Ouadeihe, K.; Benaabidate, L.; Dahire, M.; Driouch, Y.; Slimane, A.B. Nouvelle approche géologique et géodinamique du Complexe Hydrothermal de Moulay Yacoub, bordure septentrionale du Sillon Sud Rifain. *Comun. Geol.* 2006, 93, 185–204.
- Mašković, P.Z.; Manojlović, N.T.; Mandić, A.I.; Mišan, A.Č.; Milovanović, I.L.; Radojković, M.M.; Cvijović, M.S.; Solujić, S.R. Phytochemical Screening and Biological Activity of Extracts of Plant Species *Halacsya sendtneri* (Boiss.) Dörfl. *Hem. Ind.* 2012, 66, 43–51. [CrossRef]
- 64. Sadeq, O.; Mechchate, H.; Es-safi, I.; Bouhrim, M.; Jawhari, F.; Ouassou, H.; Kharchoufa, L.; AlZain, M.N.; Alzamel, N.M.; Al kamaly, O.M.; et al. Phytochemical Screening, Antioxidant and Antibacterial Activities of Pollen Extracts from *Micromeria fruticosa*, *Achillea fragrantissima*, and *Phoenix dactylifera*. *Plants* **2021**, *10*, 676. [CrossRef] [PubMed]
- Kara, M.; Assouguem, A.; Benmessaoud, S.; Imtara, H.; Mechchate, H.; Hano, C.; Zerhouni, A.R.; Bahhou, J. The Impact of Apple Variety and the Production Methods on the Antibacterial Activity of Vinegar Samples. *Molecules* 2021, 26, 5437. [CrossRef] [PubMed]
- Gulluce, M.; Sahin, F.; Sokmen, M.; Ozer, H.; Daferera, D.; Sokmen, A.; Polissiou, M.; Adiguzel, A.; Ozkan, H. Antimicrobial and Antioxidant Properties of the Essential Oils and Methanol Extract from *Mentha longifolia* L. Ssp. Longifolia. *Food Chem.* 2007, 103, 1449–1456. [CrossRef]