

Brief Report

Hydrocharis laevigata in Europe

Pablo Garcia-Murillo 

Department of Plant Biology and Ecology, Faculty of Pharmacy, University of Seville, 41012 Seville, Spain; pgarcia@us.es

Abstract: *Hydrocharis laevigata* (Humb. & Bonpl. ex Willd.) Byng & Christenh. [= *Limnobium laevigatum* (Humb. & Bonpl. ex Willd.) Heine], Hydrocharitaceae, is a floating-leaf aquatic plant that is native to inland South America. It is an invasive species in several parts of the world. Reports of its presence in Europe have been recently published: naturalised populations occur in three locations on the Iberian Peninsula. The literature also contains records of the species in Hungary and Poland. In addition, it has been observed in Sweden, Belgium, and the Netherlands. *H. laevigata* is highly adaptable and can profoundly transform habitat conditions in its invasive range, causing major issues for ecosystem conservation and human activities. Until recently, *H. laevigata* was not to be found in natural environments in Europe. Factors explaining its spread include its use as an ornamental plant, the eutrophication of inland waters, and the effects of global warming. With a focus on Europe, this short communication provides information on the species' distribution, taxonomy, biology, habitat, and negative impacts.

Keywords: aquatic plants; spongeplant; AIS; biological invasions



Citation: Garcia-Murillo, P.

Hydrocharis laevigata in Europe. *Plants* **2023**, *12*, 701. <https://doi.org/10.3390/plants12040701>

Academic Editors: Bruce Osborne and Panayiotis Dimitrakopoulos

Received: 23 January 2023

Revised: 1 February 2023

Accepted: 2 February 2023

Published: 4 February 2023



Copyright: © 2023 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The major decline in freshwater species biodiversity has become one of the most important alterations in ecosystems around the world [1,2]. Pollution, water flow restrictions, and climate change are transforming conditions in freshwater habitats, favouring the establishment of invasive species [3–5].

There is a common consensus, supported by the Red List of European Habitats [6], that the key threats to European aquatic and wetland habitats are alterations to hydrological systems, climate change, pollution, and invasive species [7]. In Europe, native freshwater biota are facing increasing competitive pressure due to the arrival and spread of floating-leaf aquatic plants (e.g., *Azolla filiculoides* and *Eichhornia crassipes*) in inland water bodies [8–15]. Of particular concern is the recent appearance of an aggressive invasive alien species (AIS), *Hydrocharis laevigata*, whose common names include South American spongeplant, smooth frogbit, or Amazon frogbit. Newly established in a range of European ponds, rivers, and reservoirs, the species serves as yet another warning about the deterioration of Europe's freshwater habitats. Its invasive dynamics are reminiscent of those of *Eichhornia crassipes*, including its striking colonisation abilities. This short communication describes *H. laevigata*'s history in Europe, including its emergence in distinct continental regions. It also provides key information about the species' taxonomy, biology, habitat, and negative impacts. To this end, it will facilitate prevention efforts aimed at invasive species, help control the spread of *H. laevigata*, and bolster the preservation of Europe's native biodiversity.

2. Results

2.1. Taxonomy and Nomenclature

Hydrocharis laevigata (Humb. & Bonpl. ex Willd.) Byng & Christenh. belongs to the family Hydrocharitaceae (superorder Lilianae). This taxon of monocots contains genera

whose members are among the world's worst aquatic invasive plants, such as *Eichhornia* Kunth and *Hydrilla* Rich.

Phylogenomic, morphological, and anatomical evidence strongly suggest a new vision of the genus *Hydrocharis* L. [16,17], one that includes the species of *Limnobium* Rich., usually considered to be a separate genus by World Flora Online (WFO) [18]. Assuming the species' reclassification, there are currently five known members of the genus *Hydrocharis* L. Three are spread throughout Afro-Asia, and the other two are found from Mexico to northern Argentina and in the Caribbean. Those previously classified in *Limonium* are native to the Americas and utilise wind pollination. The original members of *Hydrocharis* are autochthonous from the Old World and are insect pollinated [19]. Those differences aside, all five species have a close phylogenetic relationship and display similarities in their vegetative traits [19,20].

Hydrocharis laevigata (Humb. & Bonpl. ex Willd.) Byng & Christenh., Global Fl., 4: 53. 2018.
 ≡ *Salvinia laevigata* Humb. & Bonpl. ex Willd., Sp. Pl., ed. 4, 5: 537. 1810 (basionym).
 ≡ *Limnobium laevigatum* (Humb. & Bonpl. ex Willd.) Heine in Adansonia, 8: 315. 1968.
 ≡ *Hydromystria laevigata* (Humb. & Bonpl. ex Willd.) Hunz., Lorentzia, 4: 5. 1981.
 ≡ *Limnobium spongia* subsp. *laevigatum* (Humb. & Bonpl. ex Willd.) Lowden, Rhodora, 94: 129. 1992).

Type (lectotype designated by [21] (p. 220): Colombia; Bogotá. Humboldt & Bonpland (holotype: B-Willd. 20251).

= *Hydromystria stolonifera* G. Mey., Prim. Fl. Esseq. 153. 1818.

≡ *Limnobium stoloniferum* (G. Mey.) Griseb., Fl. Brit. W. I. 506. 1864.

This species has traditionally been included in the genus *Limnobium* Rich., under the name *Limnobium laevigatum* (Humb. & Bonpl. ex Willd.) Heine. Other authors have considered this taxon to be a subspecies of *L. spongia* (Bosc) Steud.

2.2. Species Description

According to [21–24], *H. laevigata* is an aquatic stoloniferous plant whose stems are floating on or suspended in the water. They develop floating or emergent leaf rosettes, which are sometimes rooted. The species' differentiated leaves are petiolate, laminate, cordate to reniform, or orbiculate, stipulate, and dimorphic. The aerial leaves have flat, leathery blades and long erect petioles measuring up to 40 cm; the floating leaves have thick spongy tissue on the lower surface and shorter petioles (minimum of 12 cm). Flowers are about 15 mm wide, unisexual, pedicelled, and enclosed in a spathe. Staminate flowers often occur in clusters of two or three; have three (rarely four) sepals and petals each; possess a variable number of stamens, whose filaments form a single column; and usually exhibit staminodes. Pistillate flowers are solitary; have rudimentary sepals and petals, although the latter may be lacking; and possess a unilocular ovary with six carpels, six styles, and divided linear stigmas that are longer than the perianth segments. Furthermore, female flowers may sometimes exhibit staminodes. The species' fruit consists of globose or ellipsoid fleshy berry-like capsules (~1.5 cm long and 0.4–3.5 mm wide) that contain as many as 100 ellipsoid seeds measuring around 10 mm long.

2.3. Habitat

Found in the tropics and subtropics [22,25], *H. laevigata* grows in a variety of freshwater habitats, such as shallow ponds; lakes; dams; reservoirs; pools; and the margins of slow shady rivers, canals, and ditches; environments with a high degree of eutrophication are particularly hospitable [26,27]. The species occurs at elevations from sea level all the way up to 2800 m. It is absent from fast-flowing watercourses [21,22,26,28], and cold temperatures limit its distribution [25].

2.4. Reproduction

Hydrocharis laevigata reproduces vegetatively via the fragmentation of the stolon segments connecting the leaf rosettes. The floating rosettes send out stolons bearing ramets on

their ends [29]. The rosettes are easily dislodged by waterfowl or large mammals in the species' native range [26], as well as by water currents or other types of water movement in general. Consequently, *H. laevigata* is able to rapidly and efficiently colonise suitable new habitats [28].

With regards to sexual reproduction, the species mainly utilises wind pollination [17,22,28]. However, water pollination has also been observed, as has insect pollination, mediated by tiny aphid nymphs (Aphididae) and the aphids' predators, ladybird beetles (Coccinellidae) [24].

After pollination, the pedicel of female flowers tips downwards, such that the fruit develops submerged in water; once it reaches maturity, it breaks open, releasing a mucilaginous mass of about 100 seeds [21].

2.5. Global Distribution

The native range of *H. laevigata* extends across the tropics of Central and South America. Naturalised populations can now be found on all continents, with the exception of Antarctica.

According to [22,24,30], this species is naturally found in water bodies in parts of the Americas with tropical, subtropical, and, sometimes, temperate climates. More specifically, these locations are the CARIBBEAN: Antigua and Barbuda, Cuba, the Dominican Republic, Guadeloupe, St. Lucia, Montserrat, Martinique, Trinidad and Tobago, and Puerto Rico; CENTRAL AMERICA: Mexico, Costa Rica, Guatemala, Nicaragua, Panama, and El Salvador; and SOUTH AMERICA: French Guiana, Guyana, Suriname, Venezuela, Brazil, Colombia, Ecuador, Peru, Argentina, Paraguay, and Uruguay. It should also be noted that there may exist some problematic records for South America. For example, although [31] stated that the species is native to the Southern Cone, this assertion is questionable in the case of Chile, as [26] indicates that the country's oldest herbarium specimen dates back to 1954. The same is likely true for Argentina: while the species' native range is thought to extend across most of the country, it certainly appears to be invasive in the southernmost regions of Argentina, where it has recently appeared for the first time [32].

As noted in [25,29,33–36], *H. laevigata* has become naturalised and spread across ASIA: Japan, Taiwan, and Indonesia; AFRICA: Zambia and Zimbabwe; AUSTRALIA: New South Wales and Queensland; NORTH AMERICA: southern Canada as well as the north-eastern and western US; and EUROPE.

2.6. Populations in Europe

Based on the scientific literature and the GBIF database, *H. laevigata* is known to occur in six European countries: Belgium, Hungary, The Netherlands, Poland, Spain, and Sweden.

The first record of its presence (as *Limnobiium laevigatum*) was in hot springs located in Komárom-Esztergom County, northern Hungary [37]. In September 2018, the species was observed in the area of Fényes springs, north of Tata. At this location, the plants were mainly in a vegetative state, although some did bear flowers. One month later, the plants had spread to the Ferenccmajori fishponds, traveling a distance of about 1.3 km. In October 2018, the species turned up at another site within the same county: Dunaalmás, a stream fed by warm water [37]. It was observed that the *H. laevigata* populations became established near hot springs (temperatures = 21–23 °C), which do not freeze in the winter. Recently, Dr. Riezinger (pers. comm.) has indicated that these populations in northern Hungary have persisted, likely because of mild winter conditions and the warm water temperatures. However, the plants occur in small groups.

On the Iberian Peninsula, the species seems to have rapidly expanded its range. Since 2018, it has been found at three locations that are quite distant from each other. The first published record for this species (as *Limnobiium laevigatum*) was associated with a reservoir near Madrid [38]. A few years later, it was observed in a stream next to a dam near Córdoba [39]. Finally, it was also reported to occur in the Guadaira, a tributary of the

Gualdalquivir, near Seville [40] (Figure 1). There are no indications that *H. laevigata* has disappeared from any of these locations in Spain.



Figure 1. (A,B) *Hydrocharis laevigata* habitat in the outskirts of Seville, Spain. (C) Male flowers, photos taken by P. Garcia Murillo in summer 2021; (D,E) habitat in the outskirts of Krakow, Poland, photos taken in November and December 2020, courtesy of Dr. Arthur Pliszko.

In 2020, *H. laevigata* was observed in southern Poland, in an artificial pond near Agatowa Street in the district of Bieżanów-Prokocim in eastern Kraków [15]. It was noted [15] that the species (as *L. laevigatum*) is likely a sporadic invader in Poland, given that other similar invasive species display constrained ranges due to the country's cold winter temperatures. Dr. Pliszko (pers. comm.) recently confirmed that this population has disappeared because of the species' vulnerability to low temperatures. It cannot survive winter conditions in Poland, outside of the habitat found in greenhouses or residences (Figure 1).

The GBIF lists nine geographical records for *H. laevigata* in Europe [41], which include Ganshoren, Belgium; the outskirts of Stockholm, Sweden; and Aalsmeer and Bleiswijk in the Netherlands. Only in the latter country are the records associated with herbarium specimens. In the other cases, there are only photographs, no herbarium samples (Appendices A and B). Two of these locations represent artificial habitats: a greenhouse in Aalsmeer in the Netherlands and the Bergianska Botanick Garden in Sweden. The remaining records for Belgium and the Netherlands were one-time events. Consequently,

they can be viewed as sporadic occurrences that have not resulted in *H. laevigata* becoming naturalised. The same is true for the record in Poland. Therefore, at present, the only naturalised *H. laevigata* populations in Europe are in Hungary and Spain (Figure 2).




Figure 2. Locations where *Hydrocharis laevigata* has been observed in Europe. Blue dots: sporadic presence. Red dots: naturalised populations. When the name of the city has a different spelling of the English alphabet, it appears below in its original spelling. Map created using The Geospatial Conservation Assessment Tool.

2.7. Invasiveness

As noted in the introduction, *H. laevigata* displays invasive dynamics reminiscent of those of *Eichhornia crassipes*, including its marked capacity for colonisation [25,32,42]. Indeed, like other invasive aquatic plant species, it is fast growing and thrives in still waters containing high concentrations of nutrients [43]. Under these conditions, it can outcompete native aquatic plant species and profoundly transforms the habitats it invades. While the impacts of this species have yet to be fully described, past observations [22,25,28,44] indicate that the floating rosettes block light from reaching the water below, indirectly fostering anoxic conditions. *H. laevigata* can thus reduce the abundance and diversity of aquatic plants and animals (e.g., fish, phytoplankton, submersed plants) by rendering

water bodies unsuitable for other organisms. The species can also generate mats that cause damage to human infrastructures (e.g., by clogging irrigation pipes or sewers, by damaging pumps or generators in hydroelectric power stations). Consequently, *H. laevigata* is considered to be one of the most harmful weeds in the United States and Australia, given its high degree of invasiveness combined with its negative impacts on water quality and aquatic biodiversity [25,29,42,44].

While *H. spongia* (= *Limnobium spongia*) is rarely cultivated, *H. laevigatum* is commonly found in aquaria and garden ponds, largely because of its ecological breadth and pronounced reproductive capacity [32,35–37,45].

2.8. Control

Typical methods for controlling *H. laevigata* are mechanical harvesting and herbicide treatments [45,46]. In the Sacramento-San Joaquin Delta, diquat dibromide has been used, either on its own or in tandem with low levels of glyphosate [46]; this work yielded unclear results. Another study [47] evaluated the performance of the herbicides imazamox, penoxsulam and topramezone. Imazamox (with 1% methylated seed oil surfactant) was found to control all the growth stages of *H. laevigata*; penoxsulam was particularly effective in targeting seedling and rosette growth. Research is currently lacking on potential biological control methods. However, [36] points out that certain North American herbivores display a dietary specialisation for American frogbit (*H. spongia*). They might therefore be candidate biological control agents.

3. Discussion

In summary, *H. laevigata* is an invasive plant species that poses a significant threat to ecosystems and certain human activities in its non-native range. Its likelihood of establishment is considerably elevated in nutrient-rich, slow-moving freshwater systems subject to mild winter temperatures [25–28], such as those found in southern Europe or the northern Mediterranean. Its invasion dynamics are similar to those of *Eichhornia crassipes* [25,32,42].

As indicated earlier, the ornamental plant trade is the main source of *H. laevigata* (and *Eichhornia crassipes*) introductions. This fact explains how the species has spread so far from its native range to parts of the world as diverse as Australia, Zimbabwe, Japan, and Spain.

The range expansion of *H. laevigata* within Europe underscores the deterioration of the continent's inland aquatic ecosystems. Such is, above all, the result of eutrophication, but other causal factors include changes to hydrological regimes (i.e., reduced levels of water flow), decreasing degrees of water transparency, and global warming, which are simultaneously transforming natural aquatic environments [5,48–53]. These ecological shifts are harmful to native aquatic plant populations while promoting the establishment of invasive species such as *H. laevigata*, leading to profound transformations in European ecosystems [12,14,54–57].

It is curious to note that, while the invasive *H. laevigata* has begun to expand its range in Europe, the native *H. morsus-ranae*, a close relative, is experiencing decline. While the latter has always been patchily distributed across western Europe, populations are shrinking in certain regions [58]. For example, on the Iberian Peninsula, *H. morsus-ranae* now occurs in a single area [59], a range constriction that took place over the last century. Probable causes include habitat degradation due to farming and industrial activities as well as changes in the hydrochemical characteristics and hydrological regimes of water bodies [58]. At the same time, *H. morsus-ranae* has become an AIS in the north-eastern USA and southern Canada [60,61].

Hence, given the plant's recent range expansion in Europe, it is crucial to maximise efforts to rapidly detect its presence. By acting now, during the early stages of the invasion, it is more likely that *H. laevigata* can be controlled [62,63]. At particular risk is the Mediterranean, where conditions in rivers, streams, lakes, ponds, reservoirs, and canals favour the arrival and establishment of *H. laevigata*.

In this regard, it should be added that the introduction and spread of IAS are a major concern for the European Union; for those reasons, Regulation (EU) No 1143/2014 of the European Parliament and of the Council of 22 October 2014 on the prevention and management of the introduction and spread of invasive alien species was published, which is being developed and updated through different implementing regulations and which establishes rules to prevent, minimize, and mitigate the adverse effects of IAS. This regulation is associated with a list of IAS: “the Union list”, which is the fundamental instrument for implementing the aforementioned regulation. *Hydrocharis laevigata* is not included in this list. Taking into account the problems that this species causes to native biodiversity and to human activities in other parts of the world, it is hoped that the information contained in this brief note will serve as a starting point to initiate actions to include *Hydrocharis laevigata* in the “the Union list”, something that will undoubtedly benefit the conservation of European biodiversity and reduce its harmful effects on human activities.

4. Materials and Methods

This study was based on an extensive literature analysis, data from field surveys, and the examination of herbarium specimens (at MA, SEV, COFC; see Appendix A).

The Geospatial Conservation Assessment Tool (<http://geocat.kew.org/>, accessed on 1 November 2022) was used to create the map (Figure 2) showing where *H. laevigata* has been observed (location descriptions: Appendices A and B).

Funding: This research was funded by the Andalusian Regional Government Research Group PLACCA (RNM116).

Data Availability Statement: Not applicable.

Acknowledgments: The author thanks Norbert Riezing from Hungary and Artur Pliszko from Poland for providing information about *H. laevigata* occurrence in those two countries. The author also expresses his appreciation to Artur Pliszko for furnishing photos of the Polish population.

Conflicts of Interest: The author declares no conflict of interest.

Appendix A

Table A1. Information on the occurrence of *Hydrocharis laevigata* in Europe compiled by GBIF (accessed: November 2022).

gbifID	Locality	State Province	DecimalLatitude	Decimal Longitude	Event Date	Basis of Record
3400985464	Aalsmeer, commercial greenhouse	Noord-Holland	52.285.699	4.818.572	2019-03-06	PRESERVED_SPECIMEN
3400971442	Bleiswijk	Zuid-Holland	52.012.874	4.549.629	2016-05-13	PRESERVED_SPECIMEN
3400962451	Bleiswijk	Zuid-Holland	52.012.874	4.549.629	2016-05-13	PRESERVED_SPECIMEN
1569965647	Ganshoren	Brussels Capital Region	50.8.804	430.617	2013-09-01	HUMAN_OBSERVATION
1569965625	Ganshoren	Brussels Capital Region	50.88.037	430.569	2013-08-25	HUMAN_OBSERVATION
1569965614	Ganshoren	Brussels Capital Region	50.88.004	430.411	2013-09-19	HUMAN_OBSERVATION
1569965606	Ganshoren	Brussels Capital Region	50.88.037	430.954	2013-09-11	HUMAN_OBSERVATION
1569965603	Ganshoren	Brussels Capital Region	50.88	430.454	2013-09-01	HUMAN_OBSERVATION
1065300305	Bergianska Botanick Garden	Stockholms stad	59.37	180.471	Not shown	LIVING_SPECIMEN

Appendix B

Table A2. Herbarium specimens of *Hydrocharis laevigata* relating to the European records.

Location	Country/Province	Date	Herbarium Codes
Aalsmeer, commercial greenhouse	Noord-Holland/The Netherlands	2019-03-06	WAG0452864
Bleiswijk	Zuid-Holland/The Netherlands	2016-05-13	WAG0453625, WAG0453626
Agatowa street	Poland/Krakow	2020-09	KRA 0557636, 0557637, 0557638;
Near Martin Gonzalo Dam	Spain/Cordoba	2018-06-10	COFC 65837
Colmenar Viejo, Grajal Dam	Spain/Madrid	2019-09-28	MA-01-00947148
Colmenar Viejo, Grajal Dam	Spain/Madrid	2021-09-08	MA-01-951386
River Guadaira, near hermitage of the Virgin of Valme	Spain/Seville	2021-06-08	SEV 289200
River Guadaira, near hermitage of the Virgin of Valme	Spain/Seville	2021-08-30	SEV 289203

References

- Jenkins, M. Prospects for biodiversity. *Science* **2003**, *302*, 1175–1177. [CrossRef]
- Nel, J.L.; Reyers, B.; Roux, D.J.; Cowling, R.M. Expanding protected areas beyond their terrestrial comfort zone: Identifying spatial options for river conservation. *Biol. Conserv.* **2009**, *142*, 1605–1616. [CrossRef]
- Markovic, D.; Carrizo, S.; Freyhof, J.; Cid, N.; Lengyel, S.; Scholz, M.; Kasperdus, H.; Darwall, W. Europe's freshwater biodiversity under climate change: Distribution shifts and conservation needs. *Divers. Distrib.* **2014**, *20*, 1097–1107. [CrossRef]
- Panlasigui, S.; Davis, A.J.; Mangiante, M.J.; Darling, J.A. Assessing threats of non-native species to native freshwater biodiversity: Conservation priorities for the United States. *Biol. Conserv.* **2018**, *224*, 199–208. [CrossRef]
- Klimaszyk, P.; Gołdyn, R. Water Quality of Freshwater Ecosystems in a Temperate Climate. *Water* **2020**, *12*, 2643. [CrossRef]
- Janssen, J.A.M.; Rodwell, J.S.; García Criado, M.; Gubbay, S.; Haynes, T.; Nieto, A.; Sanders, N.; Landucci, F.; Loidi, J.; Ssymank, A.; et al. *European Red List of Habitats. Part 2. Terrestrial and Freshwater Habitats*; Publications Office of the European Union: Luxembourg, 2016; pp. 1–34. Available online: <https://data.europa.eu/doi/10.2779/091372> (accessed on 1 November 2022).
- Grzybowski, M.; Glińska-Lewczuk, K. Principal threats to the conservation of freshwater habitats in the continental biogeographical region of Central Europe. *Biodivers. Conserv.* **2019**, *28*, 4065–4097. [CrossRef]
- Carrapiço, F. *Azolla em Portugal*; Instituto da Água Ministério do Ambiente e do Ordenamento do Território Centro de Biologia Ambiental: Lisbon, Portugal, 2001. Available online: <http://azolla.fc.ul.pt/documents/AzollaPortugal.pdf> (accessed on 1 November 2022).
- Monteiro, A.; Moreira, I.; Santos, A.C.; Serrasqueiro, P.M. *Gestão do Jacinto-Aquático (Eichhornia crassipes) na Lezíria Grande de Vila Franca de Xira*; Anais do Instituto Superior de Agronomia: Lisbon, Portugal, 2003. Available online: <https://www.repository.utl.pt/bitstream/10400.5/2288/1/REP-ANAIS-297.pdf> (accessed on 1 November 2022).
- Almeida, A.S.R.L. Avaliação da Qualidade da Água na Pateira de Fermentelos: Caso de Estudo na Área de Projecto. Master Thesis, Universidade de Aveiro, Aveiro, Portugal, 2006. Available online: <http://hdl.handle.net/10773/4662> (accessed on 1 November 2022).
- García Murillo, P.; Fernández Zamudio, R.; Cirujano Bracamonte, S.; Sousa Martín, A.; Espinar, J.M. The invasion of Donana National Park (SW Spain) by the mosquito fern (*Azolla filiculoides* Lam). *Limnetica* **2007**, *26*, 243–250. [CrossRef]
- Ruiz Téllez, T.; de Rodrigo López, E.M.; Lorenzo Granado, G.; Albano Pérez, E.; Morán López, R.; Sánchez Guzmán, J.M. The water hyacinth, *Eichhornia crassipes*: An invasive plant in the Guadiana River Basin (Spain). *Aquat. Invasions* **2008**, *3*, 42–53. [CrossRef]
- Ministerio para la Transición Ecológica. *Estrategia de Gestión, Control y Posible Erradicación del Camalote (Eichhornia crassipes). Technical report*; Ministerio para la Transición Ecológica: Madrid, Spain, 2019. Available online: https://www.miteco.gob.es/va/biodiversidad/publicaciones/estrategiadegestioneichhorniacrassipes3deoctubre2019_tcm39-502314.pdf (accessed on 1 November 2022).
- Piñero Rodríguez, M.J.; Fernandez Zamudio, R.; Arribas, R.; Gómez Mestre, I.; Diaz Paniagua, C. The invasive aquatic fern *Azolla filiculoides* negatively impacts water quality, aquatic vegetation and amphibian larvae in Mediterranean environments. *Biol. Invasions* **2021**, *23*, 755–769. [CrossRef]
- Pliszko, A.; Górecki, A. First record of *Limnobium laevigatum* (Humb. & Bonpl. ex Willd.) Heine (Hydrocharitaceae) and *Pontederia crassipes* Mart. (Pontederiaceae) in Poland. *Biol. Invasions Rec.* **2021**, *10*, 537–543. [CrossRef]
- Bernardini, B.; Lucchese, F. New phylogenetic insights into Hydrocharitaceae. *Ann. Bot.* **2018**, *8*, 45–58. [CrossRef]

17. Li, Z.Z.; Lehtonen, S.; Gichira, A.W.; Martins, K.; Efremov, A.; Wang, Q.F.; Chen, J.M. Plastome phylogenomics and historical biogeography of aquatic plant genus *Hydrocharis* (Hydrocharitaceae). *BMC Plant Biol.* **2022**, *22*, 106. [CrossRef] [PubMed]
18. WFO. *Hydrocharis laevigata* (Humb. & Bonpl. ex Willd.) Byng & Christenh. World Flora Online Data. 2021. Available online: <http://www.worldfloraonline.org/taxon/wfo-0001427027> (accessed on 1 November 2022).
19. Les, D.H. *Aquatic Plants of North America: Ecology, Life History and Systematics*; CRC Press: Boca Raton, FL, USA, 2020; Volume 2, pp. 56–58.
20. Tanaka, N.; Setoguchi, H.; Murata, J. Phylogeny of the family Hydrocharitaceae inferred from *rbcL* and *matK* gene sequence data. *J. Plant Res.* **1997**, *110*, 329–337. [CrossRef]
21. Acevedo-Rodríguez, P.; Strong, M.T. Monocotyledons and gymnosperms of Puerto Rico and the Virgin Islands. *Contrib. U. S. Natl. Herb.* **2005**, *52*, 1–415.
22. Cook, C.D.K.; Urmi-König, K. A revision of the genus *Limnobia* including *Hydromystria* (Hydrocharitaceae). *Aquat. Bot.* **1983**, *17*, 1–27. [CrossRef]
23. Lot, A.; Medina Lemos, R.; Chiang, F. *Plantas Acuáticas Mexicanas, Una Contribución a la Flora de México, Volumen I, Monocotiledóneas*; Universidad Nacional Autónoma de México—Instituto de Biología: Coyoacán, Mexico, 2013; 398p.
24. Lowden, R.M. Floral variation and taxonomy of *Limnobia* LC Richard (Hydrocharitaceae). *Rhodora* **1992**, *94*, 111–134.
25. APHIS. Weed Risk Assessment for *Limnobia laevigata* (Humb. & Bonpl. ex Willd.) Heine (Hydrocharitaceae)—South American Spongeplant, United States Department of Agriculture, Animal and Plant Health Inspection Service, Raleigh, USA. 2013. Available online: https://www.aphis.usda.gov/plant_health/plant_pest_info/weeds/downloads/wra/Limnobia_laevigatum_WRA.pdf (accessed on 1 November 2022).
26. Martín, C.S.; Contreras, D.; Vidal, O.; Solís, J.L.; Ramírez, C. Distribución en Chile y colonización del río Cayumapu (Valdivia) por el macrófito acuático invasor *Limnobia laevigata*. *Gayana Bot.* **2021**, *78*, 65–76. [CrossRef]
27. Hernández, J.; Rangel, J.O. La vegetación del humedal de Jaboque (Bogotá, DC). *Caldasia* **2009**, *31*, 355–379.
28. Ruiz-Merino, M.; Campos-Cuellar, R.; Germán-Gómez, A.; Aponte, H. Características, historia natural y aplicaciones de *Hydrocharis laevigata*: Una revisión. *Caldasia* **2022**, *44*, 432–441. [CrossRef]
29. SERCUL (2020). *Amazon Frogbit—Limnobia laevigata Ex-Situ Experiments and Field Investigations*; South East Regional Centre for Urban Landcare Inc. (SERCUL). Perth NRM.: Perth, Australia, 2020. Available online: <https://www.sercul.org.au/wp-content/uploads/2021/09/Amazon-Frogbit-Research-Report-2020web.pdf> (accessed on 1 November 2022).
30. U.S. National Plant Germplasm System. USDA, Agricultural Research Service, Germplasm Resources Information Network (GRIN Taxonomy). National Germplasm Resources Laboratory, Beltsville, USA. 2022. Available online: <https://npgsweb.ars-grin.gov/gringlobal/taxon/taxonomydetail?id=419808> (accessed on 1 November 2022).
31. Zuloaga, F.O.; Morrone, O.; Belgrano, M.J. *Catálogo de las Plantas Vasculares del Cono Sur (Argentina, Sur de Brasil, Chile, Paraguay y Uruguay). Volumen 1: Pteridophyta, Gymnospermae y Monocotyledoneae*; Smithsonian: Washington, DC, USA, 2008; 983p.
32. Pérez, C.; Seewald, P. *Limnobia laevigata* (Humb. & Bonpl. ex Willd.) Heine (Alismatales, Hydrocharitaceae), una nueva especie invasora en la cuenca del Río Negro, provincia de río negro. *Argent. His. Nat.* **2019**, *9*, 211–218.
33. Kadono, Y. Alien aquatic plants naturalized in Japan: History and present status. *Glob. Environ. Res.* **2004**, *8*, 163–169.
34. Haynes, R.R.; Holm-Nielsen, L.B. The genera of Hydrocharitaceae in the Southeastern United States. *Harv. Pap. Bot.* **2001**, *5*, 201–275.
35. Howard, G.W.; Hyde, M.A.; Bingham, M.G. Alien *Limnobia laevigata* (Humb. & Bonpl. ex Willd.) Heine (Hydrocharitaceae) becoming prevalent in Zimbabwe and Zambia. *Bioinvasions Rec.* **2016**, *5*, 221–225. [CrossRef]
36. Pratt, P.D.; Moran, P.J.; Pitcairn, M.; Reddy, A.M.; O'Brien, J. Biological control of invasive plants in California's Delta: Past, present, and future. *J. Aquat. Plant Manag.* **2021**, *59*, 55–66.
37. Riezinger, N. Újabb adventív vízínövény Magyarországon: *Limnobia laevigata* (Hydrocharitaceae). *Kitaibelia* **2019**, *24*, 9–15. [CrossRef]
38. Corro, M.; Izuzquiza, Á.; Cirujano, S. Primera cita de la especie potencialmente invasora *Limnobia laevigata* (Humb. & Bonpl. ex Willd.) Heine (Hydrocharitaceae) en la Península Ibérica. *BV News Publ. Científicas* **2019**, *8*, 75–80.
39. Martínez-Sagarra, G.; García Murillo, P.; Devesa, J.A. *Hydrocharis laevigata* (Humb. & Bonpl. ex Willd.) Byng & Christenh. (Hydrocharitaceae), novedad para la flora de Andalucía (España). *Acta Bot. Malacit.* **2021**, *46*, 93–94. [CrossRef]
40. García Murillo, P.; Boniquito, J.M.; Gutiérrez González, D.; Rodríguez Hiraldo, C. Nueva localidad ibérica de la planta exótica invasora *Hydrocharis laevigata* (Humb. & Bonpl. ex Willd.) Byng & Christenh. *Folia Bot. Extremad.* **2021**, *15*, 141–145.
41. GBIF. *Hydrocharis laevigata* (Humb. & Bonpl. ex Willd.) Byng & Christenh. GBIF Backbone Taxonomy, Checklist Dataset, GBIF Secretariat. 2022. Available online: <https://www.gbif.org/dataset/d7dddbf4-2cf0-4f39-9b2a-bb099caae36c> (accessed on 1 November 2022). [CrossRef]
42. Larson, L. *Spongeplant: A New Aquatic Weed Threat in Delta*. Cal-IPC News Spring 2011, Berkeley, USA. 2011. Available online: https://www.cal-ipc.org/wp-content/uploads/2017/03/Cal-IPC_News_2011Spring-6.pdf#page=4 (accessed on 1 November 2022).
43. Aponte, H. Productividad de *Limnobia laevigata* (Hydrocharitaceae) bajo condiciones de laboratorio. *Polibotánica* **2017**, *44*, 157–166.
44. Van de Witte, Y. *Limnobia laevigata* (South American spongeplant); CABI Compendium; CABI International: Wallingford, UK, 2022. [CrossRef]

45. DiTomaso, J.M.; Kyser, G.B.; Oneto, S.R.; Wilson, R.G.; Orloff, S.B.; Anderson, L.W.; Wright, S.D.; Roncoroni, J.A.; Miller, T.L.; Prather, T.S. *Weed Control in Natural Areas in the Western United States*; University of California Weed Research and Information Center: Davis, CA, USA, 2013; 544p.
46. Willis, B.E.; Heilman, M.A.; Bishop, W.M.; Shuler, S.W. Evaluation of Multiple Herbicides for Control of Sponge Plant (*Limnium laevigatum*). *J. Geosci. Environ. Prot.* **2018**, *6*, 56–64. [\[CrossRef\]](#)
47. California Department of Parks and Recreation. Water Hyacinth Control Program and Sponge Plant Control Program Aquatic Pesticide Application Plan. 2013. Available online: <https://dbw.parks.ca.gov/pages/28702/files/WHCP%20SCP%20Aquatic%20Pesticide%20Application%20Plan%202013.pdf> (accessed on 1 November 2022).
48. Bol, R.; Gruau, G.; Mellander, P.E.; Dupas, R.; Bechmann, M.; Skarbøvik, E.; Bieroza, M.; Djodjic, F.; Glendell, M.; Jordan, P.; et al. Challenges of reducing phosphorus based water eutrophication in the agricultural landscapes of Northwest Europe. *Front. Mar. Sci.* **2018**, *5*, 276. [\[CrossRef\]](#)
49. Charlton, M.B.; Bowes, M.J.; Hutchins, M.G.; Orr, H.G.; Soley, R.; Davison, P. Mapping eutrophication risk from climate change: Future phosphorus concentrations in English rivers. *Sci. Total Environ.* **2018**, *613*, 1510–1526. [\[CrossRef\]](#) [\[PubMed\]](#)
50. Paredes, I.; Ramírez, F.; Aragonés, D.; Bravo, M.A.; Forero, M.G.; Green, A.J. Ongoing anthropogenic eutrophication of the catchment area threatens the Doñana World Heritage Site (South-west Spain). *Wetl. Ecol. Manag.* **2021**, *29*, 41–65. [\[CrossRef\]](#)
51. Bonometto, A.; Ponis, E.; Cacciato, F.; Riccardi, E.; Pigozzi, S.; Parati, P.; Novello, M.; Ungaro, N.; Acquavita, A.; Manconi, P.; et al. A New Multi-Index Method for the Eutrophication Assessment in Transitional Waters: Large-Scale Implementation in Italian Lagoons. *Environments* **2022**, *9*, 41. [\[CrossRef\]](#)
52. Mack, L.; Andersen, H.E.; Beklioglu, M.; Bucak, T.; Couture, R.M.; Cremona, F.; Ferreira, M.T.; Hutchins, M.G.; Mischke, U.; Molina-Navarro, E.; et al. The future depends on what we do today—Projecting Europe’s surface water quality into three different future scenarios. *Sci. Total Environ.* **2019**, *668*, 470–484. [\[CrossRef\]](#) [\[PubMed\]](#)
53. De Vries, W.; Schulte-Uebbing, L.; Kros, H.; Voogd, J.C.; Louwagie, G. Spatially explicit boundaries for agricultural nitrogen inputs in the European Union to meet air and water quality targets. *Sci. Total Environ.* **2021**, *786*, 147283. [\[CrossRef\]](#)
54. Rodríguez-Garrito, E.C.; Paz-Gallardo, A.; Plaza, A. Monitoring the Spatiotemporal Distribution of Invasive Aquatic Plants in the Guadiana River, Spain. *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.* **2023**, *16*, 228–241. [\[CrossRef\]](#)
55. Van Rees, C.B.; Waylen, K.A.; Schmidt-Kloiber, A.; Thackeray, S.J.; Kalinkat, G.; Martens, K.; Domisch, S.; Lillebø, A.I.; Hermoso, V.; Grossart, H.P.; et al. Safeguarding freshwater life beyond 2020: Recommendations for the new global biodiversity framework from the European experience. *Conserv. Lett.* **2021**, *14*, e12771. [\[CrossRef\]](#)
56. Troia, A.; Ilardi, V.; Oddo, E. Monitoring of alien aquatic plants in the inland waters of Sicily (Italy). *Webbia* **2020**, *75*, 77–83. [\[CrossRef\]](#)
57. Gross, E.M.; Groffier, H.; Pestelard, C.; Hussner, A. Ecology and Environmental Impact of *Myriophyllum heterophyllum*, an Aggressive Invader in European Waterways. *Diversity* **2020**, *12*, 127. [\[CrossRef\]](#)
58. Efremov, A.N.; Grishina, V.S.; Kislov, D.E.; Mesterházy, A.; Toma, C. The genus *Hydrocharis* L. (Hydrocharitaceae): Distribution features and conservation status. *Bot. Pac.* **2020**, *9*, 83–94. [\[CrossRef\]](#)
59. García-Murillo, P.; Cirujano, S.; Medina, L.; Sousa, A. ¿Se extinguirá *Hydrocharis morsus-ranae* L. de la Península Ibérica? *Port. Acta Biol.* **2000**, *19*, 149–158.
60. Catling, P.M.; Dore, W.G. Status and identification of *Hydrocharis morsus-ranae* and *Limnium spongia* (Hydrocharitaceae) in northeastern North America. *Rhodora* **1982**, *84*, 523–545.
61. Zhu, B.; Ottaviani, C.C.; Naddafi, R.; Dai, Z.; Du, D. Invasive European frogbit (*Hydrocharis morsus-ranae* L.) in North America: An updated review 2003–16. *J. Plant Ecol.* **2018**, *11*, 17–25. [\[CrossRef\]](#)
62. Epanchin-Niell, R.; Thompson, A.L.; Treake, T. Public contributions to early detection of new invasive pests. *Conserv. Sci. Pract.* **2021**, *3*, e422. [\[CrossRef\]](#)
63. Larson, E.R.; Graham, B.M.; Achury, R.; Coon, J.J.; Daniels, M.K.; Gambrell, D.K.; Jonassen, K.L.; King, G.D.; LaRacune, N.; Perrin-Stowe, T.I.; et al. From eDNA to citizen science: Emerging tools for the early detection of invasive species. *Front. Ecol. Environ.* **2020**, *18*, 194–202. [\[CrossRef\]](#)

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.