plants

MDPI

Article

Methyl Jasmonate Synergistically Enhances Phenolic Compounds and Reactive Oxygen Species Scavenging to Alleviate Drought-Induced Oxidative Damage in *Ilex rotunda*

Junhan Guo [1,†], Omer Conteh [1,†], Ye Zheng [1], Mingzheng Duan [2,*], Huwei Yuan [1,3], Yongqing He [4], Minfen Yu [5], Wenjun Dai [1,3], Yihui Li [1,3], Songting Zhang [1,3], Qinyuan Shen [1,3], Liangye Huang [1,3], Bingsong Zheng [1,3,*] and Muhammad Junaid Rao [1,3,*]

[1] State Key Laboratory for Development and Utilization of Forest Food Resources, Zhejiang A&F University, Hangzhou 311300, China; guojh@stu.zafu.edu.cn (J.G.); contehomar343@stu.zafu.edu.cn (O.C.); 2024102012044@stu.zafu.edu.cn (Y.Z.); hwyuan@zafu.edu.cn (H.Y.); 2023602122015@stu.zafu.edu.cn (W.D.); 2024602122054@stu.zafu.edu.cn (Y.L.); 2024102012041@stu.zafu.edu.cn (S.Z.); 2024202011016@stu.zafu.edu.cn (Q.S.); 2023602121036@stu.zafu.edu.cn (L.H.)

[2] Advanced Institute of Ecological Agriculture and Biodiversity on the Yunnan-Guizhou Plateau, Zhaotong University, Zhaotong 657000, China

[3] Zhejiang Key Laboratory of Non-wood Forest Products Quality Regulation and Processing Utilization, Zhejiang A&F University, Hangzhou 311300, China

[4] Natural Resources and Planning Bureau of Qingyuan County, Lishui 323800, China; 15857810359@163.com

[5] Ningbo Forestry Development Center, Ningbo 315042, China; 15957495963@163.com

* Correspondence: mjr@zafu.edu.cn (M.J.R.); bszheng@zafu.edu.cn (B.Z.); duanmingzheng@ztu.edu.cn (M.D.); Tel.: +86-1333-600-4216 (M.J.R.)

† These authors contributed equally to this work.

Abstract

Drought stress severely compromises the physiological integrity and secondary metabolism of medicinal plants. This study integrated physiological, biochemical, and ultra-performance liquid chromatography–tandem mass spectrometry (UPLC-MS/MS) analyses to investigate the effects of exogenous methyl jasmonate (MeJA) on drought-stressed *Ilex rotunda* seedlings. Drought reduced relative water content by 29% and chlorophyll by >50%, while elevating H_2O_2 (76%) and malondialdehyde (120%). MeJA application mitigated these impairments, reducing oxidative markers by 25% and enhancing non-enzymatic antioxidant capacity, as shown by a 74% increase in DPPH radical scavenging activity and a 141% rise in total phenolic content. Hierarchical clustering analysis (HCA), principal component analysis (PCA), and orthogonal partial least squares-discriminant analysis (OPLS-DA), and pathway mapping confirmed a significant reprogramming of the phenolic metabolome, particularly within phenylpropanoid and benzoate biosynthesis pathways. Drought + MeJA-treated plants exhibited a distinct and enriched profile compared to both well-watered control and drought-stressed groups. This reprogramming specifically elevated key hydroxycinnamates, including verbascoside and neochlorogenic acid (increased by 50% and 52%, respectively), while suppressing alternative phenolic branches. These findings demonstrate that MeJA orchestrates a shift from enzymatic scavenging to a potent metabolite-based antioxidant system, positioning it as an effective elicitor for enhancing drought resilience and enriching the high-value phytochemicals in *I. rotunda*.

Keywords: *Ilex rotunda*; drought stress; methyl jasmonate; oxidative stress; phenolic metabolism; antioxidant defense; metabolomics

Check for updates

Academic Editor: Koji Mikami

Received: 26 January 2026
Revised: 15 February 2026
Accepted: 17 February 2026
Published: 19 February 2026

Copyright: © 2026 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license.

1. Introduction

Ilex rotunda Thunb. (Kurogane holly), a member of the Aquifoliaceae family, is an evergreen tree widely distributed across East and Southeast Asia, including China, Japan, and Korea [1]. Traditionally valued in East Asian medicine, its leaves, bark, and fruits are rich in phenolics, flavonoids, and triterpenoids with diverse pharmacological activities such as antioxidant, anti-inflammatory, and hepatoprotective effects [1,2]. Recent phytochemical investigations have identified *I. rotunda* leaves as a promising source of bioactive phenolic compounds, including chlorogenic acid derivatives, saponins, and caffeic acid derivatives, which exhibit strong radical scavenging and cytoprotective properties [3,4]. These compounds underpin the plant's growing industrial relevance in the nutraceutical, cosmeceutical, and functional food sectors, where natural antioxidants are increasingly sought as safer alternatives to synthetic additives [5]. Consequently, *I. rotunda* has emerged as a potential medicinal crop for sustainable production of high-value phytochemicals, aligning with global efforts to develop plant-based antioxidant resources.

However, the productivity and phytochemical quality of *I. rotunda* are highly sensitive to environmental stressors, particularly drought. As a perennial woody species, *I. rotunda* depends on continuous water availability to sustain photosynthetic efficiency and secondary metabolism [6]. Drought stress disrupts this balance by inducing stomatal closure, reducing CO_2 assimilation, and impairing chlorophyll biosynthesis, ultimately leading to growth inhibition and yield loss [7]. More critically, drought triggers excessive accumulation of reactive oxygen species (ROS) such as hydrogen peroxide (H_2O_2) and superoxide radicals, which cause oxidative damage to lipids, proteins, and nucleic acids. Plants counteract this oxidative burden through enzymatic defenses such as superoxide dismutase (SOD), catalase (CAT), and peroxidase (POD), and non-enzymatic antioxidants like phenolic compounds and carotenoids [8]. Yet, prolonged or severe drought often overwhelms these systems, resulting in oxidative stress and metabolic reprogramming that compromise both plant vitality and the accumulation of industrially relevant metabolites [9]. In *Ilex* species, drought has been shown to alter the biosynthesis of phenolic acids and flavonoids, but the direction and magnitude of these changes vary depending on stress intensity and duration [10–12]. Thus, developing strategies to mitigate drought-induced oxidative stress while maintaining or enhancing metabolite production is essential for the sustainable cultivation of *I. rotunda* as an industrial crop.

Among the various strategies explored, elicitor-based priming has gained prominence as a sustainable approach to enhance plant resilience and secondary metabolite biosynthesis [13]. Methyl jasmonate (MeJA), a volatile oxylipin derivative of jasmonic acid, functions as a key signaling molecule in plant defense and stress adaptation [14]. Exogenous application of MeJA has been widely reported to modulate physiological and biochemical responses under abiotic stresses, including drought, salinity, and temperature extremes [14,15]. Mechanistically, MeJA primes plants by activating transcriptional networks associated with antioxidant defense and secondary metabolism, particularly the phenylpropanoid pathway, which governs the biosynthesis of phenolic acids, flavonoids, and lignins [16]. In several species, MeJA treatment has been shown to enhance drought tolerance by reducing ROS accumulation, stabilizing photosynthetic pigments, and elevating antioxidant enzyme activities [17]. Moreover, MeJA acts as a potent inducer of phenolic metabolism, stimulating the accumulation of hydroxycinnamic acid derivatives, flavonols, and glycosylated phenolics in crops such as *Chrysanthemum morifolium* Ramat, *Camellia vietnamensis* Huang, and *Vigna radiata L.* [18–20]. These findings collectively underscore MeJA's dual role as a stress mitigator and metabolic enhancer, making it an attractive tool for improving both stress resilience and phytochemical yield in medicinal crops.

Despite these advances, the interactive effects of drought and MeJA on *Ilex rotunda* remain largely unexplored. While prior studies have characterized the general antioxidant potential of *I. rotunda* extracts [2,21], there is an obvious lack of integrative analyses linking physiological performance, oxidative stress markers, and metabolomic reprogramming under combined drought and MeJA treatments. Specifically, how MeJA modulates the phenolic metabolome of *I. rotunda* leaves under drought conditions remains unknown. This gap limits our understanding of the biochemical mechanisms through which MeJA confers drought tolerance and constrains the rational application of elicitation strategies for industrial metabolite enhancement. Given the commercial significance of phenolic antioxidants in *Ilex* species [2–4], elucidating this regulatory interplay is critical for optimizing cultivation and processing protocols aimed at maximizing bioactive compound yield under environmental stress.

To address this knowledge gap, the present study integrates physiological, biochemical, and metabolomic approaches to investigate the effects of exogenous MeJA on drought-stressed *Ilex rotunda* seedlings. Specifically, we aimed to (i) assess the extent to which MeJA mitigates drought-induced physiological impairment and oxidative stress, (ii) evaluate its influence on antioxidant enzyme activities and non-enzymatic defense capacity, and (iii) characterize the reprogramming of phenolic metabolites using UPLC–MS/MS-based profiling. By coupling physiological and metabolomic data, this study provides a mechanistic understanding of how MeJA orchestrates drought resilience and secondary metabolism in *I. rotunda*. Furthermore, it explores the potential of MeJA elicitation as a biotechnological strategy to enrich *I. rotunda* leaves in specific hydroxycinnamic acid and glycosylated phenolic derivatives with high industrial value.

2. Results

2.1. MeJA Mitigates Drought Effects on Physiology and Enhances Secondary Metabolism

Drought stress severely impaired leaf physiology in *Ilex rotunda*, causing visible wilting and chlorosis (Figure 1A) and reducing relative water content (RWC) by 29% (Figure 1B). This water deficit precipitated a notable 52–54% decline in chlorophyll a, b, and total chlorophyll (Figure 1C). Foliar application of MeJA significantly mitigated these effects, partially restoring RWC and limiting chlorophyll loss to only ~18% relative to the control. In contrast, photoprotective and antioxidant secondary metabolites exhibited a significant increase: drought alone raised carotenoid and total phenolic contents by 31% and 50%, respectively, while MeJA application under drought amplified these accumulations to 83% and 141% above control levels (Figure 1D,E), indicating a shift toward non-enzymatic defense compounds.

2.2. MeJA Alleviates Oxidative Stress and Modulates Antioxidant Defense

Drought stress in *Ilex rotunda* leaves induced severe oxidative stress, marked by a 76% increase in hydrogen peroxide (H_2O_2) (Figure 2A) and a 120% rise in malondialdehyde (MDA) content (Figure 2B) compared to the control, indicating significant membrane damage. In response, antioxidant enzyme activities increased significantly, SOD by 43%, CAT by 73%, and POD by 76%, reflecting an enzymatic defense activation (Figure 2C–E). Application of MeJA under drought conditions alleviated oxidative stress, reducing H_2O_2 and MDA by 25% each relative to drought-alone plants and lowering enzyme activities to intermediate levels. Notably, MeJA treatment uniquely enhanced the non-enzymatic antioxidant capacity, elevating DPPH radical scavenging by 74% over the control and 34% over drought-stressed plants (Figure 2F), indicating a shift toward a more efficient, metabolite-based defense strategy that correlated with reduced oxidative damage.



Figure 1. Methyl Jasmonate mitigates drought-induced inhibition of chlorophyll and preserves morphological integrity in *Ilex rotunda* leaves. (**A**) Representative images of whole plants (line 3.5 cm), (**B**) Relative water content (%), (**C**) Chlorophyll contents (mg/g), (**D**) Total carotenoids (mg/g), (**E**) Total phenols (mg/g) from *Ilex rotunda* subjected to three treatments: Control (CK), Drought (D), and Drought + MeJA (DMeJA). Drought stress induced severe wilting, leaf curling, and chlorosis. Data are presented as mean ± SD (n = 6). Different lowercase letters (a,b,c) indicate significant differences among treatments ($p < 0.05$, Duncan's multiple range test).

2.3. Drought and Methyl Jasmonate Significantly Remodel the Phenolic Profile

Metabolomic analysis revealed that the 324 phenolic compounds common to all treatments (Figure 3A), resulting in strong positive correlations between the groups (Figure 3B). However, drought stress significantly alters 68 compounds compared to the CK, with a

strong upregulation (47 upregulated vs. 21 downregulated) (Figure 3B). Application of MeJA under drought (DMeJA) shared a similar number of altered compounds versus CK (50 total), resulting in an equal number of up- and down-regulated compounds relative to the control (27 vs. 23). Furthermore, the direct comparison between D and DMeJA showed that MeJA application specifically upregulated 45 and downregulated 13 phenolic compounds, indicating an active hormonal reprogramming of the drought-induced phenolic metabolism.

2.4. Hierarchical Clustering Analysis Reveals Treatment-Specific Phenolic Accumulation Patterns

Hierarchical clustering analysis (HCA) of the 119 significantly altered phenolic compounds (VIP > 1) clearly delineated the three treatments into distinct clusters (Supplementary Table S2), confirming a significant alteration of the phenolic metabolome under drought and MeJA application (Figure 4A). The heatmap visualization revealed pronounced, treatment-specific accumulation trends. One major compound cluster, containing derivatives such as 3-O-feruloylquinic acid, grevilloside F, and 1-caffeoyl-beta-D-glucose, exhibited a progressive increase in abundance from CK to D to DMeJA, indicating a strong synergistic effect of drought and MeJA on their biosynthesis. Conversely, another distinct cluster, including compounds like various caffeoylquinic acid isomers, showed peak accumulation under drought stress but a partial reduction with MeJA application (DMeJA), suggesting a moderating effect of the hormone on this specific drought response. A third group, featuring compounds such as salicin and several galloyl derivatives, was notably suppressed by both stress treatments compared to the control. This structured clustering demonstrates that MeJA does not simply reverse the drought response but actively reprograms phenolic metabolism, enhancing a specific subset of compounds while tempering the accumulation of others.



Figure 2. Effects of drought and methyl jasmonate on oxidative stress markers and antioxidant responses in *Ilex rotunda* leaves. (**A**) Hydrogen peroxide (H_2O_2) content. (**B**) Malondialdehyde (MDA) content. (**C**) Superoxide dismutase (SOD) activity. (**D**) Peroxidase (POD) activity. (**E**) Catalase (CAT) activity. (**F**) DPPH radical scavenging capacity. Plants were subjected to three treatments: control (CK), drought (D), and drought with exogenous methyl jasmonate application (DMeJA). Values represent means ± standard deviation (n = 6). Different lowercase letters above bars indicate statistically significant differences among treatments according to Duncan's multiple range test ($p < 0.05$).



Figure 3. Analysis of phenolic compound profiles in *Ilex rotunda* leaves under drought and MeJA treatment. (**A**) Venn diagram illustrating the number of phenolic compounds identified as unique and shared among the Control (CK), Drought (D), and Drought + MeJA (DMeJA) treatments. (**B**) Pearson correlation matrix depicting the overall similarity between the phenolic profiles of the three treatment groups. (**C**) Bar chart of phenolic compounds that were significantly altered across comparison groups, filtered by the criteria: Fold Change > 2, variable importance in projection (VIP) > 1, and $|\mathrm{Log_2(Fold\ Change)}| > 1$.

Principal component analysis (PCA) clearly separated the phenolic profiles of the three treatments, with PC1 and PC2 explaining 64.46% and 22.41% of the total variance, respectively (Figure 4B). Drought-stressed plants (D) were distantly separated from the CK along PC1. The DMeJA treatment formed a distinct cluster in the upper-right quadrant, significantly separated from both CK and D along PC2; *t*-tests confirmed significant differences among all groups along both axes ($p < 0.05$), with D vs. DMeJA showing the highest significance. This demonstrates that MeJA application actively reconfigures the

drought-induced phenolic metabolome into a unique state, rather than merely reverting it toward the control profile.



Figure 4. Multivariate analysis of the phenolic metabolome in *Ilex rotunda* leaves under drought and MeJA treatment. (**A**) Hierarchical clustering analysis (HCA) heatmap of 119 phenolic compounds that were significantly altered (VIP > 1, $p < 0.05$; Table S2). Rows represent individual phenolic compounds, and columns represent the three treatment groups: control (CK), drought (D), and drought with methyl jasmonate (DMeJA). Color intensity indicates relative abundance (peak area) based on the Z-score, from low (green) to high (pink). (**B**) Two-dimensional score plot from principal component analysis (PCA) of the phenolic compounds. Different asterisks indicate significant separation between treatment groups along each principal component axis (* $p < 0.05$, *** $p < 0.001$, **** $p < 0.0001$, *t*-test).

2.5. OPLS-DA Modeling Confirms Reprogramming of the Phenolic Metabolome

OPLS-DA analysis provided a validated model of the treatment-specific phenolic profiles. The score plot showed clear and robust separation among all three groups

(Figure 5A). The first predictive component on *x*-axis (66.9%) effectively separated the CK from the DMeJA, while the orthogonal component on *y*-axis (21.4%) distinguished the drought-stressed group (D), visually demonstrating that MeJA application under drought establishes a unique metabolic state, distinct from both the stress response and the non-stressed condition. The corresponding S-plot identified the key phenolic metabolites driving this separation, with compounds having a variable importance in projection (VIP) > 1 located at the plot extremes (Figure 5B). A permutation test (n = 200) confirmed the model's high robustness and lack of overfitting, as indicated by the significantly high original Q^2 (0.992) and R^2Y (0.997) values compared to all permuted models ($p < 0.005$) and the negative Q^2 regression intercept (Figure 5C). This analysis confirms that drought and MeJA induce targeted, significant reprogramming of specific phenolic compounds.



Figure 5. Orthogonal partial least squares discriminant analysis (OPLS-DA) of the phenolic metabolome in *Ilex rotunda* leaves. (**A**) OPLS-DA score plot displaying separation between the control (CK), drought (D), and drought + MeJA (DMeJA) groups along the predictive component. (**B**) S-plot from OPLS-DA highlighting metabolites that contribute most to group separation (VIP > 1, shown in red; Table S2). Metabolites positioned in the upper-right and lower-left corners represent those with high covariance and correlation to the model. (**C**) Permutation test (n = 200) assessing model validity; the negative Q^2-intercept confirms that the original model is robust and not over-fitted.

2.6. Phenolic Metabolome Is Distinctively Reprogrammed by Drought and Amplified by MeJA

OPLS-DA revealed that drought stress induced a marked bipolar shift in the phenolic profile of *Ilex rotunda* leaves (Supplementary Table S3), with 20 compounds strongly downregulated and 30 significantly upregulated (Figure 6A). Among the most suppressed metabolites were glycosylated and acylated quinic acid derivatives such as 1,3 O di Caffeoyl 4 O glucoside Quinic Acid ($Log_2FC = -6.99$) and 3 O Galloyl Glucose ($Log_2FC = -6.63$). In contrast, drought triggered a pronounced accumulation of compounds including 1 O Galloyl rhamnose ($Log_2FC = 7.36$, fold change >160) and 1 Caffeoylquinic acid (Index: mws0748; $Log_2FC = 6.38$) (Figure 6A), indicating a targeted redirection of phenolic biosynthesis.

Notably, application of methyl jasmonate under drought (DMeJA) further amplified a specific subset of drought responsive metabolites (Supplementary Table S4), with 23 compounds strongly downregulated and 27 significantly upregulated (Figure 6B). While the suppression of certain compounds persisted, the DMeJA treatment obviously elevated hydroxycinnamic acid derivatives such as 1 Caffeoylquinic acid (Index: mws0748; $Log_2FC = 7.00$) and 3,6 Di O caffeoyl glucose (index: Lmsn004970; $Log_2FC = 6.53$) (Figure 6B). Other drought-induced phenolics, including 1 O Galloyl rhamnose and 7 Caffeoylsedoheptulose, were maintained at high levels.

Radar chart analysis of the top 15 altered metabolites between D and DMeJA revealed that MeJA actively redirects phenolics rather than merely amplifying the drought response (Figure 6C). While MeJA moderated the accumulation of some drought-induced compounds, it significantly elevated others. Notably, the complex gallate Maclurin-3-C-(2″-P-Hydroxybenzoyl-6″-Galloyl)Glucoside ($Log_2FC = 6.40$) and 3,6-Di-O-caffeoyl glucose ($Log_2FC = 6.72$) were substantially enhanced in DMeJA. This was accompanied by a marked increase in glycosylated derivatives such as 3′-p-Coumaroyl-2,6-di-acetyl sucrose ($Log_2FC = 4.70$) and 2-Caffeoylrhamnoside ($Log_2FC = 4.58$). This reprogramming indicates that MeJA reshapes the phenolic metabolome into a distinct, stress-adapted profile. For industrial applications, MeJA elicitation under drought offers a targeted approach to enrich leaves in specific glycosylated phenolics, which may offer enhanced bioavailability for nutraceutical and preservative uses.

2.7. Integrated Pathway and Targeted Analysis of Drought- and MeJA-Modulated Phenolic Compounds

Targeted analysis revealed that drought and MeJA treatments distinguished the phenolic profile of *Ilex rotunda* leaves (Figure 7). Drought stress alone severely reduced the abundance of salicin (−69%) and nearly depleted 3,5-dicaffeoylquinic acid (−87%), with MeJA application failing to restore these compounds. In contrast, the DMeJA treatment actively enhanced a suite of hydroxycinnamic acid derivatives. Caffeic acid 4-O-glucoside increased by 88%, while neochlorogenic acid and verbascoside rose by 52% and 50%, respectively. Isochlorogenic acid A also showed a significant 13% increase in DMeJA. This MeJA-mediated upregulation of potent antioxidant phenolics corresponds directly with the 74% increase in whole-leaf DPPH radical scavenging capacity (Figure 2F), indicating a targeted reallocation of phenylpropanoid flux toward non-enzymatic defense.

Additionally, DMeJA further elevated soluble hydroxycinnamate conjugates such as ferulic acid-4-O-glucoside (+33%) and galactosylated p-coumaric acid derivatives (up to +114%) (Figure S1). Conversely, several glycosylated protocatechuic acid derivatives were strongly suppressed, with reductions up to 89%. Together, MeJA under drought does not simply reverse the stress response but actively redirects phenolic metabolism, amplifying a specific array of antioxidant-rich hydroxycinnamate derivatives while concurrently downregulating alternative branches. This targeted reprogramming effectively configures

a metabolome primed for enhanced oxidative stress tolerance and provides a biochemical basis for using drought stress combined with MeJA elicitation as a metabolic switch to enrich *Ilex rotunda* leaves in a specific palette of useful phenolics.



Figure 6. Key phenolic metabolites discriminate treatment groups in *Ilex rotunda* leaves. (**A**) Top 50 differentially abundant compounds in Drought vs. Control (D/CK), ranked by Variable Importance in Projection (VIP). (**B**) Top 50 differentially abundant compounds in Drought + MeJA vs. Control (DMeJA/CK), ranked by VIP. *X*-axis show Log_2(Fold Change) (red: up-regulated, blue: down-regulated); asterisks indicate significance (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$). The black dot plots the VIP score (right axis). (**C**) Radar chart of the top 15 Log_2(Fold Change) values for D vs. DMeJA. Full compound names corresponding to index identifiers are provided in Supplementary Table S1.



Figure 7. Differential regulation of phenolic compounds in *Ilex rotunda* leaves under different treatments. Treatments: control (CK), drought (D), and drought + methyl jasmonate (DMeJA). Bar plots show the mean LC-MS peak area (± SD, n = 3). Different lowercase letters above bars denote statistically significant among treatments ($p < 0.05$, Duncan's multiple range test).

Pathway analysis demonstrated that MeJA under drought stress actively redirected flux within phenylpropanoid and benzoate metabolism (Figure 8). The DMeJA treatment specifically enhanced the accumulation of valuable hydroxycinnamate derivatives and glycosylated phenolics, including verbascoside, neochlorogenic acid, chlorogenate, and

rosmarinate. In contrast, drought alone primarily induced a distinct set of stress-responsive compounds, such as various sinapate and coumarate conjugates (e.g., 4-O-beta-D-Glucosyl-sinapate, p-Coumaroyl-D-glucose). A core group of upstream phenylpropanoid intermediates, like trans-cinnamate and cinnamaldehyde, was elevated in both stressed groups. This delineates a MeJA-mediated strategy that reshapes the phenolic landscape, selectively amplifying branches rich in industrially relevant antioxidant compounds while modulating the general stress response.



Figure 8. Schematic representation of the phenylpropanoid and benzoate metabolism pathways in *Ilex rotunda* leaves, highlighting the reprogramming of phenolic compounds in response to drought (D) and drought combined with methyl jasmonate (DMeJA). Key industrial metabolites, such as verbascoside, neochlorogenic acid, and caffeoylquinic acid derivatives, were significantly upregulated under DMeJA, while alternative branches, including specific glycosylated benzoates, were suppressed. This targeted redirection illustrates how MeJA optimizes the phenolic profile toward a high-value, antioxidant-rich metabolome under stress conditions.

2.8. Correlation Analysis Links Specific Phenolic Compounds with Oxidative Stress Tolerance

To further substantiate the protective roles of individual phenolic metabolites, we performed Pearson correlation analysis between the relative abundances of eight representative phenolic compounds (including verbascoside, neochlorogenic acid, and caffeic acid 4-O-glucoside), total phenolics, total carotenoids, DPPH scavenging capacity, antioxidant enzyme activities, and oxidative stress markers (H_2O_2, MDA). The resulting correlation matrix, visualized as a heatmap with hierarchical clustering (Figure 9), revealed distinct metabolite–trait associations.



Figure 9. Pearson correlation heatmap of phenolic compounds, antioxidant capacity, and oxidative stress markers in *Ilex rotunda* leaves. The heatmap visualizes Pearson correlation coefficients (r) between the relative abundance of key phenolic metabolites (rows) and physiological/biochemical parameters (columns) measured across all treatments (CK, D, DMeJA). Color intensity indicates the strength and direction of correlation: red = positive correlation, blue = negative correlation.

Verbascoside, neochlorogenic acid, caffeic acid 4-O-glucoside, and isochlorogenic acid A displayed strong positive correlations with DPPH capacity ($r > 0.90$), total phenolics ($r > 0.99$), and total carotenoids ($r > 0.99$), and strong negative correlations with H_2O_2 ($r < –0.80$) and MDA ($r < –0.85$). In contrast, salicin and 3,5-dicaffeoylquinic acid showed negative or negligible correlations with these antioxidant traits and were positively correlated with oxidative damage markers. Antioxidant enzyme activities (SOD, POD, CAT) exhibited moderate positive correlations with oxidative stress markers and weak or negative correlations with the MeJA-induced hydroxycinnamates, consistent with the observed shift from enzymatic to non-enzymatic defense. These correlation results provide quantitative evidence that the specific phenolic subclasses enriched by MeJA treatment are functionally associated with enhanced ROS scavenging and reduced lipid peroxidation, thereby directly contributing to drought tolerance.

3. Discussion

The present study demonstrates that exogenous MeJA application effectively mitigates drought-induced physiological impairment in *Ilex rotunda* while simultaneously amplifying a targeted phenolic antioxidant response. This dual action (physiological protection and metabolic enhancement) reflects a coordinated reprogramming of stress defense that integrates both primary and secondary metabolism [7,22]. The findings collectively indicate that MeJA not only alleviates drought-induced oxidative stress but also orchestras a metabolic shift from enzymatic scavenging toward a proactive, metabolite-based antioxi-

dant system, thereby establishing a unique biochemical state optimized for resilience and potential industrial exploitation.

Under drought conditions, *I. rotunda* exhibited typical symptoms of water deficit, including reduced relative water content and chlorophyll degradation (Figure 1). These physiological impairments are consistent with previous observations in woody perennials subjected to dehydration stress, where compromised photosynthetic efficiency and membrane stability are hallmarks of oxidative imbalance [11,12]. The partial restoration of water status and chlorophyll integrity by MeJA treatment underscores its role as a stress-priming signal that stabilizes cellular homeostasis [15]. Jasmonates are known to modulate stomatal behavior, osmolyte accumulation, and chloroplast protection [23], and the present results confirm that MeJA confers similar protective effects in *I. rotunda*, thereby maintaining photosynthetic competence under drought.

Beyond physiological mitigation, the most striking effect of MeJA was the reconfiguration of antioxidant defense. Drought alone triggered a strong activation of enzymatic antioxidants (SOD, CAT, and POD), reflecting a reactive defense against elevated reactive oxygen species (ROS) (Figure 2C–E). However, MeJA-treated plants exhibited reduced enzymatic activity despite lower ROS accumulation (Figure 2A,B), suggesting that oxidative stress was early reduced through alternative mechanisms. The concurrent 74% increase in DPPH radical scavenging capacity (Figure 2F) and the pronounced accumulation of phenolic antioxidants (Figure 7) indicate that MeJA redirected defense investment from enzyme-dependent detoxification toward a metabolite-based antioxidant system. This strategic shift aligns with the concept of "metabolic priming," wherein jasmonate signaling enhances the biosynthesis of redox-active secondary metabolites that provide sustained protection without the high energetic cost of continuous enzyme turnover [24].

The metabolomic data provides compelling evidence for this reprogramming. Multivariate analyses (PCA and OPLS-DA; Figures 4B and 5A) revealed that MeJA-treated plants occupied a distinct metabolic profile, separate from both control and drought-stressed groups, confirming the establishment of a unique biochemical state. This distinct clustering reflects a jasmonate-mediated redirection of phenylpropanoid flux, a hallmark of JA signaling under abiotic stress [25]. Specifically, MeJA amplified the accumulation of hydroxycinnamate derivatives such as caffeic acid 4-O-glucoside, neochlorogenic acid, verbascoside (Figure 7), ferulic acid 4-O-glucoside, and p-coumaric acid derivatives (Figure S1), compounds well-documented for their potent antioxidant and photoprotective properties [26,27]. These metabolites act as ROS scavengers and membrane stabilizers [28], providing a biochemical explanation for the observed reduction in lipid peroxidation (MDA) and hydrogen peroxide content (Figure 2A,B).

Interestingly, MeJA not only enhanced certain phenolic branches but also selectively suppressed others, including glycosylated protocatechuic acid derivatives and specific quinic acid conjugates (Figure 6A,B). Such targeted suppression suggests a regulatory trade-off favoring the synthesis of more redox-active hydroxycinnamates over less reactive phenolics [16,29]. This pattern mirrors jasmonate-induced metabolic channeling reported in *Arabidopsis* and *Camellia* species, where JA signaling redirects phenylalanine flux toward caffeoyl- and feruloyl-based compounds via transcriptional activation of *PAL*, *C4H*, and *4CL* genes while repressing competing glycosylation pathways [16,30,31]. The strong upregulation of caffeoylquinic acid derivatives (e.g., 1-caffeoylquinic acid and 3,6-di-O-caffeoyl glucose; Figure 6B) thus represents a jasmonate-driven optimization of phenolic biosynthesis toward compounds with superior radical-quenching efficiency [32,33].

The hierarchical clustering of 119 significantly altered phenolics (Figure 4A) further supports this interpretation, revealing distinct compound groups that respond synergisti-

cally to drought and MeJA. The progressive enrichment of feruloylquinic acid, grevilloside F, and 1-caffeoyl-β-D-glucose from CK→D→DMeJA treatments exemplifies a cumulative effect of stress and hormonal signaling. Such synergy has been described in *Vigna radiata* L. sprouts and other plants, where MeJA amplifies abiotic-induced phenylpropanoid metabolism to reinforce structural and antioxidant defenses [19,34,35]. The present findings extend this paradigm to *I. rotunda*, demonstrating that MeJA not only enhances drought tolerance but also redefines the qualitative composition of the phenolic metabolome.

From a mechanistic perspective, the observed metabolic reprogramming can be interpreted as a jasmonate-mediated reallocation of carbon resources. Under drought, carbon assimilation is constrained, and plants must prioritize metabolic pathways that maximize survival benefits per unit carbon [36,37]. By promoting the synthesis of multifunctional hydroxycinnamates, compounds that simultaneously serve as antioxidants, UV screens, and signaling mediators, MeJA optimizes resource use under stress. The suppression of less efficient phenolic branches further supports this adaptive economy. Collectively, these findings suggest that MeJA acts as a master regulator aligning metabolic flux with stress resilience [38], converting a reactive drought response into a proactive, preconditioned defense state.

Beyond physiological and mechanistic insights, the industrial implications of this metabolic modulation are significant. The enhanced accumulation of hydroxycinnamate derivatives and glycosylated phenolics under combined drought and MeJA treatment (Figures 6C, 7 and 8) positions *I. rotunda* as a promising source of high-value phytochemicals. Compounds such as verbascoside, neochlorogenic acid, and caffeoylquinic acid possess strong antioxidant, anti-inflammatory, and preservative properties that are highly sought after in nutraceutical and cosmetic industries [39,40]. The elicitation strategy demonstrated (combining controlled drought with MeJA foliar application) a sustainable, low-input approach to enrich foliage in these bioactive metabolites without genetic modification or synthetic additives. Such an approach aligns with the growing demand for eco-efficient production systems in industrial crop biotechnology.

This study provides correlative physiological and metabolomic evidence that MeJA reprograms phenolic metabolism under drought stress. However, several limitations require acknowledgment and targeted follow-up. First, causality between specific metabolites and stress tolerance remains unconfirmed; functional validation via gene silencing, chemical inhibition, or heterologous expression is needed. Second, net metabolite quantification cannot distinguish enhanced biosynthesis from reduced turnover. Integrating transcriptomics (RNA-seq, qRT-PCR of phenylpropanoid genes such as PAL, C4H, 4CL, COMT, and MYB regulators) and metabolic flux analysis is essential to elucidate the regulatory mechanisms underlying the observed metabolic reprogramming. Such transcriptomic experiments are currently underway in our laboratory. Third, molecular interrogation in I. rotunda is currently constrained by the absence of a published genome and validated reference genes; de novo transcriptome assembly and housekeeping gene validation are prerequisite steps. This limitation precludes, at present, the inclusion of gene expression data or enzyme activity measurements that would otherwise strengthen mechanistic interpretations. Consequently, enzyme names were not added to the pathway schematic (Figure 8) to avoid misrepresentation, as their roles in I. rotunda have not been experimentally validated. Fourth, while n = 3 is standard for deep metabolomic profiling, larger independent cohorts are required to validate biomarker reproducibility. Fifth, greenhouse findings require field validation; translational challenges, MeJA cost-effectiveness, formulation stability, and environmental heterogeneity, must be addressed through low-cost formulations, multi-site trials, cost–benefit analysis, and industry partnerships. Addressing these gaps, through ongoing transcriptomic studies, functional validation, and translational

research, will advance the current correlative framework toward a mechanistic and applied model for enhancing drought resilience and phenolic production in medicinal crops.

4. Materials and Methods

4.1. Plant Material and Experimental Condition

One-year-old *Ilex rotunda* Thunb. (Kurogane holly) seedlings were procured from the Winterberry Base of Runtu Horticulture Technology Co., Ltd. (Yuhang District, Hangzhou, China). The plants were transplanted into pots (9 cm in diameter × 12 cm in height) filled with a homogenized growth substrate composed of soil, peat, and sand in a 1:1:1 ratio. Subsequently, they were acclimatized for one month in a controlled greenhouse environment maintained at 22 ± 2 °C, with a 12 h light/12 h dark photoperiod and 65–70% relative humidity. Irrigation was performed every three days during this acclimation phase. After acclimation, plants of uniform height (approximately 45 cm) were selected and randomly distributed into three treatment groups (n = 6 biological replicates per group): (1) well-watered control (CK), (2) drought stress (D; water withheld for 10 days), (3) combined drought and MeJA treatment (DMeJA; water withheld and sprayed with 100 μM MeJA).

A 300 mM stock solution of Methyl Jasmonate (CAS No. 39924-52-2; Shanghai Yuanye Bio-Technology Co., Ltd., China) was prepared in ethanol. A preliminary dose–response experiment (0, 100, 200 μM MeJA) was conducted; 100 μM was selected as the optimal concentration based on maximal alleviation of wilting and phenolic accumulation. A 100 μM working solution was freshly prepared by diluting the stock in distilled water containing 0.1% (*v*/*v*) Tween-20 (CAS No. 9005-64-5; Shanghai Yuanye Bio-Technology Co., Ltd., Shanghai, China). Plants in the DMeJA group were sprayed to the point of run-off on days 0, 4, and 8 of the 10-day experimental periods. The application schedule (days 0, 4, 8) was designed to maintain bioactive MeJA levels throughout the 10-day drought period, based on reported foliar uptake in woody species [14]. Control and drought-stressed plants were sprayed with a solution containing equivalent concentrations of ethanol and Tween-20 but without MeJA. CK plants were watered every four days, whereas drought stress was imposed by completely withholding water from the D and DMeJA groups. Soil water content (SWC) was monitored daily by gravimetric method. Fresh soil samples (approx. 10 g) were collected from each pot, weighed (fresh weight), dried at 105 °C to constant weight in a forced-air drying oven (DHG-9240A, Shanghai Jinghong Experimental Equipment Co., Ltd., Shanghai, China), and reweighed (dry weight). SWC (%) was calculated as [(fresh weight − dry weight)/fresh weight] × 100. At harvest, SWC was 68.2 ± 3.1% (CK), 24.7 ± 2.4% (D), and 25.1 ± 2.6% (DMeJA), confirming severe and uniform drought stress.

After 10 days, when drought-induced symptoms such as wilting became apparent then leaf tissues were harvested from three randomly selected biological replicates per treatment. All collected samples were immediately flash-frozen in liquid nitrogen and stored at −80 °C for subsequent biochemical and metabolomic analyses.

4.2. Relative Water Content and Pigment Analysis

The relative water content (RWC) of leaf tissues was assessed using 1 cm diameter discs. Fresh discs were immediately weighed to determine the fresh weight (FW), subsequently floated on distilled water for 24 h in darkness to obtain the turgid weight (TW), and finally oven-dried at 65 °C until a constant mass was achieved to record the dry weight (DW). RWC was calculated using Equation (1):

$$\text{RWC (\%)} = [(\text{FW} - \text{DW})/(\text{TW} - \text{DW})] \times 100 \quad (1)$$

where FW: fresh weight, DW: dry wright, TW: turgit weight.

For chlorophyll quantification, approximately 0.1 g of fresh leaf tissue was finely cut with scissors and ground in a pre-chilled mortar with liquid nitrogen before extraction. Extraction was performed in 95% ethanol and incubated at 25 °C for 24 h in darkness. The extract was clarified by centrifugation at 5000× *g* for 10 min and the supernatant collected. The absorbance of the supernatant was measured at 665 nm, 649 nm, and 470 nm to determine chlorophyll *a*, *b*, and carotenoids respectively [41]. A 95% ethanol solution served as the blank. Absorbance readings were obtained using a SPARK multimode microplate reader (Model SPARK, TECAN, Switzerland; Ref. 30086376). The following equations were used:

$$\text{Chlorophyll a (mg/g FW)} = (13.95 \times A_{665} - 6.88 \times A_{649}) \times V/(1000 \times W).$$

$$\text{Chlorophyll b (mg/g FW)} = (24.96 \times A_{649} - 7.32 \times A_{665}) \times V/(1000 \times W).$$

$$\text{Total carotenoids (mg/g FW)} = [(1000 \times A_{470} - 2.05 \times \text{Chl a} - 114.8 \times \text{Chl b})/245] \times V/(1000 \times W).$$

where V = volume of extract (mL), W = fresh weight (g).

4.3. Determination of Total Phenolic Compounds and DPPH Radical Scavenging Capacity

The concentrations of total phenolic compounds and DPPH radical scavenging capacity in leaf tissues were quantified using commercial assay kits obtained from Suzhou Michy Biomedical Technology Co., Ltd. (Suzhou, Jiangsu, China). For each analysis, 0.1 g of freshly harvested leaf material was homogenized following the manufacturer's protocol for the respective kit. The total phenolic content (TP) was measured at 700 nm employing the Total Phenolic Compounds Assay Kit (Catalog No: Mo119A). The DPPH free radical scavenging activity was evaluated at 515 nm using the Total Antioxidant Capacity (T-AOC) Assay Kit (Catalog No: M0112A). Absorbance values for all assays were recorded with a SPARK multimode microplate reader (TECAN, Männedorf, Switzerland).

4.4. Determination of Oxidative Stress Markers

The levels of malondialdehyde (MDA) and hydrogen peroxide (H_2O_2) were quantified using commercial assay kits in accordance with the manufacturer's instructions. For each determination, 0.1 g of freshly collected leaf tissue was homogenized and processed following the respective assay protocols. Lipid peroxidation was assessed using the Lipid Peroxidation Assay Kit (Cat. No. M0106A) at 532 nm, while H_2O_2 concentration was measured with the Hydrogen Peroxide Assay Kit (Cat. No. M0107A) at 415 nm, both supplied by Suzhou Michy Biomedical Technology Co., Ltd. (Suzhou, Jiangsu, China). Absorbance readings were obtained using a SPARK multimode microplate reader (TECAN, Männedorf, Switzerland).

4.5. Antioxidant Defense Enzyme Activities

The activities of principal antioxidant enzymes were quantified in leaf tissues using commercial assay kits procured from Suzhou Michy Biomedical Technology Co., Ltd. (Suzhou, Jiangsu, China). For each analysis, 0.1 g of freshly harvested leaf material was homogenized following the manufacturer's specified protocols. Superoxide dismutase (SOD) activity was evaluated using the WST-8 method (Catalog No. M0102A) by monitoring the decrease in absorbance at 450 nm. Peroxidase (POD) activity was determined at 470 nm employing a Peroxidase Assay Kit (Catalog No. M0105A). Catalase (CAT) activity was measured through the ultraviolet absorption method (Catalog No. M0103A), based on the rate of hydrogen peroxide (H_2O_2) decomposition at 240 nm. All absorbance measurements

were obtained using a SPARK multimode microplate reader to ensure analytical precision and consistency across assays.

4.6. Profiling of Phenolic Compounds by UPLC–MS/MS

Metabolite profiling of *Ilex rotunda* leaf tissues was carried out using an Ultra-Performance Liquid Chromatography–Tandem Mass Spectrometry (UPLC–MS/MS) system (Wuhan Metware Biotechnology Co., Ltd., Wuhan, China). The analytical platform consisted of an ExionLC™ AD UPLC (SCIEX) coupled to a triple quadrupole-linear ion trap mass spectrometer (QTRAP® 6500+, SCIEX).

Freeze-dried tissue samples were finely ground, and 30 mg of the resulting powder was extracted with 600 µL of pre-chilled methanol:water (7:3, v/v) containing 2-chlorophenylalanine (1 µg/mL) as an internal standard for quality assurance. The mixture was vortexed thoroughly, subjected to ultrasonic extraction in an ice-water bath for 30 min, and centrifuged at 13,000× g rpm for 20 min at 4 °C. The supernatant was then filtered through a 0.22 µm filter membrane prior to analysis.

Chromatographic separation was achieved on an Agilent SB-C18 column (2.1 × 100 mm, 1.8 µm) maintained at 40 °C. The mobile phase comprised 0.1% formic acid in water (solvent A) and acetonitrile (solvent B). A linear gradient elution was applied, increasing from 5% to 95% solvent B over 9 min at a flow rate of 0.35 mL min^{-1}. Data acquisition was performed in scheduled multiple reaction monitoring (MRM) mode. The optimized ion source parameters were as follows: ion spray voltages of +5500 V (ESI^+) and –4500 V (ESI^-), source temperature 500 °C, curtain gas 25 psi, and nebulizer gases GS1 and GS2 set to 50 psi and 60 psi, respectively.

For each metabolite, precursor-to-product ion transitions, declustering potentials, and collision energies were optimized using authentic standards and the Metware Database (MWDB). Compound identification was based on the alignment of retention times, fragmentation patterns, and MS/MS spectra with those in the reference database [42]; confidence levels are provided in Supplementary Table S1. Three independent biological replicates per treatment were analyzed, which is sufficient for robust multivariate modeling (OPLS-DA) when combined with stringent significance thresholds [43,44]. Comprehensive identification parameters and raw peak areas for all detected metabolites are presented in Supplementary Table S1.

4.7. Multivariate Statistical Analyses

For the metabolomic dataset of *Ilex rotunda*, relative quantification values were normalized using unit variance scaling before statistical evaluation. Principal Component Analysis (PCA) was employed to examine the overall clustering trends among the four experimental treatments (CK, D, and DMeJA). To further discriminate between groups and identify metabolites contributing most strongly to class separation, an Orthogonal Projections to Latent Structures Discriminant Analysis (OPLS-DA) model was developed. Metabolites with a Variable Importance in Projection (VIP) score exceeding 1.0 were regarded as significant contributors [43].

Differentially accumulated metabolites (DAMs) were determined using a dual-criterion approach, combining an absolute $\log_2$ fold change ($|\log_2 FC| \geq 1.0$ with a Student's t-test p-value < 0.05, followed by false discovery rate (FDR) correction for multiple comparisons [44]. Hierarchical cluster analysis (HCA) was subsequently applied to the normalized metabolite profiles to visualize accumulation trends across all samples [45].

Duncan's multiple range test was conducted to assess statistical significance among physiological and biochemical parameters at $p < 0.05$. Venn diagrams illustrating unique and overlapping DAMs among pairwise comparisons were generated using the EVenn

online platform (https://www.bic.ac.cn/EVenn/# (accessed on 4 December 2025)) [46]. All multivariate analyses, statistical tests, and graphical visualizations were performed using R software (version 4.1.2).

5. Conclusions

This study elucidates that methyl jasmonate (MeJA) confers drought tolerance in *Ilex rotunda* through a dual mechanism: mitigating physiological impairment and strategically reprogramming phenolic metabolism. By reducing oxidative damage and enhancing non-enzymatic antioxidant capacity, MeJA shifts the defense strategy from energy-intensive enzymatic scavenging toward a more sustainable, metabolite-based system. Crucially, metabolomic and pathway analysis revealed that MeJA does not merely reverse the drought response but establishes a unique biochemical state by redirecting flux within phenylpropanoid and benzoate metabolism. This active reprogramming selectively enriches leaves in high value hydroxycinnamate derivatives like verbascoside and caffeoylquinic acids, while modulating alternative branches. These findings provide a mechanistic foundation for using MeJA as a sustainable elicitation strategy. The combined application of moderate drought and MeJA priming represents a promising, low-input biotechnological approach to enhance the resilience of *I. rotunda* cultivation and boost the targeted production of specific phenolic antioxidants for industrial use.

Supplementary Materials: The following supporting information can be downloaded at: https://www.mdpi.com/article/10.3390/plants15040648/s1, Supplementary Figure S1: Treatment-specific accumulation of selected phenolic metabolites in *Ilex rotunda* leaves. Supplementary Table S1: UPLC-MS/MS dataset of all phenolic metabolites detected in control (CK), drought-stressed (D), and MeJA-treated drought-stressed (DMeJA) *Ilex rotunda* leaves. Supplementary Table S2: Phenolic metabolites significantly altered (VIP > 1, $p < 0.05$) among all three treatment groups (CK, D, DMeJA). Supplementary Table S3: Differentially accumulated phenolic metabolites (VIP > 1, $p < 0.05$) in drought-stressed vs. control plants (D vs. CK). Supplementary Table S4: Differentially accumulated phenolic metabolites (VIP > 1, $p < 0.05$) in MeJA-treated drought-stressed vs. control plants (DMeJA vs. CK).

Author Contributions: Conceptualization, M.J.R.; methodology, M.J.R., O.C., and J.G.; software, J.G., Q.S., and L.H.; validation, J.G., H.Y., Y.Z., M.D., Y.H., M.Y., W.D., Y.L. and B.Z.; formal analysis, J.G., O.C., M.J.R., H.Y. and B.Z.; investigation, O.C., Y.Z., M.D., Y.H., M.Y., W.D., Y.L. and S.Z.; resources, Y.Z., H.Y., J.G. and B.Z.; data curation, O.C., Q.S., L.H., J.G., W.D., M.J.R. and B.Z.; writing—original draft preparation, M.J.R.; writing—review and editing, M.J.R., M.D. and B.Z.; visualization, J.G., Y.Z., M.D., Y.H., M.Y., W.D., Y.L., Q.S., L.H. and B.Z.; supervision, M.J.R., H.Y. and B.Z.; project administration, M.J.R.; funding acquisition, M.J.R. All authors have read and agreed to the published version of the manuscript.

Funding: This study was funded by the National Natural Science Foundation of China (W2533063); Ningbo Key Research and Development Project (2024Z268, 2019B10014); Central Funded Forestry Science and Technology Promotion and Demonstration Project ([2023] TS 03-1, [2020] TS 07); Open Foundation of State Key Laboratory of Subtropical Silviculture, Zhejiang A&F University (SKLSS-KF2023-07); Open Subject of Ningbo Key Laboratory of Characteristic Horticultural Crops in Quality Adjustment and Resistance Breeding (NBYYL2023004, NBYYL2023003); Ningbo Forestry Development Center Project (2023); Forestry Science and Technology Development Project of the National Forestry and Grassland Administration (KJZXSA2019034); Overseas Expertise Introduction Project for Discipline Innovation (111 Project D18008).

Institutional Review Board Statement: Not applicable.

Data Availability Statement: The original contributions presented in this study are included in the article or in the Supplementary Materials. Further inquiries can be directed to the corresponding authors.

Acknowledgments: Not applicable.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Zeng, W.; Cui, H.; Yang, W.; Zhao, Z. A Systematic Review: Botany, Phytochemistry, Traditional Uses, Pharmacology, Toxicology, Quality Control and Pharmacokinetics of Ilex Rotunda Thunb. *J. Ethnopharmacol.* **2022**, *298*, 115419. [CrossRef] [PubMed]
2. Duan, M.; Bao, L.; Jiang, T.; Song, K.; Chen, Y.; He, S.; Duan, X.; Ikram, M.; Yang, S.; Rao, M.J. Comparative LC-MS/MS Metabolomics of Wild and Cultivated Strawberries Reveals Enhanced Triterpenoid Accumulation and Superior Free Radical Scavenging Activity in *Fragaria nilgerrensis*. *Horticulturae* **2025**, *11*, 1417. [CrossRef]
3. Ngoc, P.H.; An, T.C.; Van Dung, L.; Dung, L.T.; Van Trung, P. UHPLC-ESI-QTOF-MS/MS Profiling of Saponins and Tyrosinase Inhibitory Activity of Ilex Cochinchinensis, Ilex Annamensis, and Ilex Rotunda Leaves from Vietnam. *RSC Adv.* **2025**, *15*, 43774–43791. [CrossRef] [PubMed]
4. Kim, C.-K.; Yu, J.; Le, D.; Han, S.; Yu, S.; Lee, M. Anti-Inflammatory Activity of Caffeic Acid Derivatives from Ilex Rotunda. *Int. Immunopharmacol.* **2023**, *115*, 109610. [CrossRef]
5. Cheng, G.; Yan, Y.; Zheng, B.; Yan, D. The Applications of Plant Polyphenols: Implications for the Development and Biotechnological Utilization of Ilex Species. *Plants* **2024**, *13*, 3271. [CrossRef]
6. Li, J.; Wang, Z.; Xue, L. Changes of Fluorescence Parameters of Greening Seedlings of Seven Species under Drought Stress. *BMC Plant Biol.* **2016**, *16*, 85.
7. Rao, M.J.; Zheng, B. The Role of Polyphenols in Abiotic Stress Tolerance and Their Antioxidant Properties to Scavenge Reactive Oxygen Species and Free Radicals. *Antioxidants* **2025**, *14*, 74. [CrossRef]
8. Rao, M.J.; Duan, M.; Zhou, C.; Jiao, J.; Cheng, P.; Yang, L.; Wei, W.; Shen, Q.; Ji, P.; Yang, Y.; et al. Antioxidant Defense System in Plants: Reactive Oxygen Species Production, Signaling, and Scavenging During Abiotic Stress-Induced Oxidative Damage. *Horticulturae* **2025**, *11*, 477. [CrossRef]
9. Xu, Y.; Fu, X. Reprogramming of Plant Central Metabolism in Response to Abiotic Stresses: A Metabolomics View. *Int. J. Mol. Sci.* **2022**, *23*, 5716. [CrossRef]
10. Avico, E.H.; Acevedo, R.M.; Duarte, M.J.; Rodrigues Salvador, A.; Nunes-Nesi, A.; Ruiz, O.A.; Sansberro, P.A. Integrating Transcriptional, Metabolic, and Physiological Responses to Drought Stress in Ilex Paraguariensis Roots. *Plants* **2023**, *12*, 2404. [CrossRef]
11. Acevedo, R.M.; Avico, E.H.; González, S.; Salvador, A.R.; Rivarola, M.; Paniego, N.; Nunes-Nesi, A.; Ruiz, O.A.; Sansberro, P.A. Transcript and Metabolic Adjustments Triggered by Drought in Ilex Paraguariensis Leaves. *Planta* **2019**, *250*, 445–462. [CrossRef]
12. Tienda-Parrilla, M.; López-Hidalgo, C.; Guerrero-Sanchez, V.M.; Infantes-González, Á.; Valderrama-Fernández, R.; Castillejo, M.-Á.; Jorrín-Novo, J.V.; Rey, M.-D. Untargeted MS-Based Metabolomics Analysis of the Responses to Drought Stress in *Quercus ilex* L. Leaf Seedlings and the Identification of Putative Compounds Related to Tolerance. *Forests* **2022**, *13*, 551. [CrossRef]
13. Jeyasri, R.; Muthuramalingam, P.; Karthick, K.; Shin, H.; Choi, S.H.; Ramesh, M. Methyl Jasmonate and Salicylic Acid as Powerful Elicitors for Enhancing the Production of Secondary Metabolites in Medicinal Plants: An Updated Review. *Plant Cell Tissue Organ Cult.* **2023**, *153*, 447–458. [CrossRef] [PubMed]
14. Rehman, M.; Saeed, M.S.; Fan, X.; Salam, A.; Munir, R.; Yasin, M.U.; Khan, A.R.; Muhammad, S.; Ali, B.; Ali, I. The Multifaceted Role of Jasmonic Acid in Plant Stress Mitigation: An Overview. *Plants* **2023**, *12*, 3982. [CrossRef] [PubMed]
15. Hakeem, A.; Li, S.; Nasiru, M.M.; Mustafa, G.; Elatafi, E.; Shangguan, L.; Fang, J. Methyl Jasmonate Acts as a Crucial Player in Abiotic Stress Responses in Grape. *Stresses* **2025**, *5*, 40. [CrossRef]
16. Li, C.; Xu, M.; Cai, X.; Han, Z.; Si, J.; Chen, D. Jasmonate Signaling Pathway Modulates Plant Defense, Growth, and Their Trade-Offs. *Int. J. Mol. Sci.* **2022**, *23*, 3945. [CrossRef]
17. Đurić, M.; Subotić, A.; Prokić, L.; Trifunović-Momčilov, M.; Milošević, S. Alterations in Physiological, Biochemical, and Molecular Responses of Impatiens Walleriana to Drought by Methyl Jasmonate Foliar Application. *Genes* **2023**, *14*, 1072. [CrossRef]
18. Qiu, Y.; Zhang, L.; Yang, D.; Chen, J.; Zhang, X.; Zhang, H. Effects of Exogenous Methyl Jasmonate on the Growth and Physiological Characteristics of Medicinal Chrysanthemum Morifolium Ramat under Drought Stress. *Sci. Hortic.* **2025**, *354*, 114278. [CrossRef]
19. Dong, H.; Long, H.; He, N.; Xiong, Y.; Liao, X.; Chen, T.; Guo, X. Methyl Jasmonate Synergistically Increased Polyphenol Accumulation with Blue Light in Mung Bean (*Vigna radiata* L.) Sprouts. *J. Plant Growth Regul.* **2025**, *44*, 3752–3761. [CrossRef]

20. Yan, H.; Zheng, W.; Wang, Y.; Wu, Y.; Yu, J.; Xia, P. Integrative Metabolome and Transcriptome Analysis Reveals the Regulatory Network of Flavonoid Biosynthesis in Response to MeJA in Camellia Vietnamensis Huang. *Int. J. Mol. Sci.* **2022**, *23*, 9370. [CrossRef]
21. Kim, C.-K.; Ahn, J.; Yu, J.; Le, D.; Han, S.; Lee, M. Analysis of Antioxidant Constituents from Ilex Rotunda and Evaluation of Their Blood–Brain Barrier Permeability. *Antioxidants* **2022**, *11*, 1989. [CrossRef]
22. Rao, M.J.; Duan, M.; Ikram, M.; Zheng, B. ROS Regulation and Antioxidant Responses in Plants Under Air Pollution: Molecular Signaling, Metabolic Adaptation, and Biotechnological Solutions. *Antioxidants* **2025**, *14*, 907. [CrossRef]
23. Ahmad Lone, W.; Majeed, N.; Yaqoob, U.; John, R. Exogenous Brassinosteroid and Jasmonic Acid Improve Drought Tolerance in *Brassica rapa* L. Genotypes by Modulating Osmolytes, Antioxidants and Photosynthetic System. *Plant Cell Rep.* **2022**, *41*, 603–617. [CrossRef] [PubMed]
24. Hajam, Y.A.; Lone, R.; Kumar, R. Role of Plant Phenolics against Reactive Oxygen Species (ROS) Induced Oxidative Stress and Biochemical Alterations. In *Plant Phenolics in Abiotic Stress Management*; Springer: Singapore, 2023; pp. 125–147.
25. Pérez-Llorca, M.; Pollmann, S.; Müller, M. Ethylene and Jasmonates Signaling Network Mediating Secondary Metabolites under Abiotic Stress. *Int. J. Mol. Sci.* **2023**, *24*, 5990. [CrossRef] [PubMed]
26. López-Herrador, S.; Corral-Sarasa, J.; González-García, P.; Morillas-Morota, Y.; Olivieri, E.; Jiménez-Sánchez, L.; Díaz-Casado, M.E. Natural Hydroxybenzoic and Hydroxycinnamic Acids Derivatives: Mechanisms of Action and Therapeutic Applications. *Antioxidants* **2025**, *14*, 711. [CrossRef] [PubMed]
27. Shahidi, F.; Chandrasekara, A. Hydroxycinnamates and Their in Vitro and in Vivo Antioxidant Activities. *Phytochem. Rev.* **2010**, *9*, 147–170. [CrossRef]
28. Khawula, S.; Gokul, A.; Niekerk, L.-A.; Basson, G.; Keyster, M.; Badiwe, M.; Klein, A.; Nkomo, M. Insights into the Effects of Hydroxycinnamic Acid and Its Secondary Metabolites as Antioxidants for Oxidative Stress and Plant Growth under Environmental Stresses. *Curr. Issues Mol. Biol.* **2023**, *46*, 81–95. [CrossRef]
29. Figueroa-Macías, J.P.; García, Y.C.; Núñez, M.; Díaz, K.; Olea, A.F.; Espinoza, L. Plant Growth-Defense Trade-Offs: Molecular Processes Leading to Physiological Changes. *Int. J. Mol. Sci.* **2021**, *22*, 693. [CrossRef]
30. De Geyter, N.; Gholami, A.; Goormachtig, S.; Goossens, A. Transcriptional Machineries in Jasmonate-Elicited Plant Secondary Metabolism. *Trends Plant Sci.* **2012**, *17*, 349–359. [CrossRef]
31. Yadav, V.; Wang, Z.; Wei, C.; Amo, A.; Ahmed, B.; Yang, X.; Zhang, X. Phenylpropanoid Pathway Engineering: An Emerging Approach towards Plant Defense. *Pathogens* **2020**, *9*, 312. [CrossRef]
32. Shindo, K.; Iwamoto, N.; Usami, M.; Saito, A.; Sato, M.; Sugaya, M.; Miyashita, N.; Murahama, M.; Higashimura, Y.; Takemura, M. 3′-Caffeoylquercetin Glycosides and 4′-Caffeoylkaempferol Glycosides—Novel Antioxidant Flavonoids Discovered in the Freesia Yellow Flowers. *Antioxidants* **2025**, *14*, 158. [CrossRef]
33. Boulebd, H.; Carmena-Bargueño, M.; Pérez-Sánchez, H. Exploring the Antioxidant Properties of Caffeoylquinic and Feruloylquinic Acids: A Computational Study on Hydroperoxyl Radical Scavenging and Xanthine Oxidase Inhibition. *Antioxidants* **2023**, *12*, 1669. [CrossRef] [PubMed]
34. Nie, R.; Wu, C.; Ji, X.; Li, A.; Zheng, X.; Tang, J.; Sun, L.; Su, Y.; Zhang, J. Methyl Jasmonate Orchestrates Multi-Pathway Antioxidant Defense to Enhance Salt Stress Tolerance in Walnut (*Juglans regia* L.). *Antioxidants* **2025**, *14*, 974. [CrossRef] [PubMed]
35. Yin, Q.; Feng, Z.; Ren, Z.; Wang, H.; Wu, D.; Jaisi, A.; Yang, M. Integrative Physiological, Metabolomic and Transcriptomic Insights into Phenylpropanoids Pathway Responses in Nicotiana Tabacum under Drought Stress. *Plant Stress* **2025**, *16*, 100815. [CrossRef]
36. Qin, K.; Ye, X.; Luo, S.; Fernie, A.R.; Zhang, Y. Engineering Carbon Assimilation in Plants. *J. Integr. Plant Biol.* **2025**, *67*, 926–948. [CrossRef]
37. Stefaniak, E.Z.; Tissue, D.T.; Dewar, R.C.; Medlyn, B.E. Optimal Carbon Storage during Drought. *Tree Physiol.* **2024**, *44*, 34–45. [CrossRef]
38. Li, S.; Cheng, Z.; Li, Z.; Dong, S.; Yu, X.; Zhao, P.; Liao, W.; Yu, X.; Peng, M. MeSPL9 Attenuates Drought Resistance by Regulating JA Signaling and Protectant Metabolite Contents in Cassava. *Theor. Appl. Genet.* **2022**, *135*, 817–832. [CrossRef]
39. Neelam; Khatkar, A.; Sharma, K.K. Phenylpropanoids and Its Derivatives: Biological Activities and Its Role in Food, Pharmaceutical and Cosmetic Industries. *Crit. Rev. Food Sci. Nutr.* **2020**, *60*, 2655–2675. [CrossRef]
40. Vertuani, S.; Beghelli, E.; Scalambra, E.; Malisardi, G.; Copetti, S.; Toso, R.D.; Baldisserotto, A.; Manfredini, S. Activity and Stability Studies of Verbascoside, a Novel Antioxidant, in Dermo-Cosmetic and Pharmaceutical Topical Formulations. *Molecules* **2011**, *16*, 7068–7080. [CrossRef]
41. Ptak, A.; Simlat, M.; Morańska, E.; Skrzypek, E.; Warchoł, M.; Tarakemeh, A.; Laurain-Mattar, D. Exogenous Melatonin Stimulated Amaryllidaceae Alkaloid Biosynthesis in in Vitro Cultures of *Leucojum aestivum* L. *Ind. Crops Prod.* **2019**, *138*, 111458. [CrossRef]
42. Chen, W.; Gong, L.; Guo, Z.; Wang, W.; Zhang, H.; Liu, X.; Yu, S.; Xiong, L.; Luo, J. A Novel Integrated Method for Large-Scale Detection, Identification, and Quantification of Widely Targeted Metabolites: Application in the Study of Rice Metabolomics. *Mol. Plant* **2013**, *6*, 1769–1780. [CrossRef]

43. Thévenot, E.A.; Roux, A.; Xu, Y.; Ezan, E.; Junot, C. Analysis of the Human Adult Urinary Metabolome Variations with Age, Body Mass Index, and Gender by Implementing a Comprehensive Workflow for Univariate and OPLS Statistical Analyses. *J. Proteome Res.* **2015**, *14*, 3322–3335. [CrossRef]
44. Colquhoun, D. An Investigation of the False Discovery Rate and the Misinterpretation of P-Values. *R. Soc. Open Sci.* **2014**, *1*, 140216. [CrossRef]
45. Chanana, S.; Thomas, C.S.; Zhang, F.; Rajski, S.R.; Bugni, T.S. HCAPCA: Automated Hierarchical Clustering and Principal Component Analysis of Large Metabolomic Datasets in R. *Metabolites* **2020**, *10*, 297. [CrossRef]
46. Yang, M.; Chen, T.; Liu, Y.; Huang, L. Visualizing Set Relationships: EVenn's Comprehensive Approach to Venn Diagrams. *Imeta* **2024**, *3*, e184. [CrossRef]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.