

Table S2. Mean \pm SE (sample size n inside parentheses) of final germination, and of estimated or derived parameters of Weibull equations, lag, rate, duration and shape of germination of *Cistus ladanifer* seeds incubated under constant temperature (20 °C) and photoperiod (8 h) after heat treatments (15 min.).

Temperature of heat treatment (°C)	Final germination (%)	Lag of germination (day)	Rate of germination (day)	Duration of germination (day)	Shape of germination
10	31.7 \pm 5.0 (4)	5.0 \pm 0.03 (3)	5.8 \pm 0.41 (3)	16.8 \pm 5.2 (3)	2.9 \pm 0.8 (3)
20	27.3 \pm 2.5 (8)	4.7 \pm 0.99 (5)	4.8 \pm 0.55 (5)	15.3 \pm 3.3 (5)	3.5 \pm 1.2 (5)
30	32.6 \pm 1.2 (4)	4.0 \pm 0.99 (3)	5.0 \pm 0.01 (3)	8.1 \pm 0.7 (3)	5.4 \pm 0.8 (3)
40	41.8 \pm 4.4 (8)	2.8 \pm 0.55 (4)	6.3 \pm 0.76 (4)	14.2 \pm 2.8 (4)	3.6 \pm 0.8 (4)
50	34.5 \pm 4.5 (4)	3.9 \pm 0.07 (2)	4.9 \pm 0.07 (2)	10.8 \pm 0.5 (2)	3.1 \pm 0.1 (2)
60	46.3 \pm 8.4 (8)	4.0 \pm 0.53 (6)	6.0 \pm 0.49 (6)	14.4 \pm 2.2 (6)	4.2 \pm 1.3 (6)
70	48.4 \pm 6.9 (8)	3.1 \pm 0.59 (7)	7.9 \pm 0.98 (7)	16.9 \pm 2.4 (7)	3.5 \pm 0.4 (7)
80	89.0 \pm 5.5 (4)	2.3 \pm 0.20 (4)	10.8 \pm 1.76 (4)	21.0 \pm 5.3 (4)	4.2 \pm 0.7 (4)
90	100.0 \pm 0 (4)	2.0 \pm 0.03 (4)	7.0 \pm 0.55 (4)	9.6 \pm 0.2 (4)	8.4 \pm 1.7 (4)
100	85.8 \pm 7.5 (4)	6.5 \pm 0.87 (3)	6.4 \pm 1.93 (3)	18.9 \pm 2.8 (3)	2.2 \pm 0.4 (3)
110	50.9 \pm 9.0 (8)	5.5 \pm 0.89 (7)	6.1 \pm 0.79 (7)	12.9 \pm 1.8 (7)	4.1 \pm 0.8 (7)
120	26.3 \pm 9.8 (8)	9.2 \pm 0.91 (3)	4.8 \pm 1.95 (3)	12.4 \pm 2.9 (3)	2.8 \pm 1.1 (3)
130	15.5 \pm 10.2 (8)	9.0 \pm 1.02 (2)	5.0 \pm 1.00 (2)	13.1 \pm 0.9 (2)	3.5 \pm 0.9 (2)
140	0 \pm 0 (4)	–	–	–	–
150	0 \pm 0 (4)	–	–	–	–

Differences in sample sizes between final germination and Weibull parameters result from replicates where no germination occurred or from replicates where fitting Weibull equations was not possible.