Review

Environmental Impact Assessment of Agricultural Production Using LCA: A Review

Rahmah Alhashim ¹, Raveendranpillai Deepa ² and Aavudai Anandhi ^{3,*}

¹ School of the Environment, Florida Agricultural and Mechanical University, Tallahassee, FL 32307, USA; rahmah1.alhashim@famu.edu

² College of Agriculture and Food Sciences, Florida Agricultural and Mechanical University, Tallahassee, FL 32307, USA; deepa1.raveendranpil@famu.edu

³ Biological Systems Engineering, Florida Agricultural and Mechanical University, Tallahassee, FL 32307, USA

* Correspondence: anandhi@famu.edu; Tel.: +1-850-412-5000

Abstract: Life cycle impact assessment (LCA) provides a better understanding of the energy, water, and material input and evaluates any production system's output impacts. LCA has been carried out on various crops and products across the world. Some countries, however, have none or only a few studies. Here, we present the results of a literature review, following the PRISMA protocol, of what has been done in LCA to help stakeholders in these regions to understand the environmental impact at different stages of a product. The published literature was examined using the Google Scholar database to synthesize LCA research on agricultural activities, and 74 studies were analyzed. The evaluated papers are extensively studied in order to comprehend the various impact categories involved in LCA. The study reveals that tomatoes and wheat were the major crops considered in LCA. The major environmental impacts, namely, human toxicity potential and terrestrial ecotoxicity potential, were the major focus. Furthermore, the most used impact methods were CML, ISO, and IPCC. It was also found that studies were most often conducted in the European sector since most models and databases are suited for European agri-food products. The literature review did not focus on a specific region or a crop. Consequently, many studies appeared while searching using the keywords. Notwithstanding such limitations, this review provides a valuable reference point for those practicing LCA.

Keywords: meta-analysis; GHG emission; ecotoxicity; agriculture; crop production; LCA

Citation: Alhashim, R.; Deepa, R.; Anandhi, A. Environmental Impact Assessment of Agricultural Production Using LCA: A Review. *Climate* **2021**, *9*, 164. <https://doi.org/10.3390/cli9110164>

Academic Editor: Steven McNulty

Received: 30 August 2021

Accepted: 29 October 2021

Published: 5 November 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Food supply chains (FSCs) are very complex. There are many components involved in FSCs that process, produce, package, store, transfer, distribute, and market food products to final consumers [1]. Each element in the FSC process is essential, as in any other supply chain; a change in one component affects the others. The relationship between the food system and the economy, environment, and society is mentioned by some organizations and agencies, such as the Food and Agricultural Organization (FAO), Institute of Medicine (IOM), and National Research Council (NRC), when they define the FSC [2]. Therefore, the most crucial question is as follows: Which food production system is more sustainable for the environment and communities?

There are many concerns about food resources and massive population growth, such as meeting the food demand for the world's population, production, and food consumption [1]. The total crop production must double or increase by at least 70% to meet the increasing world population's demand by 2050 [3]. Models have estimated that a 2.4% annual increase in crop yield is necessary to reach the 2050 demand [4]. The rise in food demand results in substantial energy and resource use by the food supply chain, leading to different environmental impacts. Many organizations have mentioned environmental impacts associated with food production, including the use of land, water, and climate

change. Significant environmental challenges that humans face are primarily due to climate change and the predicted future shortage of fossil fuels [5]. Farming methods, fertilizers, pesticides, water pumping, tractors to prepare the land, and transport of the crops or final food products via railroads, trucks, airplanes, or ships can all impact the environment. Lastly, food processing and food preservation methods such as refrigeration and packaging also contribute to environmental damages. There are many production sectors involved in environmental impacts, and one of them is the agricultural sector.

According to the Environmental Protection Agency (EPA), agricultural chemicals and pesticide manufacturing are two of the 68 area source groups that account for 90% of the overall emissions of the 30 urban air toxins. For example, in 2018, greenhouse gas (GHG) emissions from the agriculture economic sector accounted for 9.9% of total US greenhouse gas emissions. Furthermore, GHG from agriculture has increased by 10.1% since 1990 [6]. One of the direct greenhouse gases is nitrous oxide. Agricultural soil management operations such as synthetic and organic fertilizers and other cropping techniques, the management of manure, and the burning of agricultural wastes produce nitrous oxide. Agricultural soil management is the major source of N_2O emissions in the US, accounting for around 75% of total emissions [7]. Agricultural soils, for example, are a major source of NO_x pollution in California, with soil NO_x emissions in the state's Central Valley region being particularly high. Therefore, it is necessary to quantify the impacts of agricultural products along the food supply chain for sustainable production and consumption systems.

Since the number of operations in the food system is large and complex, many studies have used the life cycle assessment (LCA) methodology as a tool to study the overall resources used and the environmental impact of food products over its entire life cycle [8]. It is best known for its qualitative and quantitative analysis of a product's environmental aspects over its whole life cycle [9]. Products in this context include both goods and services [10]. Environmental impacts in the LCA context refer to the adverse effects on the areas of concern such as the ecosystem, human health, and natural resources. Due to the limitation of raw materials and energy resources, LCA has been used since the 1960s to find solutions for sustainable productions [11].

Such research on the crop supply chain provides helpful information from the economic, social, and environmental perspectives. Using the LCA offers a better understanding of the energy, water, and material input and evaluates the outputs' impacts. Thus, decision-makers in various fields can regulate new policies and use modern practices to improve the production supply chains. As observed in previous studies [9,12], many authors have used LCA to address environmental impacts over the entire life cycle of crops. However, the world's largest industrial sector, the food supply chain, involves various crops and products that still need to be addressed by the LCA.

Therefore, this study's broad objective is to synthesize the LCA studies relating to different environmental impacts from agricultural production to support stakeholders with decision-making. Besides, an in-depth analysis of the various steps involved in LCA is provided.

2. Materials and Methods

A literature review of published articles in international journals was undertaken using the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) protocol to address the research aims.

2.1. Eligibility Criteria

The studies that applied the following selection criteria were chosen to reduce the number of articles: (i) using the LCA method, (ii) including GHG in their impact category and/or ecotoxicity, and (iii) researching agriculture products. A total of 36 research articles were eliminated because they were about FSCs and not GHG/ecotoxicity as an effect cate-

gory, did not apply the LCA methodology, or utilized the LCA method for nonagricultural products. The LCA studies were analyzed extensively considering four phases of the LCA:

- Goal and scope definition,
- Life cycle inventory,
- Life cycle impact assessment,
- Life cycle interpretation/recommendation options.

2.2. Search Strategy

The literature review was done through the Google Scholar database. The keyword “LCA crop production” was used in the initial step, which yielded 59,100 studies as of July 2021. Later, more specific keywords were used, such as “agri-food supply chain and LCA” and “agri-food supply chain and GHG” combined with different fruit and vegetable products such as corn, peanuts, wheat, tomato, and apple. Nevertheless, the number of studies available remained enormous, the largest number of articles we got when we used the above key word with different crops was 7330, while the smallest number was 1820. A total of 110 articles were downloaded and analyzed. Twenty-nine studies were excluded because they were about FSCs and not about GHG/ecotoxicity as an effect category, or because they utilized the LCA method.

Furthermore, seven more were excluded because they used the LCA method for non-agricultural products. Accordingly, we ended up with 74 articles after applying the selection criteria. Figure 1 shows the steps used throughout the review and the inclusion criteria for the literature.

Figure 1. Steps followed for review and the inclusion/exclusion criteria.

2.3. Categorization

The data obtained from the reviewed articles included the year of study, the aim of the study, and the different steps involved in LCA assessment, which are discussed in the results section. The timeline, different components, the approach of the LCA, application of the LCA concept in the impact analysis, and suggestions for a sustainable food system are all covered.

2.4. Data Analysis

The analysis was carried out by obtaining the necessary information from the literature, as given in Tables A1 and A2 (Appendix A). Then, the information was visualized by means of collapsible trees, bar charts, doughnut figures, and word clouds after the information was classified into different result sections. Word clouds have evolved as a straightforward and visually appealing technique of text representation. They are used in a variety of contexts to offer an overview by reducing text down to the most frequently occurring terms. This is usually done statistically as a pure text summary [13,14]. Word clouds can be the initial step to refine the important concepts of results, which could save a great deal of time for other researchers since they already know where to start and the most common terms and ideas [15]. Pie and doughnut charts represent the relationship of parts with the whole [16,17]. Collapsible trees, bar charts, and doughnut figures are designed to provide greater numerical detail. Combining word clouds and bar charts allowed presenting both qualitative and quantitative information on LCA results.

The collapsible tree diagram was created with R software Version 3.6.1, and the bar and doughnut figures were created with Microsoft Excel. When making word cloud figures using the word cloud online website (<https://www.jasondavies.com/wordcloud/> accessed on November 2021), each word must be typed correctly since the size and the color of the words in the figure are affected by the number of words entered. Therefore, it is essential to make sure that the number of entered words is accurate.

Lastly, the study was organized in IMRAD format, which is the most common format for scientific papers. The term represents the first letters of the words introduction, materials and methods, results, and discussion. IMRAD format facilitates knowledge acquisition and enables easy evaluation of an article [18]. Currently, IMRAD is used by the majority of academic publications. Before the IMRAD structure, all academic writing followed the IBC (introduction, body, and conclusion) pattern. The IMRAD format is only a more specified variant of the IBC format. [19]. It is important to keep in mind that no one journal follows a standard or consistent format. Each journal has its structure, yet they all have a guideline for authors [20].

3. Results

3.1. Snapshot of Selected Studies

The characteristics of publications during 1998–2021 are displayed in Figure 2 to obtain an overview of LCA research. The number of publications per year has increased steadily since 2008, following development of the ISO standard.

Critiques of the ISO 14040 series pre-2006 were that LCA is too nascent [21], and ISO 14040 does not address uncertainty, weighting, valuation, and allocation [22].

The release of the latest version of the ISO 14040 standard in 2006 explains why LCA research is attracting more attention. Moreover, some have recently gone so far as to state that the ISO 14040: 2006 series “has proven a suitable tool for sustainability assessment” [13,14]. Fava et al. (2009) claimed that ISO 14040 should be the basis for future LCA studies [23].

Figure 2. Frequency of studies related to LCA of agricultural production from 1998 to 2021 ($n = 74$).

Studies found that the most common tool to study the impact on the environment associated with a product over its life cycle in the agri-food sector was the LCA ISO 14040 standard [14,15]. LCA ISO 14040 has four main phases: (1) goal and scope, which is the essential component of the LCA, (2) qualitative and/or quantitative inventory analysis of the used resources and the emissions released from the life cycle of a product, (3) life cycle impact assessment, which can be divided into classification, characterization, and evaluation, and (4) the interpretation, involving the identification of key issues, evaluation (including checking completeness, sensitivity, and consistency), and development of conclusions together with recommendations, as defined by ISO 14043 (Figure 3). The details of each phase are discussed below.

Figure 3. Overview of life cycle assessment (LCA) phases.

3.2. Phase 1: Goal and Scope Definition

3.2.1. Goal

According to Lee and Inaba (2004), the following questions should be addressed to set up the goal: Why perform LCA, who is the target audience, and what is the product under the LCA study [10]? These were recognized from the reviewed articles while examining the first phase of the LCA, as given in Figure 4. Some of the studies stated the answers to these questions directly, whereas others addressed them indirectly. Figures 5–7 show the most common responses to each question.

Figure 4. Phase 1 (goal and scope definition) of life cycle assessment (LCA).

Figure 5. Quantitative and qualitative representation of the common aims of LCA from the literature.

Figure 6. Target audiences in the literature represented as bars (quantitative) and a word cloud (qualitative).

Aims of LCA

As indicated in the literature, LCA studies can be partitioned into two major categories: descriptive and comparative. Descriptions aim to recognize the natural load of a chosen framework, while comparisons aim to differentiate between two frameworks. Among the discussed papers, 48 were descriptive, while 30 were comparative. As noted, the most common aim was to assess agricultural production, cultivation, processing, packaging, transport, and emission at all production stages to recognize the vast issues and to propose reasonable alternatives that decrease the environmental effects (Figure 5). The purpose of this review was to better understand how to use LCA to evaluate the environmental impact of agricultural production. The least common goal was to compare LCA to other methods, which may be due to the difficulty of making a fair comparison in terms of method performance.

Target Audience

The target audience defines who undertakes or commissions an LCA and for whom. It is critical to understand who will use the LCA results to provide them with helpful information. The majority of articles have multiple target audiences (TAs). Politicians working on climate change, decision-makers, and policymakers on global warming potential (GWP) footprints related to food and common agricultural policy (CAP) were the most common TAs, with 10 studies. Additionally, several studies targeted government sectors such as food sector policymakers, the country's agriculture sector, and the fruit and vegetable sector. Following that, the producers, namely, the farmers and the producing industry, were targeted in eight studies, six of which provided information to the consumer on a local and international scale (see Figure 6). People working on social and economic development, such as government policymakers for sustainable consumption and production, future ecolabeling programs, and those working to improve the environmental and financial sustainability of existing agricultural systems, were also targeted. Another target audience was represented by the Florida food, agri-food, and citrus industries. As shown in Figure 6, only 35 of the 74 research articles analyzed clearly stated their target audience. The frequency of target audiences is also displayed as a word cloud for a rapid overview.

Figure 7. Common agricultural products used in LCA studies.

Agricultural

We divided the products into 11 categories: tomato, fruits, citrus, vegetable, fresh salad, grains, seeds, oil, sugar, flower, and trees, as shown in Figure 7. The most common product was tomato; 13 studies analyzed tomato production, including fresh tomato, canned tomato (whole peeled, paste, and diced), and ketchup. The second most common product was wheat with nine studies. Because some studies involved more than one crop, that explains why the same reference was used for multiple crop groups and why the number of studies on the chart exceeds the number of studies covered. Tomato production was separated into three categories since three types of tomato products (fresh tomato, canned tomato, and tomato ketchup) were considered, as indicated in the diagram.

3.2.2. Scope

The scope defines the product system boundaries that determine which unit processes should be included in the LCA analysis and which should be excluded. Table A2 (Appendix A) includes more information on all 74 studies, including their inputs and outputs inside and outside of the scope. Most studies (14) contained three to four phases in their boundaries, as shown in Figure 8A. There are two explanations for not including the eliminated phases in the majority of articles. The first is a lack of data and knowledge about individual inputs, making it difficult to get a decent overall view. Secondly, some authors excluded the minor influence stages because it was impossible to include all phases.

Figure 8. (A) Number of stages included in the systems obtained from the literature; (B) quantitative and qualitative illustration of the starting and ending points of the included stages as boundaries.

Since we are looking at the agri-food supply chain, most of the articles noticeably had similar steps when designing their boundaries. Depending on the selected crop and the target audience, there were slight differences in the scope's starting point and

3.2.4. Data Quality Requirement

The reliability of the results from LCA studies strongly depends on how data quality requirements are met. The following parameters should be considered: time-related coverage (selected year), geographical coverage (study area), and technology coverage (technology used in the processes stages). This paper examined the temporal and spatial data in detail and the used machinery in general.

It is understood from the literature review that most studies collected their data for a single year of cultivation (Figure 10B). The spatial scale of the analysis (global or regional) depends on the impact category. For example, global warming is a worldwide issue, whereas acidification is a regional issue. Furthermore, two countries were commonly represented in the evaluated research, Italy and the United States, with 17 and 14 studies, respectively (Figure 10A). When it comes to the technology used in each activity, the majority of the tools mentioned were agricultural equipment, which is to be expected given that we are investigating crop production.

Figure 10. (A) Quantitative (bars) and qualitative (word cloud) representation of the geographic coverage considered in the reviewed studies; (B) donut chart depicting the temporal scales used in the literature.

3.3. Phase 2: Life Cycle Inventory

The second step of the LCA is the life cycle inventory analysis (LCI). The product's life cycle inventory results in an LCA study are obtained by summing up all fractional contributions of the input and output from each unit process in the product's production system. Thus, LCI generates quantitative environmental information of a product throughout its entire life cycle.

Most studies at this stage specified the input material (water, fertilizer, pesticide, diesel, etc.) in each process of the production included in the scope, as well as the output (harvested crop, waste, emission to the air, soil, and water, etc.). Furthermore, they mentioned the sources of the inventory data (Figure 11), typically being from primary and/or secondary data sources. Primary data are obtained from specific processes throughout the life cycle of the researched product. Process activity data (physical measures of a process that results in GHG emissions or removal), direct emissions data (determined through direct monitoring, stoichiometry, mass balance, or similar methods) from a specific site, or data averaged across all sites containing the specific process are all examples of primary data [25]. Secondary data are collected from government departments, organizational records, and studies that previously gathered information from primary sources and made it available to other researchers.

Figure 11. Phase 2 (inventory) of life cycle assessment (LCA).

About 48% of the studies used secondary data, 13% used primary data, and 35% used both. One study collected data from a real farm experience. Three authors conducted interviews with owners to collect the data. Two studies used surveys with specific questions to collect the required information. One study mentioned that the source was primary, but the article did not specify their method. Seven studies utilized primary data, while the other nine used secondary data. The authors of the examined research utilized two types of secondary data methods: databases and previous studies. Eleven of the studies used databases, while five of them used previous studies. Five writers, on the other hand, gathered inventory data from databases and prior studies. Twenty-six studies utilized both primary and secondary approaches to reduce the uncertainty of their findings (Figure 12A,B).

Figure 12. (A) Data sources of the inventory stage rendered as a pie chart; (B) breakdown of primary and secondary data into various sources as obtained from the studies.

3.4. Phase 3: Life Cycle Impact Assessment

In life cycle impact assessment (LCIA), the significance of a product system's potential environmental impacts, based on life cycle inventory results, is evaluated using LCIA. The LCIA consists of several elements: classification, characterization, normalization, and weighting. Of these four elements, normalization and weighting are considered optional, while the first two are mandatory elements in LCIA [10] (Figure 13). As shown in Figure 14, all 74 reviewed studies completed the classification and characterization phases, whereas 14 studies completed normalization and 10 completed weighting. Few studies included the waiting stage since it is optional and challenging.

The first step is classification, which involves identifying the impact assessment method. The most common standard method was the CML with various versions, such as CML 2 baseline 2000 V2/world, developed by the Center for Environmental Studies, and CML 2000 produced by the Center of Environmental Science of Leiden University. The second most common methods were ISO 14044 (2006), ISO (2000), and ISO 14040, followed by many other methods, such as IPCC 2001 GWP 100, proposed by the Intergovernmental Panel on Climate Change. For more information about the methods used in the studies, see Figure 15. The model used to calculate the impact is determined by the impact category the author intends to examine. As a result, LCA, ISO, and IPCC were the most commonly used impact methods since they provide categorization factors for ecotoxicity and climate change, which were among the criteria used to select articles for this review.

Figure 13. Phase 3 (impact assessment) of life cycle assessment (LCA).

Figure 14. Quantitative and qualitative representation of the frequency of components of the LCIA phase in the reviewed studies.

Figure 15. LCIA methods obtained from the literature review denoted by means of bar plots and a word cloud (classification).

Choosing the correct method for the LCA's impact assessment stage depends on the impact category under investigation. Each method has categories; for example, CML 2000 has 10 environmental impact categories: abiotic depletion, global warming, ozone layer depletion, human toxicity, freshwater aquatic ecotoxicity, marine aquatic ecotoxicity, terrestrial ecotoxicity, photochemical oxidation, acidification, and eutrophication.

In the process to quantify the impact of a procedure or material used, impact categories are first chosen, followed by quantifying environmental impact in each impact category using the equivalency approach. This process is termed "characterization" [10]. Characterization includes the emissions to air, soil, and water, as represented in Figure 16. The most prevalent impact categories in the 74 papers were human toxicity and ecotoxicity, with 48 and 41 studies, respectively. Moreover, 34 studies included global warming potential as an effect category, whereas marine pollution (26 articles), freshwater aquatic ecotoxicity

(23 articles), and acidification potential (22 articles) were topics of the remaining studies (Figure 16).

Figure 16. Illustration of LCIA impact categories from the literature (characterization).

3.5. Phase 4: Life Cycle Interpretation/Recommendation Options

The primary purpose of interpretation, which is the last phase of the LCA, is to use the inventory results and impact assessment analysis to evaluate the starting point for product improvement. The starting point is to understand the process tree and then identify the key issues, i.e., the key processes, materials, activities, components, or even

life cycle stages in developing a product. The primary purpose is followed up with improvement recommendations to find more environmentally friendly designs and/or process modification. Studies applied dominance analysis and marginal analysis to identify the key issues. The dominant aspects of the inventory table may be revealed by studying the environmental elements of a process matrix. An arbitrarily chosen criterion, such as “contribution greater than 1% of the total impact”, can be applied in identifying key issues from the matrix. Marginal analysis illustrates the changes in the process to which the intervention, effect, or index is most sensitive. In theory, marginal analysis is a powerful tool in determining product improvement options [8,26].

Many studies stated that, for a complete understanding of the significant driver of the impacts, it is necessary to include all stages and material used through a product’s life cycle, which is very challenging due to a lack of information and databases. However, depending on the aim of the LCA research, the literature review revealed a number of critical concerns, such as emissions from chemical and energy usage, the cultivation method used, land-use problems, and consumption waste.

Furthermore, studies in the literature proposed several recommendations for improving the agri-food system and reducing environmental consequences. One of them was adhering to the EPA and USDA pesticide and fertilizer guidelines. A frequent proposal was to use agricultural waste as animal feed. The most common request, however, was to enhance production without increasing inputs (Figure 17).

Figure 17. Phase 4 (interpretation/recommendation) of life cycle assessment (LCA).

4. Discussion

The present study reviewed articles related to the environmental impacts of agricultural production in LCA assessment. The main steps in conducting an LCA are defining the purpose of the study and boundary stages involved in the analysis, collecting the data of the inventory phase, estimating the impact of the involved process and used material, and then identifying the key issues, followed up with improvement recommendations. Most studies followed these steps, and some of them had common impact categories. However, implementing LCA is challenging and necessitates meticulous data collection.

4.1. Choice of Time, Spatial Domain, and Elementary Flows in LCA

Nearly 17% of studies did not mention the temporal scale of their analyses, depicting the inherent limitation of ISO 14040/ISO 14044 in considering the time period of evolution and process variations pertaining to diverse impact categories. The highest temporal resolution obtained from the literature was seasonal (4% of studies). The choice of time

in LCA depends on the spatial and temporal scale of the impact categories considered. For example, the temporal scale of ecotoxicity varies from hours to years. On the other hand, ecotoxicity impacts have multiple transport pathways such as air, water, and soil emissions with diverse temporal scales. Establishing a time frame for the evaluation in LCA is challenging, as both very lengthy and very short periods of assessment are not practicable depending on the topic of the LCA. Extremely short timescales violate the concept of intergenerational equality, whereas extremely long ones marginalize short-term actions, lowering the incentive to act [27]. Consequently, care should be taken when defining the temporal scale of inventory flows.

About half of the studies (49%) used secondary data collection for the LCA, acquiring data from websites and previous studies. The studies that constituted primary datasets were fewer due to the trouble of obtaining data at the desired spatial/temporal resolution for the inventory flows. The selection of impact categories and spatial domains (Figure 16) clearly reflects a preference for secondary datasets. The major categories studied were human toxicity potential and terrestrial ecotoxicity (the primary contributor being agricultural pesticide emissions). Studies used the approximated characterization factor from models for a particular spatial and temporal horizon to assess the potential impacts. Multimedia chemical exposure models such as CalTOX [28], USES-LCA [28,29], IMPACT 2002 [30], and USEtox [31] can provide the time-dependent concentrations of a chemical in the environmental compartments of air, soil, water, plants, and sediments. The potential impacts are characterized on the basis of the chemical's fate in an environmental partition and its effect.

4.2. Impact Assessment

The quantity of the input material at each stage of the crop production chain can reduce GHG, as well as emissions, including energy use (diesel, fuel, electricity) both on farm (crop production, machinery use) and off farm (transportation, refrigeration). Additional emissions include fertilizer production and use (N, P₂O₅, K₂O), pesticide use (fungicide, herbicide, insecticide), raw material production and transportation, packaging production, and disposal (Table A2). These sources of emissions contribute to environmental impacts in various ways, including human toxicity, terrestrial toxicity, freshwater toxicity, aquatic toxicity, global warming, and acidification (Figure 16). It has been demonstrated that low-input crops have minimal impacts, but high-input crops have high impacts [32]. Furthermore, the type of input can affect the rate of the impacts. For example, replacing Thomas slag with triple superphosphate reduced the toxicity associated with the presence of heavy metals [33]. Simultaneously, replacing urea with ammonium nitrate reduced the influence of fertilization on eutrophication and acidity induced by ammonia volatilization [34].

4.3. LCA as a Tool in Environmental Policy Decisions

In order to achieve the population demand in the future, increasing food production is not the only pathway to increase food availability. Increased food production necessitates either more land or increased fertilizer and pesticide use on current arable land, with negative environmental consequences such as elevated GHG emissions, biodiversity loss, water contamination, and soil erosion [35]. That explains why, among the LCA papers, the most common target audiences were policymakers and producers, whereby policymakers regulate new policies for upcoming issues and producers follow these rules. The LCA methodology can be used to identify parameters and their variability in order to assist producers, wholesale and retail consumers, and policymakers in aligning their practices and purchasing decisions with low-carbon goals. LCA can also be used to analyze different production systems in order to quantify differences in input consumption and environmental consequences. The key parameters and their variability are then addressed to offer stakeholder metrics for evaluating and aligning their agricultural processes, purchasing decisions, and policies to optimize production supply chains.

4.4. Challenges in Collecting the Information and Limitations

Obtaining each LCA component from the reviewed studies is not simple for the reader due to the authors' descriptive and nonexhaustive approach. Section 3 shows that diverse communities can benefit from this study on a local, international, and global scale. Hence, the author could have used a table or a flow chart to present the flow of components and stages to summarize the four phases and their components to enable the reader to focus on helpful information.

Another challenge is to identify what information needs to be included in the phases of the LCA. One of the essential characteristics of phase one of the LCA is using a functional unit; some authors mentioned it in the goal section while others mentioned it in the scope section. Noticeably, studies with an economic purpose often did not clearly report the functional unit.

The necessity of incorporating all production processes and their input materials, analyzing all phases to understand the environmental effect, and obtaining an optimal outcome from the LCA analysis of food production systems was emphasized by researchers. However, that is neither possible nor practical because of data limitations and cost restrictions [10]. Accordingly, the minor influential stages were excluded. Hence, most studies focused on a single phase of the food production chain. For example, some studies focused on the cultivation phase because they considered that the food production system's environmental impact mainly comes from farming activities.

The literature review did not focus on a specific region or a crop. Consequently, many studies appeared while searching using the keywords. Therefore, we included 74 articles related to LCA in agricultural production in general, as well as GHG emissions and ecotoxicity as an LCA impact category.

4.5. Assumptions Used, Benefits, and Recommendations

The LCA of crops along a food supply chain can provide helpful information from an economic, social, and environmental perspective. Using the LCA, stakeholders can better understand the energy, water, and material input and evaluate the outputs' environmental impacts. Thus, they can regulate new policies and use modern practices to improve the production supply chains.

A substantial understanding of each phase of the LCA is required to present an accurate food product's environmental impact. This paper clearly explains the LCA's major components that can serve as a primer for the scientific community. Specifically, because LCA is a systematic tool that allows for analyzing a product throughout its life cycle, LCA is used to study the economic value and importance from the local and global perspectives.

If the final product's functional unit is introduced at either the goal or the scope stage, the study results would be unaffected from our perspective. However, we recommend illustrating the input's measurement unit and the outputs while illustrating the production scope, followed by a table of units to be more readable for the audience to understand at which stage the inputs are being used and to represent the elementary flows. Defining the system boundary determines the impact pathway for an impact category that links the elementary flows from inventory to the endpoint of analysis. It is clear that the system boundary processes need to be defined according to the study's goal and the impact category. Furthermore, the functional unit must be clearly defined to explain the elementary flows from inventory to the endpoint. It is essential to know the impact category that the LCA aims to estimate, which processes are related to it, and their cause-effect relationships. The impact assessment studies were mostly conducted in the European sector since most models and databases are suited for European agri-food products.

4.6. Research Gaps

The information obtained from the literature sheds light on some of the future research needs: (a) the impact of land use on GHG emissions [36], (b) LCA applications based on irrigation techniques using solar energy dealing with waste streams [37], (c) LCA of processed and homegrown vegetables [38], (d) packaging of foods with eco-design

solutions [8], and (e) applications of LCA in organic agricultural practices, fertilization practices, mulching and milling techniques, and achievable production yields [39]. Some studies have called for more LCA applications in non-European and non-OECD countries to make their agri-food sector more environmentally friendly [40]. Therefore, it is understood that LCA can be used to make the agri-food supply chain more sustainable.

The inventory flows obtained from the present review point to the inter-dependency of three sectors in LCA: energy, food, and water. Consequently, policymakers can use LCA as a tool to spot the crucial areas that need improvisation within the framework of the food–energy–water nexus. Moreover, it is imperative to understand the drivers of environmental policy for selecting an environmentally friendly agri-food supply system. The regional variation of this nexus calls for more regional LCA assessments based on the allocation of resources. More research is needed to explore future scenarios [41] that drive resource consumption and policy design for long-term sustainability utilizing the LCA framework.

Funding: This material was based on work partially supported by the USDA-National Institute of Food and Agriculture’s Evans-Allen Project, grant 11979180/2016-01711 and 1890 Institution Capacity Building grants 2017-38821-26405, the National Science Foundation under grant No. 1735235, awarded as part of the National Science Foundation Research Traineeship as well as the Saudi Arabian Cultural Mission (SACM) under grant No. KSA10009393.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Data sharing does not apply to this article as no new data were created or analyzed in this study.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A

Table A1. Common aims in the selected studies.

Aim	Type of Aim	Studies
Evaluate the impact of all or most stages of FSC	Descriptive	[12,38,42–69]
Determine environmental differences of different cultivation options	Comparative	[24,36,39,70–77]
Estimate the impact of energy consumption	Descriptive	[42,43,48,50,78–80]
Investigate the impact of different fertilization rates and type	Comparative	[34,70,81–84]
Quantify impacts associated during cultivation cycle of a crop using life cycle analysis	Descriptive	[32,85–88]
Investigate the impact of different pesticide rates and type	Comparative	[28,89–92]
Evaluate the suitability of LCA	Descriptive	[34,93,94]
Compare the energy and GHG of regional and national scale	Comparative	[40,95,96]
Compare the impact of the cultivation of different crops	Comparative	[97–99]
Provide datasets on several agricultural products	Descriptive	[47,100]
Compare two different methods	Comparative	[101–103]

Table A2. Inventory data of the selected studies.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
1	[42]	Primary data Not mentioned how	Field preparation Seeding Post seeding weed control Creation of irrigation ditches Irrigation Supporting with reeds Fertilization Plant protection Harvest Life cycle inventory data (per 1 t of beans produced) and (per 1 ha cultivated)	Diesel Seeds Manure Water (electricity) Herbicides, insecticides, fungicides N fertilizer, P ₂ O ₅ , K ₂ O Manure cattle, sheep Seaweeds Land occupation	60 kw tractor 60 kw tractor kg kg ton ton m ² /year	Emissions to air, water, and soil) harvested beans	kg
2	[43]	Primary data (real farm) Secondary data (previous studies)	Cultivation and crop Orange transport Selection and washing Primary extraction	Fertilizers: N, P ₂ O ₅ , K ₂ O Water Diesel HDPE bins Electric energy Water Recycled water	kg MJ kg MJ kg MJ kg	CO ₂ , CO, NO _x , SO ₂ , N ₂ O, NH ₃ Oranges Wastes (leaves, rejected, citrus) Wastewater purification plant Scraps	kg kg kg kg kg kg

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
3	[44]	Primary data (Interview) secondary data (Databases)	Crop management practices				
			Maintenance of watering canals				
			Bank management				
			Plowing				
			Fertilizing				
			Harrowing				
			Sowing				
			Application of plant protection products				
			Harvesting				
			Fertilizers				
			Cuoio torrefatto (12% N);				
			ORVET 8 (8% N);	Excavation hydraulic digger			
			Urea (46% N);	Ploughing	m ³	Direct field emissions (CH ₄ , NH ₃ , etc.)	
			Calce Fosfopotassica (8% P ₂ O ₅ –22% K ₂ O–20% CaO);	Tillage, plowing	ha		
			Complezzo (18% N–36% K ₂ O);	Fertilizing, by broadcaster	ha	Indirect emissions from combustion delivered refined rice	kg
			ORVET (10% N–5% P ₂ O ₅ –15% K ₂ O);	Tillage, harrowing, by rotary harrow	ha		
			Complezzo (11% N–12% P ₂ O ₅ –36% K ₂ O)	Sowing	ha	Rice byproducts:	
			Pesticides	Application of plant protection products, by field sprayer	ha	husk, flour, broken grains, green grains	
			Gulliver	Combine harvesting	kg/ha		
			Londax 60 DF–Square 60 WDG	12% N, 46% N, 21% P ₂ O ₅ , 50% K ₂ O			
			Pull 52 DF				
			Sunrice				
			Karmex				
			Buggy–Clinic 360				
			Stratos ultra				
			Aura				
			K-Othrine				
			Dipterex				
			Heteran				
			Nominee				
			Rifit				
			Cannicid–Poladan				

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
4	[104]	Secondary data (previous studies)	Fertilizer production (process gas and fuel) Arable farming P fertilizer application Fertilizer production (effluents) Arable farming (volatilization) Fertilizer production (nitric acid production) Arable farming (denitrification/nitrification) Arable farming (leaching) P fertilizer production (effluents)	NA	NA	Fossil fuels (oil, natural gas, hard coal, lignite) Minerals (phosphate rock, potash) Land Cd CH ₄ , CO ₂ , CO, NO _x , particles, SO ₂ , NMVOC N _{tot} NH ₃ N ₂ O NO ₃ -N P _{tot}	NA
5	[95]	Secondary data (databases)	Field production Diced tomato processing Tomato paste processing Diced tomato packaging Tomato paste consumer packaging Transport: long-haul truck, rail	Fertilizers (synthetic/organic) Crop protection (chemical/organic) Energy (diesel, gas, electricity) Seeds/plants Water Energy Chemicals Packaging materials Fuel use efficiency	L/mt km	Field emissions of N ₂ O during tomato production Field emissions of CO ₂ GHG emissions associated with the production of seeds and transplants Emissions intensity	kg CO ₂ /mt km
6	[78]	Primary (survey and interview) Secondary data (databases and previous studies)	Pesticides Fertilizers Machinery Energy water	NA	NA	Emission from direct energy consumption and field emission Harvested apple	1 ton
7	[8]	Primary data (real farm) Secondary data (databases)	NA	Steel, aluminum, concrete, glass fiber resin, plastic Water Fertilizer, manure Pesticide Packaging Diesel	kg m ³ kg kg kg kg	Organic waste Construction waste Packaging Plastics oils Hazardous waste	kg kg kg kg kg kg

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
8	[45]	Primary data (interview) Secondary data (databases)	Motion of tractors Conveying and unloading Optical selection Washing Peeling Crushing and pulping for the juice Sorting Can filling and pasteurization Water purification Palletizing Irrigation Tomato fertilization Plant protection Tomato fruit transport Packaging	Diesel Electricity Natural gas Water N, P ₂ O ₅ , K ₂ O Insecticide, fungicide Tin can, label, carton tray, plastic film, pallet, box for transport, plastic boxe	kg kWh/can kWh/can m ³ /Can kg L g/can	The resulting impact was provided as output.	NA
9	[46]	Primary data (Surveys)	Resources Raw materials and fossil fuels Electric and thermal energy	Occupation, permanent crop, fruit, extensive Transformation, to permanent crop, fruit, extensive Transformation, from pasture and meadow Water, process, unspecified natural origin Fertilizer N, P ₂ O ₅ , K ₂ O Pesticides Planting Irrigating Pesticide treatments Transport Power saw Petrol unleaded at a refinery Diesel at refinery Lubricating oil Sawmill Transport, lorry 16–32 ton, EURO Orchard end of life	ha·year ha ha m ³ ton ton ha m ³ ha kton·km P kg kg kg P ton·km P	Emission in water Nitrogen, total Phosphorus, total Potassium Waste treatments Disposal, hazardous waste, 25% water, to hazardous waste incineration	ton ton ton kg

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
10	[47]	Primary data (interview) Secondary data (databases)	Fuels, fertilizers, pesticides, water use, agricultural machinery models and use, yield, harvest schedule, distance and means of transport to the packing facility.	NA	NA	Air emission Water and soil waste	NA
11	[48]	Primary data (Survey) Secondary data (databases)	Life cycle inventory data for greenhouse tomato and cucumber (per 1 ton of produced crop). Energy coefficients of different inputs and output used	Machinery Labor Diesel fuel Electricity Natural gas Nitrogen Phosphate Potassium Sul Farmyard manure Pesticides Water for irrigation Plastic 1. Machinery Tractor, self-propelled Stationary Equipment implemented, machinery 2. Human labor 3. Natural gas 4. Diesel fuel 5. Biocide Herbicide, fungicide, insecticide 6. Fertilizers: N, P ₂ O ₅ , K ₂ O 7. Micro (M) 8. Farmyard: manure 9. Water for Irrigation 10. Electricity 11. Seeds	kg h L kWh m ³ kg kg kg kg kg kg m ³ kg kg-year kg-year h m ³ L kg kg kg m ³ kWh kg	Tomato/cucumber	kg

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
12	[70]	Primary data (real farm) Secondary data (databases)	1- Preliminary considerations Doses of fertilizing products applied				
			2- Stage of compost production (CP) Collection and transport of the organic waste Industrial composting process Biofilter characteristics and gaseous emissions	Fertilizer application Compost HNO ₃ , KNO ₃ , KPO ₄ H ₂ , K ₂ SO ₄	g·m ⁻² g·m ⁻² L·m ⁻²		
			3- Stage of mineral fertilizer production (FP)	Nitrogen application organic, mineral	g·m ⁻² L·m ⁻²		
			4- Stage of compost transport	Irrigation water	m ³ ·FU ⁻¹		
			5- Stage of mineral fertilizers transport (FT)	Per area	t·ha ⁻¹		
			6- Stage of cultivation (Cu)	Per ton tomato	mm		
			Fertigation infrastructure substage (CuF)	Open field (OF)	g		
			Phytosanitary substances substage (CuP)	Commercial yield, Total yield	t·ha ⁻¹		
			Machinery and tools substage (CuM)	Tomato average diameter	t·ha ⁻¹		
			Irrigation substage (CuI)	Tomato average weight	mm		
			Post-application emissions sub-stage (CuE)	Greenhouse (GH)	g		
			Nursery plants substage (CuN)	Commercial yield	T MAL		
			Management of waste generated in the cultivation stage	Total yield diameter			
				Tomato average weight			
				Trucks			
			7- Greenhouse (G) Greenhouse structure substage (GS) Greenhouse management substage (GM) Avoided burdens of dumping OFMSW and BA in landfill				

Table A2. *Cont.*[illegible]

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
15	[96]	Secondary data (databases)	Fertilizer production Pesticide production Production of greenhouse infrastructure	Mineral fertilizer N Mineral fertilizer P Mineral fertilizer K Manure compost Organic fertilizer Steel Aluminum Glass Plexiglas Plastic Iron Concrete Rockwool	N kg·ha ⁻¹ ·year ⁻¹ P kg·ha ⁻¹ ·year ⁻¹ K kg·ha ⁻¹ ·year ⁻¹ N kg·ha ⁻¹ ·year ⁻¹ N kg·ha ⁻¹ ·year ⁻¹ kg·ha ⁻¹ ·year ⁻¹ kg·ha ⁻¹ ·year ⁻¹ kg·ha ⁻¹ ·year ⁻¹ kg·ha ⁻¹ ·year ⁻¹ kg·ha ⁻¹ ·year ⁻¹ kg·ha ⁻¹ ·year ⁻¹ kg·ha ⁻¹ ·year ⁻¹ kg·ha ⁻¹ ·year ⁻¹ kg·ha ⁻¹ ·year ⁻¹	Machine use Energy demand heating changes in soil organic carbon	h ha ⁻¹ GJ year ⁻¹ N ₂ O emissions direct N ₂ O emissions indirect Humus sequestration
16	[81]	Primary data (real farm)	NA	N min in the soil in spring Mineral N fertilizer rate Atmospheric N deposition Net N mineralization during vegetation Mineralization of N from sugar beet leaves (easily degradable part) Mineralization of N from sugar beet leaves (slowly degradable part)	NA	NH ₃ volatilization N ₂ O emission N removal with beets N content of leaves N uptake of winter wheat in autumn	One ton of grain
17	[49]	Primary data (interview) Secondary data (databases)	Greenhouse Training system Irrigation system	Low-density Polyethylene Sawn timber Steel Wire Polyethylene Sawn timber Wire Polyethylene Polyvinylchloride	k m ³ kg kg kg m ³ kg kg kg kg	Fresh tomato Air emissions NH ₃ N ₂ O-N NO _x -N Water emissions N-NO ₃	t kg·ha ⁻¹ kg·ha ⁻¹

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
18	[50]	Primary data (real farm) Secondary data (previous studies and databases)	Cultivation and crop Primary process (citrus selection and washing, extraction) Secondary process (refining; centrifugation) Secondary process (refining; pasteurization and cooling) Concentration and cooling Packaging and storage Transport of final products	Fertilizers Water Diesel Electric energy Water Recycled water Water-oil emulsion Electric energy Cooling water Raw juice Methane Electric energy Steam Electric energy Methane Steam Cooling water electric energy Essential oil Electric energy Natural juice Concentrated juice HFO, Diesel	NA	Air emissions Amount of citrus fruit Wastes (scraps, leaves, rejected citrus) Wastewater to a purification plant Scraps to pressing process Essential oil to packaging and storage Wet wastes Wastewater to purification plant Natural and concentrated juice Concentrated juice	NA
19	[40]	Secondary data (previous studies)	(larvae/fingerlings, fertilizers, and feeds).	NA	NA	nitrogen and phosphorus emissions	NA

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
20	[51]	Primary data (reports) Secondary data (databases)				Cane products	
						Cane	
						Agr. Wastes	
					ha/year	Emissions	t/day
					kg/ha-year	N ₂ O	t/day
					kg/ha-year	N total to water	kg/day
					kg/ha-year	Pesticides to water Pesticides to soil	kg/day
			Land use		kg/ha-year	Sugar	kg/day
			Pesticides		kg/ha-year	Molasses	kg/day
			Fertilizer use	Diuron, Glyphosate, Gesapox 80,	kg/ha-year	Electr. to network	kg/day
			Fuel use	MSMA 72, Amine Salt, Isoctilic ester	kg/day	Alcohol	t/day
			Seed use	48, Asulox 40, Goxone, Amigan 65,	kg/ha-year	Biogas	GJ/day
			Sun use	Merlin 75, Sulfatante 90, Unspecified	GJ/day	Ash (P ₂ O ₅ equiv.)	t/day
			Agr. operations	Urea, P ₂ O ₅ , K ₂ O	ha/year	Ash (K ₂ O equiv.)	t/day
			Lime hydrated	Diesel	ha/year	Sludge/wastewater/cake (urea equiv.)	t/day
			Cane	Cane seed	ha/year	Sludge/wastewater/cake (P ₂ O ₅ equiv.)	t/day
			Cane transport	Solar energy	ha/year	Sludge/wastewater/cake (K ₂ O equiv.)	t/day
			River water	Harvesting	t/day	Emissions to air	t/day
			Air	Fertilizing	t/day	PM ₁₀	t/day
			Softened water	Planting	t/day	Nitrogen oxides	t/day
			Ammonium sulfate	Irrigating	t/day	Emissions to water	t/day
			Sulfuric acid	NaOH 50% in H ₂ O	km	Wastewater	t/day
			Yeast	HCl 30% in H ₂ O	t/day	Inorganic solids	t/day
			Transport of filter cake		t/day	Total nitrogen	t/day
			Transport of ashes		t/day	Chemical oxygen demand	t/day
					km	Total phosphorus	t/day
						Emissions to soil	
						Ashes	
						Filter cake	
21	[52]	Primary data (interview) Secondary data (databases)	Seed production and transport	NA	NA	Emission to air and water	NA
			Fertilizer protection and transport			Solid emission	
			Pesticide production and transport				
			Machinery protection and maintenance				
			Energy carriers and protection				

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
22	[53]	Secondary data (databases)	Cultivation: Plastic cover Greenhouse Transportation: small truck, truck, sea, pre-cooling, and storage	fuel consumption, refrigeration, driving	L/t km kWh/m ³ /year	Waste management (CO ₂ emission, t/t) Paper, board, plastics CO ₂ emission from packaging, transportation, and storage Transportation Farm to packing house Packinghouse to wholesale	kg/t kg/t kg/t·km kg/t
23	[54]	Secondary data (databases)	Cattle manure Fuel use for various types of driving machinery and for different loads Low power Medium power High power Combine Willow harvester	N, P ₂ O ₅ , K ₂ O fertilizer Slurry Power	mg/kg mg/kg kw	Willow Straw Wheat	mg/kg mg/kg mg/kg
24	[82]	Secondary data (both)	Yields for main products Straw yields and crop residues Moisture content Quantity of seed Use of machinery (number of passes) Sowing and harvest date Quantity of fertilizers Types of fertilizers in integrated systems Types of fertilizers in organic systems Pesticide applications Chemical seed dressing Machinery classes Tractor harvester Trailer machinery, tillage Slurry tank	Steel, unalloyed Steel, alloyed Other metals Rubber Plastics Others (glass, paints, etc.)	NA	Ammonia emissions Nitrate leaching P-emissions N ₂ O emissions Heavy-metal emissions Pesticide applications Tractor combustion emissions	NA

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
25	[97]	Secondary data (both)		Fertilizers and lime			
				Nitrogen fertilizer (urea and diammonium phosphate)			
				Phosphate fertilizer (diammonium phosphate)			
				Potassium fertilizer (potassium chloride)	kg N	Ammonia (NH ₃)	kg/xkg
			Inventory of agricultural inputs	Agricultural lime (calcic carbonate)	kg P ₂ O ₅	Nitrates (NO ₃)	kg/xkg
			Agrochemical types and application rates	Pesticides: Clopyralid, Haloxyfop, Picloram, Glyphosate, Linuron, Thiophanate-methyl, Prochloraz	kg K ₂ O kg CaCO ₃ kg	Nitrous oxide (N ₂ O)	kg/xkg
			Seeding rate	Seed	kg	Nitrogen oxides (NO _x)	kg/xkg
			Irrigation water intake	Seed for sowing	m ³	Phosphates (PO ₄)	kg/xkg
			Fuel consumption in agricultural operations	Irrigation requirement	kg	Carbon dioxide (CO ₂)	kg/xkg
			Operating rate in machinery	Irrigation water intake	kg	Glyphosate (main pesticide in rapeseed)	kg/xkg
			Agricultural machinery type	Diesel consumption: plowing, harrowing, crushing	kg	Linuron (main pesticide in sunflower)	kg/xkg
			Seed yield	sowing, spraying, weeding, hilling/fertilizing harvest	t/ha		
				Tractor for field operations			
				Tools and harvester			
				Seed yield			

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
26	[55]	Secondary data (databases)	Inventory data on wheat production (1995–2011, year ^{−1}). Wheat grown in paddy fields and Wheat grown in upland fields	Production costs			
				Seed			
				Chemical fertilizers			
				Purchased manure			
				Pesticides 49858			
				Fossil fuels 14760			
				Electricity			
				Land improvement and irrigation			
				Agricultural services		Wheat straw	
				Buildings		Wheat	
				Agricultural machinery		Air-emission sources included fossil	
				Fossil fuels		fuel combustion, fertilizer	
				Heavy oil		application, and crop residue	
				Diesel oil	yen·ha ^{−1}	incorporation	
				Kerosene	L·ha ^{−1}	Emissions in fossil fuel combustion	
				Gasoline	kg·ha ^{−1}	were calculated using the CO ₂ , CH ₄ ,	
				Motor oil	kg N·ha ^{−1}	and N ₂ O emission factors	
				Premixed fuel		and the NO _x and SO _x emission	
				Calcium carbonate		factors	
				fertilizer		The CO ₂ emission factor of calcium	
				Nitrogen balance		carbonate fertilizer on a weight	
				Chemical fertilizers		the basis was 12%	
				Purchased manure			
				Atmospheric deposition			
				Wheat straw (incorporated)			
				Wheat			
				Wheat straw (total)			
				Denitrification			
				Ammonia volatilization			
				Surplus			

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
27	[105]	Primary data (real farm) Secondary data (previous studies and databases)	Farming				
			Irrigation				
			Soil management				
			Pest treatment				
			Fertilization	Water	m ³		
			Pruning	Pesticides	kg		
			Harvesting	Fertilizers	kg	Olive mill	L
			Olive oil mill	Diesel	kg	Wastewater	L
			Washing	Lubrication oil	L	Water from washing	L
			Milling	water	m ³	Virgin olive	L
			Pressing	Electric energy	kWh	Exhausted pomace	kg
			Decantation	Water	L	Pomace oil	kg
			Oil pomace mill	Electric energy	kWh		
			Pitting	Hexane	kg		
			Drying				
			Solvent extraction				
			Dysventilation and condensation				
28	[56]	Primary data (interview)		N, P, K			
				Lubricating oils			
				Seeds			
				Tomatoes			
				Sugar beets			
				Tomato paste			
				Raw sugar			
				Sugar solution		CH ₄ , N ₂ O, CO NMHC	kg per FU
			Fertilization	Vinegar		Biological oxygen demand (BOD)	kg per FU
			Pesticides	Spice emulsion		NO _x	g per FU
			Packaging	Salt		Other organic compounds	m ³ per FU
			Transportation	Tomato ketchup		Water emissions	kg soil per FU
				Packaging system for tomato paste		Soil emissions	
				Packaging system for ketchup			
				Transportation			
				Shopping			
				Household phase			
				Electricity production			
				Waste management			

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
29	[72]	Secondary data (databases)	Primary input and output flow from the case study farms during broccoli cropping Flow Inventory of retail-to-grave processes RDC Retailer Household	Occupation, arable land			
				Plants (plugs)			
				CO ₂ from air fixed in crop			
				Tractor use			
				Diesel (for field operations)			
				Steel (spare parts replacement)	m ² ·year		
				Labor (labor-intensive operations)	number		
				Diesel (for workers' transport)	kg CO ₂		
				Plastic (fleece, mulch ...)	hours		
				Pesticides (unspecified)	L		
				Fertilizers: N, P, K	kg		
				Manure/organic fertilizers	kg N, kg		
				Irrigation	P ₂ O ₅ , kg		
				Bluewater, surface water	K ₂ O		
				Bluewater, groundwater	kg		
				Infrastructure (pipes, sprinklers ...)	m ³		
				Electricity (pumps)	m ³	Crop	kg
				Input packed broccoli to RDC	kg	Soil emissions (literature)	kg CO ₂
				Diesel for transport to RDC	kWh	CO ₂ from soil	kg CH ₄
				From Spain	kg	CH ₄ from soil	kg NH ₃
				From the UK	kg	NH ₃ from soil	kg NO _x
				Electricity RDC storage	kg	NO _x from soil	kg N ₂ O
				Input packed broccoli to retailer	MJ	N ₂ O from soil	kg NO ₃
				Diesel for transport to retailer	kg	NO ₃ from soil	kg PO ₄
				Electricity retailer storage and display	MJ	PO ₄ from soil	kg C
				Solid waste from retailer to landfill	kg	Change in soil organic carbon (SOC)	
				Broccoli	kg		
				LDPE packaging	kg		
				Diesel for solid waste transport	MJ		
				Input broccoli to household	MJ		
				Petrol for transport to household	MJ		
				Diesel for transport to household	L		
				Electricity home storage	kg		
				Electricity cooking	kg		
				Natural gas cooking	kg		
				Tap water	L		
				Solid waste from household to landfill	kg		
				Broccoli			
				LDPE packaging			
				Diesel for solid waste transport			
				Cooking wastewater to WWTP			
				Cooked broccoli (input to human excretion)			

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
30	[38]	Secondary data (databases)	Data inventory for the agricultural phase Data inventory for the processing phase (data refer to FU)	Seeds	Mg	Emissions to air	
				Compost from cow and horse manure	g	Carbon dioxide	
				Fosetyl-Al	mg	Carbon monoxide	g
				[Thio]carbamate-compounds	mg	Nitrogen oxides	mg
				[Sulfonyl]urea-compounds	mg	Particulate hydrocarbons	mg
				Diesel fuel	g	Dinitrogen monoxide	mg
				Water	dm ³	Ammonia	mg
				Electricity for irrigation	kWh	Benfluralin	mg
				LDPE film (greenhouse)	mg	Fosetyl-Al	mg
				Land	m ²	Propamocarb	mg
				Salad (<i>Valerianella locusta</i>)	g	Emissions to water	mg
				Salad	g	Benfluralin	mg
				Electricity	kWh	Fosetyl-Al	mg
				Water	dm ³	Propamocarb	mg
				Sodium hypochlorite	mg	Emissions to soil	mg
				PP film	g	Benfluralin	mg
						Fosetyl-Al	mg
						Propamocarb	mg
						Salad bag (130 g)	p
						Salad scraps	g
						PP film waste	
						Wastewater	
31	[98]	Secondary data (previous studies)	NA	NA	NA	NA	NA

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
32	[73]	Primary data (interview) Secondary data (databases)	Main characteristics of the life cycle inventory of the studied conventional (Con) and organic (Org) groups of fruit tree orchards crops in Spain. Data refer to 1 ha and year unless otherwise stated	Drip irrigation			
				Surface irrigation			
				Water use	% of cases		
				Electricity	% of cases		
				Presence of cover crops	m ³		
				Machinery use	kWh		
				Fuel consumption	%		
				Mulching plastic	h		
				Mineral nitrogen	L		
				Mineral phosphorus	kg		
				Mineral potassium	kg N	Soil emissions	kg N ₂ O
				Manure	kg P ₂ O ₅	Direct nitrous oxide	kg N ₂ O
				Slurry	kg K ₂ O	Indirect nitrous oxide	kg CH ₄
				Cover crop seeds	mg	Methane	kg C
				Other organic fertilizers	mg	Carbon	
				Total carbon inputs	kg		
				Total nitrogen inputs Synthetic pesticides	kg		
				Sulfur	kg		
				Copper	kg active matter		
				Paraffin	kg		
				Natural pesticides	kg		
				Production	kg		
				Yield			

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
33	[57]	Secondary data (previous studies)	LCI to produce a single oil palm seedling	Electricity	kWh	Emissions to air	
				Diesel	L	NH ₃	
				Polybag	kg	N ₂ O	
				Water	L	NO	
				Fertilizer: N, P ₂ O ₅ , K ₂ O	kg	N ₂	
				Thiocarbamate	kg	Glyphosate	
				Pyrethroid	kg	Metsulfuron-methyl	
				Organophosphate	kg	Glufosinate ammonium	
				Dithiocarbamate	kg	Paraquat	
				Unspecified pesticide	kg	Emissions to water	kg/t FFB
				Urea/sulfonylurea	kg	NO ₃ ⁻	kg/t FFB
				Glyphosate	kg	PO ₄ ⁻³	
				Transportation Van	tkm	Glyphosate	Leached out and runoff
						Metsulfuron-methyl	g/t FFB
						Carbofuran	
						Glufosinate ammonium	
						Paraquat	
						Emissions to soil	
						Glyphosate	
						Metsulfuron-methyl	
						Carbofuran	
						Glufosinate ammonium	
						Methamidophos	
						Paraquat	
34	[58]	Primary data (real farm) Secondary data (databases)	Fertilizer doses, application emissions, and irrigation water (per ha) for lettuce and escarole crops in the open field (OF), plastic mulch (PM), plastic mulch combined with fleece system (PM F), and greenhouse (GH) systems. Characteristics of materials and electricity and diesel consumption (per ha) included in the inventory. PY polyethylene, PP polypropylene.	Fertilizer doses			
				N optimum			
				P ₂ O ₅	kg		
				K ₂ O	m ²		
				Mulch	m ²	Air emissions	
				Fleece		NH ₃ -N	kg
				Main pipe 1	m	NO ₂ -N	
				Main pipe 2	m	Water emissions	kg
				Main pipe 3	m	NO ₃ -N	m ³
				Secondary pipes	m	Irrigation water	
				Drip irrigation pipes (laterals)	kg		
				Pumps	MJ		
				Electricity (pumps)	MJ		
				Electricity (climate system)			
				Diesel (crop management)			

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
35	[59]	Primary data (interview)	Principal inputs involved in the analysis of the “Delizie di Bosco del Piemonte” production chain for raspberries and giant American blueberries	Substratum	L·ha ⁻¹	GWP (global warming potential) IPCC 100a Nonrenewable energy	kg CO ₂ eq MJ primary
			Nursery	Black PE	kg·ha ⁻¹		
			Rooting	White PE	kg·ha ⁻¹		
			Mulching	Metal supports	kg·ha ⁻¹		
			Covering	PVC piping	kg·ha ⁻¹		
			Covering	PVC tubing	kg·ha ⁻¹		
			Fertigation system	Compost mix	kg·ha ⁻¹		
			Fertigation system	Water	m ³ ·ha ⁻¹		
			Fertigation	PVC	kg·ha ⁻¹		
			Nozzles	Electrical energy	kWh·m ⁻³		
			Cold storage	Plow or cultivator	h·ha ⁻¹		
			Field	Harrow	h·ha ⁻¹		
			Soil preparation	Bed-former	h·ha ⁻¹		
			Soil preparation	Diesel consumption	L·h ⁻¹		
			Mulching	PE sheeting	kg·ha ⁻¹		
			Total processes	PVC piping	kg·ha ⁻¹		
			Mulching	PVC tubing	kg·ha ⁻¹		
			Irrigation system	Water	m ³ ·ha ⁻¹		
			Irrigation system	Electrical energy for the well	kWh·ha ⁻¹		
			Irrigation	Manure	t·ha ⁻¹		
			Irrigation	Compost	t·ha ⁻¹		
			Base fertilization	White PE	kg·ha ⁻¹		
			Total fertilization	Metal supports	kg·ha ⁻¹		
			Covering	p.a.	kg·ha ⁻¹		
			Covering	Electrical energy	kWh·kg ⁻¹		
			Plant protection treatments	Electrical energy	kWh·kg ⁻¹		
			Post-harvesting	PE tray	g·kg ⁻¹		
			Refrigeration	PE wrapping	g·kg ⁻¹		
			Flow packaging				
			Flow packaging				
			Flow packaging				

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
36	[60]	Secondary data (databases)	Rice production tillage, growing, harvest	Machines, materials		Rice field Pollution (emissions) Product, byproduct Rice field product, byproduct, pollution	
37	[36]	Primary data (survey) Secondary data (previous studies and databases)		Seed Power tiller diesel fuel use GHG intensity diesel fuel Power tiller life expectancy Power tiller weight Tractor L diesel fuel/h Tractor weight Embodied GHG of steel Bullocks Allocation to straw Tractors embodied emission Fertilizers Pesticides Manure Nitrogen use efficiency	kg CO ₂ eq·ha ⁻¹ L/h kg CO ₂ eq·L ⁻¹ Years kg L/h kg kg CO ₂ eq·kg steel ⁻¹ kg CO ₂ eq·h ⁻¹ kg CO ₂ eq·h ⁻¹ kg CO ₂ eq·kg ⁻¹ CO ₂ -eq kg/kg CO ₂ -eq·t ⁻¹	Methane emissions Nitrous oxide emissions SRI CH ₄ and N ₂ O emissions Electricity-based emissions from irrigation. Embodied GHG emissions associated with electricity Harvest Soil organic carbon	kg CO ₂ -eq·ha ⁻¹ kg CO ₂ eq·ha ⁻¹ kg CO ₂ eq·ha ⁻¹ kg CO ₂ eq·kWh ⁻¹ GHG emissions·h ⁻¹ kg CO ₂ eq·ha ⁻¹
38	[61]	Primary data (interview)	Primary production Grading and packing Regional distribution center Supermarkets	Piscicide production N fertilizer production Tools Machinery Water Compost Field diesel Packaging Electricity Electricity Pallets and packaging Electricity Pallets and packaging	NA	Land-use change Direct emission Nitrate Nitrous oxide Ammonia Waste Waste waste	NA
39	[62]	Primary data (reports) Secondary data (previous studies)	Orchard establishment inputs Agricultural stage inputs Retail stages inputs Consumption stages inputs	Water Electricity Diesel Machinery Materials Transport	L kW kg kg kg tkm	Apple Peach (NPK) NO _x N ₂ O Machinery production emissions and diesel consumed for machinery operations	kg kg

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
40	[74]	Secondary data (both)	Annual chemical inputs for managing a mature orange grove in Florida Chemical mowing Herbicide spray Pesticide spray Fertilization Use of energy products for undertaking various cultural activities at a mature orange grove in Florida Site preparation Management of a mature orange grove	Roundup weather max			
				Solcam 80 DF			
				Karmex WP			
				Roundup weather max			
				Prowl H20			
				Simazine 4L			
				Roundup weather max			
				Mandate			
				Direx 4L			
				Roundup weather max			
				Spray oil	mL/ha		
				Copper (Kocide 3000)	kg/ha		
				Agrimek (if no mite resistance)	kg/ha		
				Zn, Mn, B	mL/ha		
				Lorsban 4EC	mL/ha		
				Copper (Kocide 3000)	mL/ha		
				Spray Oil	mL/ha		
				MgO	mL/ha		
				Dolomite	mL/ha		
				Mowing (mechanical)	mL/ha	Emission from energy use	g CO ₂ eq./FU
				Mowing (chemical)	L/ha	Emission from material use	g CO ₂ eq./FU
				Discing	kg/ha		
				Soil shaping	mL/ha		
				Planting	kg/ha		
				Mowing (mechanical) Mowing (chemical)	mL/ha		
				Fertilization (16–0–16–4 MgO)	kg/ha		
				Fertilization (lime)	L/ha		
				Herbicide	kg/ha		
				Pesticide	kg/ha		
				Conditioning			
				Topping			
				Hedging			
				Brush removing			
				Chopping brush			
				Dead tree removal			
				Irrigation			
				Fruit picking			
				Transporting pickers Roadsiding fruit			

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
41	[75]	Primary data (survey) Secondary data (databases)	Principal inputs involved in the production and distribution chain (scenarios 1 and 2) for strawberries				
			Nursery	Substratum	L·ha ⁻¹		
			Rooting	Black PE	kg·ha ⁻¹		
			Mulching	White PE	kg·ha ⁻¹		
			Covering	Metal supports	kg·ha ⁻¹		
			Covering	PVC piping	kg·ha ⁻¹		
			Fertigation system	PVC tubing	kg·ha ⁻¹		
			Fertigation system	Compost mix	kg·ha ⁻¹		
			Fertigation	Water	m ³ ·ha ⁻¹		
			Cold storage	Electrical energy	kWh·m ⁻³		
			Field	Plow or cultivator	h·ha ⁻¹		
			Soil preparation	Harrow	h·ha ⁻¹		
			Soil preparation	Bed-former	h·ha ⁻¹		
			Mulching	Diesel consumption	L·h ⁻¹	GWP (global warming potential) IPCC 100a	kg CO ₂ eq·UF ⁻¹
			Total processes	PE sheeting	kg·ha ⁻¹	Non-renewable energy	kg CO ₂ eq·UF ⁻¹
			Mulching	PVC piping	kg·ha ⁻¹		
			Irrigation system	PVC tubing	kg·ha ⁻¹		
			Irrigation system	Water	m ³ ·ha ⁻¹		
			Irrigation	Electrical energy for the well	kWh·ha ⁻¹		
			Irrigation	Manure	t·ha ⁻¹		
			Base fertilization	Compost	t·ha ⁻¹		
			Total fertilization	White PE	kg·ha ⁻¹		
			Covering	Metal supports	kg·ha ⁻¹		
			Covering	p.a.	kg·ha ⁻¹		
			Plant protection treatments	Electrical energy	kWh·kg ⁻¹		
			Post-harvesting	PE tray	g·kg ⁻¹		
			Refrigeration	PE wrapping	g/kg		
			Flow packaging				
			Flow packaging				

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
42	[93]	Secondary data (databases)	Life cycle inventory data for watermelon cultivation (per ha). Characterization factors of inputs used in watermelon production. Parameters and coefficients of objective functions.	1. Human labor (man/woman)			
				2. Diesel fuel			
				Plowing			
				Discing			
				Ditcher			
				3. Machinery			
				Tractor and self-propelled			
				Implement and machinery			
				4. Fertilizers	h		
				Nitrogen (N)	L		
				Phosphate (P ₂ O ₅)	kg		
				Potassium (K ₂ O)	kg		
				Microelements	kg		
				5. Farmyard manure	kWh		
				6. Electricity	kg		
				7. Chemicals	kg		
				Fungicide	kg		
				Insecticide	kg		
				8. Seeds	L		
				9. Plastics	kg	Watermelon	kg
				Machinery	kg	On-farm emissions	kg
				Diesel fuel	kg	N fertilizer	kg
				Chemical fertilizers	kWh	Diesel fuel	MJ
				(a) Urea	kg		
				(b) Phosphate (P ₂ O ₅)	kg		
				(c) Potassium (K ₂ O)	kg		
				Manure	kg		
				Pesticides	kg		
				Electricity	L		
				Plastics	kWh		
				Constanta	kg		
				N	kg		
				K ₂ O	kg		
				P ₂ O ₅	kg		
				Manure	MJ		
				Diesel			
				Electricity			
				Seed			
				Chemicals			
				Machinery			
				Plastic			
				Water			

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
43	[76]	Secondary data (databases)	NA	Mulching film for pot production (PP) Wind-stopper (galvanized iron) Hydraulic pipe/micro pipe (PEHD/PELD/PVC) Taps (PEHD/PVC) Tunnel cover Tunnel structure (galvanized iron) Poles (galvanized iron/wood) Sprinklers (galvanized iron) Hydraulic fittings (PE) Solenoid (PVC) Support canes (bamboo) Black clip (PP) Plates PP black wire (nylon) Green thread (PVC) Iron wire (galvanized iron) Elastics/hooks/butterfly valve (PE) Irrigation bar (aluminum) Block (concrete) Covering (gravel/volcanic stones) Raincoat towel (PVC/PP/PEHD) A chain-link fence (galvanized iron) Centrifugal/submersible pump (Cast iron/stainless steel) Electrical panel (PEHD/copper) Burlap (jute) String (sisal) Wire basket (iron) Plastic net (PP) Plastic box (PP)	NA	Total yearly GHG emissions are divided into different categories (kg CO ₂ eq/m ² /year) NFS $\frac{1}{4}$ nursery farm structure; AGS $\frac{1}{4}$ aboveground structures; IC $\frac{1}{4}$ inputs of cultivation; P $\frac{1}{4}$ packaging; EFS $\frac{1}{4}$ emissions from soil	NA
44	[106]	Secondary data (databases)	NA	Strawberry (nursery field) PE punnet PE plastic film End-of-life Transport Electricity	NA	Nonrenewable energy IPCC GWP 100a	MJ·UF ^{−1} kg CO ₂ eq·UF ^{−1}

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
45	[79]	Primary data (surveys)	Gasoline at the refinery (US) Diesel at the refinery (US) Urea ammonium nitrate (UAN), (US) Monoammonium phosphate (US) Waxes/paraffin at the refinery (US) Potassium sulfate, at regional storage (Europe) Mined from natural sources, only transport is modeled Fishmeal Potassium carbonate, at the plant (Europe) Sulfur (elemental) at the refinery (US) Yeast (surrogate data, yeast produced as a co-product) Serenade is a strain of <i>Bacillus subtilis</i> (Swiss) Glyphosate, at regional storehouse (Europe) Diphenyl-ether compounds at regional storehouse (Europe) Phtalamide compounds at regional storehouse (Europe) Pesticide unspecified, at regional storehouse RER Developed based on Recycled Organics Unit (2006), updated with regionally appropriate LCI datasets Electricity grid mix (West US) Modeled based on power rating and hours of operation (California model) and Diesel (US) Truck (combination)—diesel rail (US diesel)	Gasoline Diesel Urea ammonium nitrate (UAN) Monoammonium phosphate (MAP) Adjuvant (stylet oil) Potassium sulfate Phytamin component: seabird guano Phytamin component: fishmeal Phytamin component: potassium carbonate Sulfur dust Serenade Roundup Ultra Max Goal 2XL Chateau, Pristine (Boscalid and Pyraclostrobin) Compost production Electricity Equipment operation Truck, rail shipping International shipping	NA	IPCC Tier 2 emissions were used to calculate the field-based N ₂ O emissions from fertilizer and compost application and vineyard plant matter, including leaves, clippings, and cover crop residue following mowing (Intergovernmental Panel on Climate Change 2006; Point et al. 2012).	NA

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
46	[63]	Secondary data (databases)	Fossil energy life cycle factors for agricultural inputs	Nitrogen	MJ/kg	Direct N ₂ O emissions from agricultural	NA
				Phosphorous	MJ/kg	Emissions (e.g., volatile organic compound	
				Potassium	MJ/kg	(VOC), carbon monoxide (CO),	
				Lime	MJ/kg	carbon dioxide (CO ₂), nitrogen	
				Sulfur	MJ/kg	monoxide (NO), nitrogen dioxide	
				Micronutrients	MJ/kg	(NO ₂), nitrous oxide (N ₂ O),	
				Cover crop seed	MJ/kg	particulate matter (PM10),	
				Herbicide	MJ/kg	particulate matter (PM2.5), sulfur	
				Insecticide	MJ/L	dioxide (SO ₂), sulfur trioxide (SO ₃),	
				Fungicide	MJ/L	methane (CH ₄))	
				Gasoline	MJ/kg	Emissions and energy use in	
				Diesel	MJ/h	transportation	
				Plastic	MJ/kWh		
				Agriculture machinery Electricity			
47	[24]	Secondary data (databases)	Electricity production	Electricity	MJ	CO ₂	kg
			Oil production Plastic P1				
			production				
			Gutter A1 production				
			Gutter A1 use and demolition				
			Incineration of P1/A1				
			Recycling process Material B				
			production				
			Product system			CH ₄	kg
			Electricity production			N ₂ O	kg
			Oil production Plastic P2			NO _x	kg
			production			SO ₂	kg
			Gutter A2 production				
			Gutter A2 use and demolition				
			Incineration of P2/A2				
			Recycling process Material B				
			production				
			Product system				

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
48	[64]	Secondary data (databases)	Planting and maintenance Harvesting and baling Receiving/storage Drying and chopping Pelletizing/cooling/screening Packing and storage	Seed			
				Fertilizer			
				Pesticide/herbicide	kg·ha ⁻¹	Strawbale	kg
				Land use	ha	CO ₂	g CO ₂ eq
				Machinery	kg·ha ⁻¹	N ₂ O	g CO ₂ eq
				Fuel	ha	CH ₄	g CO ₂ eq
				Machinery	MJ	SO ₂	g SO ₂ eq
				Fuel	MJ	PO ₄	g PO ₄ eq
				Electricity	kWh	Pellet	kg
				Air			
				Plastic bag			
49	[65]	Primary data (real farm) Secondary data (database and previous studies)	Production characteristics Greenhouse plastic Water consumption Growing media Fertilizer Pesticide Electric power Diesel and petrol Post-harvest chemicals	Plastic consumption	g		
				Rejected steams	#		
				Power consumption	kWh		
				Diesel	g		
				Petrol	g		
				Cardboard box	g		
				Bunching paper	g		
				Rubber band	g	Roses	Bunch
				Strapping roll	g	CO ₂	g
				Water	L	CH ₄	g
				Substrate (red ash)	g	N ₂ O	g
				Pesticide	g		
				Pesticide empty containers	g		
				Calcium nitrate	g		
				Other fertilizers	g		
				Acids	g		
				Post-harvest chemicals	g		
				Post-harvest water use	L		

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
50	[32]	Secondary data (database and previous studies)	Production of crop inputs, production and use of diesel, and field emissions	N (ammonium nitrate)	kg/ha	Hemp	ha
				P ₂ O ₅ (triple superphosphate)	kg/ha	Sunflower	ha
				K ₂ O (potassium chloride)	kg/ha	Rapeseed	ha
				CaO	kg/ha	Pea	ha
				Seed for sowing	kg/ha	Wheat	ha
				Pesticide (active ingredient)	kg/ha	Maize	ha
				Diesel	kg/ha	Potato	ha
				Natural gas (for grain drying)	kg/ha	Sugar beet	ha
				Agricultural machinery	kg/ha	NH ₃ -N	emissions/kg
				Grain dry matter yield	kg/ha	NO ₃ -N	emissions/kg
				Stem/straw dry matter yield	kg/ha	N ₂ O-N	emissions/kg
				Sugar/tuber dry matter yield	kg/ha	PO ₄ -P	emissions/kg
				Followed bycatch crop (%)	kg/ha		
				Succeeding crop NO ₃ -N emitted	kg/ha		
51	[100]	Secondary data (database and previous studies)	Infrastructure:			Potatoes organic, at the farm	
						Rapeseed extensive, at the farm	kg
						Wheat grains conventional, Barrois, at the farm	kg
						Carbon dioxide CO ₂	g/kg
						Sulfur dioxide SO ₂	g/kg
						Lead Pb	g/kg
						Methane CH ₄	g/kg
						Benzene C ₆ H ₆	g/kg
						Particulate Matter PM	g/kg
						Cadmium Cd	g/kg
						Chromium Cr	g/kg
						Copper Cu	g/kg
						Monoxide N ₂ O	g/kg
						Nickel Ni	
			Fieldwork processes:				
				Mineral fertilizers	kg		
				Organic fertilizers	kg		
				Pesticides	kg		
				Seed	kg		
				Feed	kg		
			<ul style="list-style-type: none"> • Buildings • Machinery • Soil cultivation • Fertilization • Sowing • Chemical plant protection • Mechanical treatment • Harvest • Transport 				

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
52	[85]	Primary data (survey)	Tractors and equipment Buildings required energy Carriers Mineral fertilizer Tree nursing Constructions for hail protection Water for irrigation Application of compost	Pesticides			
				Fungicide			
				Insecticide			
				Herbicide			
				Other plants			
				treatment products	kg active matter		
				Fertilizers	kg N		
				N-fertilizer	kg K ₂ O		
				Ca- and Mg-fertilizer	kg P ₂ O ₅	Total receipts	USD·ha ⁻¹
				(kg Ca, Mg)	kg	Yield	t·ha ⁻¹
				K-fertilizer	kg		
				P-fertilizer	kg		
				Machinery	m ²		
				Diesel			
				Tractor			
				Equipment			
				Buildings			
53	[89]	Secondary data (databases)	Pesticide	Energy input	MJ eq	CH ₄	t CO ₂
			Seeds				eq·ha ⁻¹ ·year ⁻¹
			PK fertilizer			N ₂ O	kg N
			N fertilizer				eq·ha ⁻¹ ·year ⁻¹
			Machinery mulching			CO ₂	t CO ₂
			Machinery irrigation				eq·ha ⁻¹ ·year ⁻¹
			Machinery pesticide			Ph	kg N
			Machinery fertilization				eq·ha ⁻¹ ·year ⁻¹
			Machinery weeding			NH ₃	kg N
			Machinery soil tillage				eq·ha ⁻¹ ·year ⁻¹
			Machinery harvest				kg N
			Machinery sowing			NO ⁻³	eq·ha ⁻¹ ·year ⁻¹

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
54	[77]	Primary data (farmers) Secondary data (databases and references)	Transportation Fertilization Pesticides Irrigation	Inputs		Grape	
				1. Diesel fuel		Ammonia (NH ₃)	kg
				2. Transportation		Ammonia (NH ₃)	kg·ha ⁻¹
				3. Human labor		Benzene	kg·ha ⁻¹
				4. Chemical fertilizers		Benzo (a) pyrene	kg·ha ⁻¹
				(a) Nitrogen	L	Cadmium (Cd)	kg·ha ⁻¹
				(b) Phosphate	kg	Carbon dioxide (CO ₂)	kg·ha ⁻¹
				(c) Potassium	h	Carbon dioxide (CO ₂) from urea.	kg·ha ⁻¹
				(d) Sulfur	kg	Carbon monoxide (CO)	kg·ha ⁻¹
				5. Manure	kg	Chromium (Cr)	kg·ha ⁻¹
				6. Chemical pesticides	m ³	Copper (Cu)	kg·ha ⁻¹
				(a) Fungicide	MJ	Diazinon	kg·ha ⁻¹
				(b) Insecticide		Dinitrogen monoxide (N ₂ O)	kg·ha ⁻¹
				7. Irrigation water		Dinitrogen monoxide (N ₂ O)	kg·ha ⁻¹
				The total energy input		Dinitrogen monoxide (N ₂ O) from atmospheric deposition	kg·ha ⁻¹
						Hydrocarbons (HC, as NMVOC)	kg·ha ⁻¹
						Methane (CH ₄)	kg·ha ⁻¹
						Nickel (Ni)	kg·ha ⁻¹
						Nitrate (NO ₃)	kg·ha ⁻¹
						Nitrogen oxide (NO _x)	kg·ha ⁻¹
						Nitrogen oxides (NO _x)	kg·ha ⁻¹
						PAH (polycyclic hydrocarbons)	kg·ha ⁻¹
						Particulates (b2.5 mm)	kg·ha ⁻¹
						Phosphorus emissions from fertilizers application emitted into groundwater.	kg·ha ⁻¹
						Selenium (Se)	kg·ha ⁻¹
						Sulfur dioxide (SO ₂)	kg·ha ⁻¹
						Tillet	
						Zinc (Zn)	

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
55	[83]	Primary data (field experiment) Secondary data (database)	Fertilization Cutting preparation Spraying Ploughing Disking Harrowing Marking Spraying Mechanical weeding Fertilizing Lignin production and application Harvest Transport Liquidation	Tractor/harvester Machinery Diesel fuel	kg·ha ⁻¹	CO ₂ PM SO ₂ p	kg Mg ⁻¹ CO ₂ eq kg MP 10 eq kg SO ₂ eq kg p eq
56	[90]	Secondary data (databases and previous studies)	Pesticide application	Active ingredients of the pesticide	t	Active ingredients emissions	Unitless
57	[66]	Primary (survey) Secondary data (database and previous studies)	Nursery Tomato cultivation Packaging Transportation	Reporting period Country (Production site) Growing period Greenhouse structure Substrate Greenhouse heating CO ₂ enrichment Yield Fertilization Irrigation Energy consumption	ton·ha ⁻¹ kg N·ha ⁻¹ , kg P ₂ O ₅ ·ha ⁻¹ , kg K ₂ O·ha ⁻¹ water m ³ ·ha ⁻¹ kWh·ha ⁻¹	Nitrogen oxides, phosphates, and pesticides emissions nitrous oxide, and ammonia	g N eq g P eq

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
58	[86]	Secondary data (databases)	Applying farmyard manure Land preparation Planting Fertilizing Harvesting	N-based fertilizers P-based fertilizers K-based fertilizers Pesticides Farmyard manure Microelements Diesel fuel Water	kg kg kg kg t kg L m ³	Carbon dioxide (CO ₂)	kg
						Sulfur dioxide (SO ₂)	
						Methane (CH ₄)	
						Benzene	
						Cadmium (Cd)	
						Chromium (Cr)	
						Copper (Cu)	
						Dinitrogen monoxide (N ₂ O)	
						Nickel (Ni)	
						Zinc (Zn)	
						Benzo(a)pyrene	
						Ammonia (NH ₃)	
						Selenium (Se)	
						PAH (polycyclic hydrocarbons)	
						Hydrocarbons (HC, as NMVOC)	
						Nitrogen oxides (NO _x)	
						Carbon monoxide (CO)	
						Particulates (<2.5 µm)	
59	[67]	Secondary data (databases)	Fertilization, split fertilization, chemical fallow, liming, sowing and spraying at the farm	NPK-fertilizer	kg·ha ⁻¹	N ₂ O NH ₃ NO _x	kg N ₂ O-N/kg N input kg NH ₃ -N NO _x -N/kg
				N-fertilizer	kg·ha ⁻¹		
				Roundup (glyphosate)	kg·ha ⁻¹		
				Dolomite (CaO)	kg·ha ⁻¹		
				Celest Formula M (fludioxonil)	kg·ha ⁻¹		
				Starane XL (fluroxypyr/florasulam)	kg·ha ⁻¹		
				Fastac 50 (alpha-ceypermethrin)	kg·ha ⁻¹		
60	[103]	Primary data (real farm) Secondary data (databases)	production, transport to the farm and use on the farm	Fertilizers, pesticides, field materials, pesticide spray equipment, irrigation system and packaging manufacturing	NA	Pesticide emission	NA

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
61	[87]	Secondary data (databases and previous studies)	<ul style="list-style-type: none">• Production of nitrogenous mineral fertilizer• Transportation of organic fertilizer• Production of phosphorus mineral fertilizer• Production of potassium mineral fertilizer• Production of lime• Production of agricultural equipment• Production of seeds for sowing/default seeds harvested crop scenario <p>Production of diesel</p>	Fertilizer N, P, K Lime Fuel Seeds Agricultural equipment	NA	Agricultural engine emissions (CO, HC, NO _x , SO ₂ , PM, CO ₂) (EPA 2004) Direct field emission from fertilization (NO ₃ , NH ₄ , N ₂ O, NO _x , CO ₂ , PO ₄) Hemp straws and seeds	NA
62	[91]	Secondary data (databases)	Pesticide application	S-Metolachlor (H) Simazine (H) Glyphosate (H) Glufosinate ammonium (H) Dimethenamid-P (H) Atrazine (H) Alachlor (H) Acetochlor (H) 2,4-D-dimethylammonium . . . 2,4-D-2-ethylhexyl ester (H) Fipronil (I) Chlorpyrifos (I)	kg/kg corn kg/kg corn kg/kg corn kg/kg corn kg/kg corn kg/kg corn kg/kg corn kg/kg corn kg/kg corn kg/kg corn	Pesticide emission to air, surface water, and groundwater	%

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
63	[101]	Primary data (real farm) Secondary data (databases)	Manufacture of greenhouse components, substrate, fertilizers, and pesticides. the electricity production mix; and transport and disposal of materials	greenhouse components Water consumption and fertilizer and pesticide doses applied	kg, m ² , m ³ M ³ kg, L kg	N ₂ O	kg
						NO _x	kg
						NH ₃	kg
						Azoxystrobin	kg
						Chlorothalonil	kg
						Clofentezine	kg
						Fenbutatin oxide	kg
						Mancozeb	kg
						Spinosad	kg
						Copper chloride oxide, hydrate	kg
						Concrete	kg
						Plastics	kg
						Substrate	kg
64	[94]	Primary data (survey) Secondary data (databases and previous studies)	<ul style="list-style-type: none"> Agricultural field operations (including plowing, harrowing, sowing, chemical weed control, harvesting, straw baling); Seeds, fertilizers, and pesticides production. Grain drying. Nitrogen and phosphate (fertilizers)' emissions; and Pesticide's emissions. 	Yield			
				Grain	t/ha		
				Straw	t/ha		
				Agricultural field operations	number of		
				Ploughing	repetitions (rep)		
				Harrowing by rotary harrow	rep		
				Sowing	rep		
				Fertilizing by broadcaster	rep		
				Slurry spreading	rep		
				Pest control application by field	rep	Fertilizers' emissions	
				Sprayer	rep	NH ₃	kg/ha
				Harvesting	rep	N ₂ O	kg/ha
				Bailing	rep	NO ₃	kg/ha
				Transport (tractor and trailer)	rep	PO ₄	kg/ha
				Grain drying	rep	Pesticides' emissions	
				Seeds	kg/ha	Tribenuron-methyl	kg/ha
				Fertilizers	kg N/ha	Pyraclostrobin	kg/ha
				Calcium ammonium nitrate (CAN)	kg N/ha	Tebuconazole	kg/ha
				Ammonium nitrate	kg N/ha	Pirimicarb	kg/ha
				Pig slurry	kg N/ha	Thifensulfuron-methyl (difensulfuron)	kg/ha
				Dairy cattle slurry	g/ha	2,4-D-Bromoxynil	
				Pesticides	g/ha		
				Tribenuron-methyl	g/ha		
				Pyraclostrobin	g/ha		
				Tebuconazole	g/ha		
				Pirimicarb	g/ha		
				Difensulfuron	g/ha		
				2,3-D-Bromoxynil			

Table A2. *Cont.*[illegible]

Table A2. *Cont.*

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
68	[99]	Primary data (survey) Secondary data (databases)	Direct agricultural inputs Production of the different agricultural inputs,	Information about tractors and implements, labor hours, and input rates such as agrochemicals and water use) nitrogen (urea and ammonium nitrate), phosphorous or potassium-based fertilizers and herbicides (terbutylazine, alachlor, lumax, and S-metolachlor		NH ₃ N ₂ O NO ₃	kg N ₂ O-N·ha ⁻¹ ·kg ⁻¹
69	[92]	Secondary data (databases and previous studies)	Pesticide application	Abamectin Azadirachtin Chlorpyrifos Clofentezine Copper oxychloride Fenazaquin Fenbutatin-oxyde Fluroxypyr Fosetyl-Al Glufosinate-ammonium Glyphosate Hexythiazox Imazalil Imidacloprid Insecticide Lambda-cyhalothrin Mancozeb MCPA Paraquat Propargite Pyridaben Pyriproxyfen Spinosad Tebufenpyrad Thiabendazole White mineral oil (paraffin oil)	kg·m ⁻² kg·m ⁻²	Pesticide emission	NA

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
70	[88]	Primary data (survey and interview) Secondary data (Databases and previous studies)	NA	Cut flowers Carnation support net Plastic cover material Putty for sun protection Water for the putty used for sun protection Transporting of cut flowers to Athens (2 times per week) Electricity consumption for refrigeration of the cut flowers, water pumping Fertilizers Water for plant protection Fungicides, Pesticides Soil disinfection (once every three years) Water for soil disinfection, plant watering (3 times per week) Humidification	Stems/year kg/year kg/year kg/year m ³ /year km/year kWh/year kg N/year, kg P/year, kg K/year m ³ /year kg/year kg/year m ³ /year	N ₂ O, NO _x , and ammonia	NA
71	[28]	Secondary data (databases and previous studies)	Pesticide application	Abamectine Azoxystrobin Benomyl Bromopropylate Captan Cyromazine Deltamethrin Fenarimol Iprodione Kresoxim-metil Mancozeb Pimetrozine	kgai·FU ⁻¹ kgai·FU ⁻¹ kgai·FU ⁻¹ kgai·FU ⁻¹ kgai·FU ⁻¹ kgai·FU ⁻¹ kgai·FU ⁻¹ kgai·FU ⁻¹ kgai·FU ⁻¹ kgai·FU ⁻¹ kgai·FU ⁻¹ kgai·FU ⁻¹	Pesticide emission to air, soil, and water	NA

Table A2. Cont.

#	Reference	Data Source	Practice	Input	Unit	Output	Unit
72	[12]	Secondary data (databases)	Fertilizer application Seeds use Plant protection production application Agriculture activity	N, P, K fertilizer	NA	NH ₃ NO _x N ₂ O NO ₃ PO ₄ ⁻³ P HM ^a Heavy metal Active ingredients CO ₂ NMVOC PM	NA
73	[102]	Secondary data (databases)	NA	Pesticide N Fertilizer P Fertilizer	NA	N ₂ O, air emission P, water Emission NO ₃ ⁻ , water Emission Pesticides, water emission	NA
74	[69]	Primary data (surveys) Secondary data (databases)	Soil tillage Seedbed preparation Owing Fertilization Plant protection Harvest Stubble cultivation Transport to the farm and grain drying	Fertilizer use Number of passes for fertilizer spreading Pesticide use (active ingredients) Herbicides Fungicides Insecticides Other pesticides Total pesticides Number of passes for pesticide spraying	kg N·ha ⁻¹ ·year ⁻¹ kg P ₂ O ₅ ·ha ⁻¹ ·year ⁻¹ kg K ₂ O·ha ⁻¹ ·year ⁻¹ ha ⁻¹ ·year ⁻¹ kg·ha ⁻¹ ·year ⁻¹ kg·ha ⁻¹ ·year ⁻¹ kg·ha ⁻¹ ·year ⁻¹ kg·ha ⁻¹ ·year ⁻¹ kg·ha ⁻¹ ·year ⁻¹ kg·ha ⁻¹ ·year ⁻¹ ha ⁻¹ ·year ⁻¹	Yields Gross energy yield Raw protein yield Gross margin	kg DM·ha ⁻¹ ·year ⁻¹ GJ·ha ⁻¹ ·year ⁻¹ kg·ha ⁻¹ ·year ⁻¹ D·ha ⁻¹ ·year ⁻¹

References

- Kucukvar, M.; Ismaen, R.; Onat, N.C.; Al-Hajri, A.; Al-Yafay, H.; Al-Darwish, A. Exploring the Social, Economic and Environmental Footprint of Food Consumption: A Supply Chain-Linked Sustainability Assessment. In Proceedings of the 2019 IEEE 6th International Conference on Industrial Engineering and Applications (ICIEA), Tokyo, Japan, 12–15 April 2019; IEEE: Piscataway, NJ, USA; pp. 733–742.
- Institute of Medicine and National Research Council. 2 Overview of the U.S. Food System. In *A Framework for Assessing Effects of the Food System*; The National Academies Press: Washington, DC, USA, 2015. Available online: <https://www.nap.edu/read/18846/chapter/5> (accessed on 28 September 2021). [CrossRef]
- Tilman, D.; Balzer, C.; Hill, J.; Befort, B.L. Global Food Demand and the Sustainable Intensification of Agriculture. *Proc. Natl Acad. Sci. USA* **2011**, *108*, 20260–20264. [CrossRef] [PubMed]
- Brandt, K.; Barrangou, R. Applications of CRISPR Technologies Across the Food Supply Chain. *Annu. Rev. Food Sci. Technol.* **2019**, *10*, 133–150. [CrossRef] [PubMed]
- FAO; WFP; IFAD. *The State of Food Insecurity in the World 2012. Economic Growth Is Necessary but Not Sufficient to Accelerate Reduction of Hunger and Malnutrition*; FAO: Rome, Italy, 2012; p. 5. Available online: <https://www.fao.org/3/i3027e/i3027e00.pdf> (accessed on 28 September 2021).
- Sources of Greenhouse Gas Emissions. US EPA. Available online: <https://www.epa.gov/ghgemissions/sources-greenhouse-gas-emissions> (accessed on 28 July 2021).
- US EPA. Overview of Greenhouse Gases. Available online: <https://www.epa.gov/ghgemissions/overview-greenhouse-gases> (accessed on 28 July 2021).
- Cellura, M.; Longo, S.; Mistretta, M. Life Cycle Assessment (LCA) of Protected Crops: An Italian Case Study. *J. Clean. Prod.* **2012**, *28*, 56–62. [CrossRef]
- Von Blottnitz, H.; Curran, M.A. A Review of Assessments Conducted on Bio-Ethanol as a Transportation Fuel from a Net Energy, Greenhouse Gas, and Environmental Life Cycle Perspective. *J. Clean. Prod.* **2007**, *15*, 607–619. [CrossRef]
- Lee, K.-M.; Inaba, A. *Life Cycle Assessment Best Practices of ISO 14040 Series*; Center for Ecodesign and LCA (CEL), Ajou University: Suwon, Korea, 2004; p. 96.
- Buyle, M.; Braet, J.; Audenaert, A. Life Cycle Assessment in the Construction Sector: A Review. *Renew. Sustain. Energy Rev.* **2013**, *26*, 379–388. [CrossRef]
- Corrado, S.; Castellani, V.; Zampori, L.; Sala, S. Systematic Analysis of Secondary Life Cycle Inventories When Modelling Agricultural Production: A Case Study for Arable Crops. *J. Clean. Prod.* **2018**, *172*, 3990–4000. [CrossRef] [PubMed]
- Atenstaedt, R. Word Cloud Analysis of the BJGP. *Br. J. Gen. Pract.* **2012**, *62*, 148. [CrossRef] [PubMed]
- Heimerl, F.; Lohmann, S.; Lange, S.; Ertl, T. Word Cloud Explorer: Text Analytics Based on Word Clouds. In Proceedings of the 2014 47th Hawaii International Conference on System Sciences, Waikoloa, HI, USA, 6–9 January 2014; pp. 1833–1842.
- DePaolo, C.A.; Wilkinson, K. Get Your Head into the Clouds: Using Word Clouds for Analyzing Qualitative Assessment Data. *TechTrends* **2014**, *58*, 38–44. [CrossRef]
- Present Your Data in A Doughnut Chart. Available online: <https://support.microsoft.com/en-us/office/present-your-data-in-a-doughnut-chart-0ac0efde-34e2-4dc6-9b7f-ac93d1783353> (accessed on 4 October 2021).
- What Is A Pie Chart and When to Use It. Available online: <https://www.storytellingwithdata.com/blog/2020/5/14/what-is-a-pie-chart> (accessed on 4 October 2021).
- Wu, J. Improving the Writing of Research Papers: IMRAD and Beyond. *Landsc. Ecol.* **2011**, *26*, 1345–1349. [CrossRef]
- Vyas, V.d.H.A. Tantraguna—The Ancient Criteria for Scientific Writing. *Ayu* **2016**, *37*, 158–162. [CrossRef] [PubMed]
- Nair, P.K.R.; Nair, V.D. Organization of a Research Paper: The IMRAD Format. In *Scientific Writing and Communication in Agriculture and Natural Resources*; Nair, P.K.R., Nair, V.D., Eds.; Springer International Publishing: Cham, Switzerland, 2014; pp. 13–25.
- Hertwich, E.G.; Pease, W.S. ISO 14042 Restricts Use and Development of Impact Assessment. *Int. J. Life Cycle Assess.* **1998**, *3*, 180–181. [CrossRef]
- Pryshlakivsky, J.; Searcy, C. Fifteen Years of ISO 14040: A Review. *J. Clean. Prod.* **2013**, *57*, 115–123. [CrossRef]
- Fava, J.; Baer, S.; Cooper, J. Increasing Demands for Life Cycle Assessments in North America. *J. Ind. Ecol.* **2009**, *13*, 491–494. [CrossRef]
- Huijbregts, M. Uncertainty and Variability in Environmental Life-Cycle Assessment. *Int. J. Life Cycle Assess.* **2002**, *7*, 173. [CrossRef]
- Bhatia, P.; Cummis, C.; Draucker, L.; Rich, D.; Lahd, H.; Brown (WBCSD), A. *Greenhouse Gas Protocol Product Life Cycle Accounting and Reporting Standard*; World Resources Institute (WRI) and World Business Council for Sustainable Development (WBCSD): Washington, DC, USA, 2011.
- Heijungs, R.; Guinée, J.B. (Eds.) *Environmental Life Cycle Assessment of Products*; Centre of Environmental Science: Leiden, The Netherlands, 1992.
- Lueddeckens, S.; Saling, P.; Guenther, E. Temporal Issues in Life Cycle Assessment—A Systematic Review. *Int. J. Life Cycle Assess.* **2020**, *25*, 1385–1401. [CrossRef]

28. Antón, A.; Castells, F.; Montero, J.I.; Huijbregts, M. Comparison of Toxicological Impacts of Integrated and Chemical Pest Management in Mediterranean Greenhouses. *Chemosphere* **2004**, *54*, 1225–1235. [[CrossRef](#)] [[PubMed](#)]
29. Huijbregts, M.A.J.; Struijs, J.; Goedkoop, M.; Heijungs, R.; Jan Hendriks, A.; van de Meent, D. Human Population Intake Fractions and Environmental Fate Factors of Toxic Pollutants in Life Cycle Impact Assessment. *Chemosphere* **2005**, *61*, 1495–1504. [[CrossRef](#)] [[PubMed](#)]
30. Jolliet, O.; Margni, M.; Charles, R.; Humbert, S.; Payet, J.; Rebitzer, G.; Rosenbaum, R. IMPACT 2002+: A New Life Cycle Impact Assessment Methodology. *Int. J. Life Cycle Assess.* **2003**, *8*, 324. [[CrossRef](#)]
31. de García, S.O.; García-Encina, P.A.; Irusta-Mata, R. The Potential Ecotoxicological Impact of Pharmaceutical and Personal Care Products on Humans and Freshwater, Based on USEtoxTM Characterization Factors. A Spanish Case Study of Toxicity Impact Scores. *Sci. Total Environ.* **2017**, *609*, 429–445. [[CrossRef](#)]
32. Van Der Werf, H.M.G. Life Cycle Analysis of Field Production of Fibre Hemp, the Effect of Production Practices on Environmental Impacts. *Euphytica* **2004**, *140*, 13–23. [[CrossRef](#)]
33. Charles, R.; Jolliet, O.; Gaillard, G.; Pellet, D. Environmental Analysis of Intensity Level in Wheat Crop Production Using Life Cycle Assessment. *Agric. Ecosyst. Environ.* **2006**, *113*, 216–225. [[CrossRef](#)]
34. Brentrup, F.; Küsters, J.; Lammel, J.; Barraclough, P.; Kuhlmann, H. Environmental Impact Assessment of Agricultural Production Systems Using the Life Cycle Assessment (LCA) Methodology II. The Application to N Fertilizer Use in Winter Wheat Production Systems. *Eur. J. Agron.* **2004**, *20*, 265–279. [[CrossRef](#)]
35. Nesheim, M.C.; Oria, M.; Yih, P.T.; Committee on a Framework for Assessing the Health, Environmental, and Social Effects of the Food System; Food and Nutrition Board; Board on Agriculture and Natural Resources; Institute of Medicine; National Research Council. *Environmental Effects of the U.S. Food System*; National Academies Press: Washington, DC, USA, 2015.
36. Gathorne-Hardy, A. A life cycle assessment (LCA) of greenhouse gas emissions from SRI and flooded rice production in SE India. *Taiwan Water Conserv. J.* **2013**, *61*, 111–125.
37. Hanieh, A.A.; Hasan, A.; Assi, M. Date Palm Trees Supply Chain and Sustainable Model. *J. Clean. Prod.* **2020**, *258*, 120951. [[CrossRef](#)]
38. Fusi, A.; Castellani, V.; Bacenetti, J.; Cocetta, G.; Fiala, M.; Guidetti, R. The Environmental Impact of the Production of Fresh Cut Salad: A Case Study in Italy. *Int. J. Life Cycle Assess.* **2016**, *21*, 162–175. [[CrossRef](#)]
39. Tasca, A.L.; Nessi, S.; Rigamonti, L. Environmental Sustainability of Agri-Food Supply Chains: An LCA Comparison between Two Alternative Forms of Production and Distribution of Endive in Northern Italy. *J. Clean. Prod.* **2017**, *140*, 725–741. [[CrossRef](#)]
40. Reckenholz-Tänikon, A. Towards a Sustainable Management of the Food Chain. In Proceedings of the Research Station ART 6th International Conference on Life Cycle Assessment in the Agri-Food Sector, Zurich, Switzerland, 12–14 November 2008.
41. O'Neill, B.C.; Kriegler, E.; Ebi, K.L.; Kemp-Benedict, E.; Riahi, K.; Rothman, D.S.; van Ruijven, B.J.; van Vuuren, D.P.; Birkmann, J.; Kok, K. The Roads Ahead: Narratives for Shared Socioeconomic Pathways Describing World Futures in the 21st Century. *Glob. Environ. Chang.* **2017**, *42*, 169–180. [[CrossRef](#)]
42. Abeliotis, K.; Detsis, V.; Pappia, C. Life Cycle Assessment of Bean Production in the Prespa National Park, Greece. *J. Clean. Prod.* **2013**, *41*, 89–96. [[CrossRef](#)]
43. Beccali, M.; Cellura, M.; Iudicello, M.; Mistretta, M. Resource Consumption and Environmental Impacts of the Agrofood Sector: Life Cycle Assessment of Italian Citrus-Based Products. *Environ. Manag.* **2009**, *43*, 707–724. [[CrossRef](#)] [[PubMed](#)]
44. Blengini, G.A.; Busto, M. The Life Cycle of Rice: LCA of Alternative Agri-Food Chain Management Systems in Vercelli (Italy). *J. Environ. Manag.* **2009**, *90*, 1512–1522. [[CrossRef](#)] [[PubMed](#)]
45. Garofalo, P.; D'Andrea, L.; Tomaiuolo, M.; Venezia, A.; Castrignanò, A. Environmental Sustainability of Agri-Food Supply Chains in Italy: The Case of the Whole-Peeled Tomato Production under Life Cycle Assessment Methodology. *J. Food Eng.* **2017**, *200*, 1–12. [[CrossRef](#)]
46. Radzimska, M.; Garbowska, B.; Jakubowska, D. Health Quality and Nutritional Value of Rye Bread Produced on a Small and Large Scale in Poland. *Ital. J. Food Sci.* **2013**, *25*, 126.
47. Ingwersen, W.W. Life Cycle Assessment of Fresh Pineapple from Costa Rica. *J. Clean. Prod.* **2012**, *35*, 152–163. [[CrossRef](#)]
48. Khoshnevisan, B.; Rafiee, S.; Omid, M.; Mousazadeh, H.; Clark, S. Environmental Impact Assessment of Tomato and Cucumber Cultivation in Greenhouses Using Life Cycle Assessment and Adaptive Neuro-Fuzzy Inference System. *J. Clean. Prod.* **2014**, *73*, 183–192. [[CrossRef](#)]
49. Bojacá, C.R.; Wyckhuys, K.A.; Schrevens, E. Life Cycle Assessment of Colombian Greenhouse Tomato Production Based on Farmer-Level Survey Data. *J. Clean. Prod.* **2014**, *69*, 26–33. [[CrossRef](#)]
50. Beccali, M.; Cellura, M.; Iudicello, M.; Mistretta, M. Life Cycle Assessment of Italian Citrus-Based Products. Sensitivity Analysis and Improvement Scenarios. *J. Environ. Manag.* **2010**, *91*, 1415–1428. [[CrossRef](#)]
51. Contreras, A.M.; Rosa, E.; Pérez, M.; Van Langenhove, H.; Dewulf, J. Comparative Life Cycle Assessment of Four Alternatives for Using By-Products of Cane Sugar Production. *J. Clean. Prod.* **2009**, *17*, 772–779. [[CrossRef](#)]
52. Hospido, A.; Milà i Canals, L.; McLaren, S.; Truninger, M.; Edwards-Jones, G.; Clift, R. The Role of Seasonality in Lettuce Consumption: A Case Study of Environmental and Social Aspects. *Int. J. Life Cycle Assess.* **2009**, *14*, 381–391. [[CrossRef](#)]
53. Roy, P.; Nei, D.; Okadome, H.; Nakamura, N.; Orikasa, T.; Shiina, T. Life Cycle Inventory Analysis of Fresh Tomato Distribution Systems in Japan Considering the Quality Aspect. *J. Food Eng.* **2008**, *86*, 225–233. [[CrossRef](#)]

54. Broek, R.; Treffers, D.-J.; Meeusen, M.; Wijk, A.; Nieuwlaar, E.; Turkenburg, W. Green Energy or Organic Food? A Life-Cycle Assessment Comparing Two Uses of Set-Aside Land. *J. Ind. Ecol.* **2001**, *5*, 65–87. [\[CrossRef\]](#)
55. Masuda, K. Measuring Eco-Efficiency of Wheat Production in Japan: A Combined Application of Life Cycle Assessment and Data Envelopment Analysis. *J. Clean. Prod.* **2016**, *126*, 373–381. [\[CrossRef\]](#)
56. Andersson, K.; Ohlsson, T.; Olsson, P. Screening Life Cycle Assessment (LCA) of Tomato Ketchup: A Case Study. *J. Clean. Prod.* **1998**, *6*, 277–288. [\[CrossRef\]](#)
57. Choo, Y.M.; Muhamad, H.; Hashim, Z.; Subramaniam, V.; Puah, C.W.; Tan, Y. Determination of GHG Contributions by Subsystems in the Oil Palm Supply Chain Using the LCA Approach. *Int. J. Life Cycle Assess.* **2011**, *16*, 669–681. [\[CrossRef\]](#)
58. Romero-Gómez, M.; Audsley, E.; Suárez-Rey, E.M. Life Cycle Assessment of Cultivating Lettuce and Escarole in Spain. *J. Clean. Prod.* **2014**, *73*, 193–203. [\[CrossRef\]](#)
59. Girgenti, V.; Peano, C.; Bounous, M.; Baudino, C. A Life Cycle Assessment of Non-Renewable Energy Use and Greenhouse Gas Emissions Associated with Blueberry and Raspberry Production in Northern Italy. *Sci. Total Environ.* **2013**, *458*, 414–418. [\[CrossRef\]](#) [\[PubMed\]](#)
60. Breiling, M.; Hoshino, T.; Matsushashi, R. *Contributions of Rice Production to Japanese Greenhouse Gas Emissions Applying Life Cycle Assessment as a Methodology*; The University of Tokyo: Tokyo, Japan, 1999; p. 32.
61. Kulak, M.; Graves, A.; Chatterton, J. Reducing Greenhouse Gas Emissions with Urban Agriculture: A Life Cycle Assessment Perspective. *Landsc. Urban Plan.* **2013**, *111*, 68–78. [\[CrossRef\]](#)
62. Vinyes, E.; Asin, L.; Alegre, S.; Muñoz, P.; Boschmonart, J.; Gasol, C.M. Life Cycle Assessment of Apple and Peach Production, Distribution and Consumption in Mediterranean Fruit Sector. *J. Clean. Prod.* **2017**, *149*, 313–320. [\[CrossRef\]](#)
63. Tabatabaie, S.M.H.; Murthy, G.S. Cradle to Farm Gate Life Cycle Assessment of Strawberry Production in the United States. *J. Clean. Prod.* **2016**, *127*, 548–554. [\[CrossRef\]](#)
64. Li, X.; Mupondwa, E.; Panigrahi, S.; Tabil, L.; Adapa, P. Life Cycle Assessment of Densified Wheat Straw Pellets in the Canadian Prairies. *Int. J. Life Cycle Assess.* **2012**, *17*, 420–431. [\[CrossRef\]](#)
65. Sahle, A.; Potting, J. Environmental Life Cycle Assessment of Ethiopian Rose Cultivation. *Sci. Total Environ.* **2013**, *443*, 163–172. [\[CrossRef\]](#)
66. Payen, S.; Basset-Mens, C.; Perret, S. LCA of Local and Imported Tomato: An Energy and Water Trade-Off. *J. Clean. Prod.* **2015**, *87*, 139–148. [\[CrossRef\]](#)
67. Roer, A.-G.; Korsæth, A.; Henriksen, T.M.; Michelsen, O.; Strømman, A.H. The Influence of System Boundaries on Life Cycle Assessment of Grain Production in Central Southeast Norway. *Agric. Syst.* **2012**, *111*, 75–84. [\[CrossRef\]](#)
68. Matsuura, M.I.S.F.; Dias, F.R.T.D.; Picoli, J.F.; Lucas, K.R.G.; de Castro, C.; Hirakuri, M.H. Life-Cycle Assessment of the Soybean-Sunflower Production System in the Brazilian Cerrado. *Int. J. Life Cycle Assess.* **2017**, *22*, 492–501. [\[CrossRef\]](#)
69. Nemecek, T.; Von Richthofen, J.-S.; Dubois, G.; Casta, P.; Charles, R.; Pahl, H. Environmental Impacts of Introducing Grain Legumes into European Crop Rotations. *Eur. J. Agron.* **2008**, *28*, 380–393. [\[CrossRef\]](#)
70. Martínez-Blanco, J.; Muñoz, P.; Antón, A.; Rieradevall, J. Assessment of Tomato Mediterranean Production in Open-Field and Standard Multi-Tunnel Greenhouse, with Compost or Mineral Fertilizers, from an Agricultural and Environmental Standpoint. *J. Clean. Prod.* **2011**, *19*, 985–997. [\[CrossRef\]](#)
71. Meisterling, K.; Samaras, C.; Schweizer, V. Decisions to Reduce Greenhouse Gases from Agriculture and Product Transport: LCA Case Study of Organic and Conventional Wheat. *J. Clean. Prod.* **2009**, *17*, 222–230. [\[CrossRef\]](#)
72. Canals, L.M.I.; Muñoz, I.; Hospido, A.; Plassmann, K.; McLaren, S.; Edwards-Jones, G.; Hounsome, B. Life cycle assessment (LCA) of domestic vs. imported vegetables. In *Case Studies on Broccoli, Salad Crops and Green Beans*; In RELU Project REW-224-25-0044; Centre for Environmental Strategy, University of Surrey: Guildford, UK, 2008.
73. Aguilera, E.; Guzmán, G.; Alonso, A. Greenhouse Gas Emissions from Conventional and Organic Cropping Systems in Spain. II. Fruit Tree Orchards. *Agron. Sustain. Dev.* **2015**, *35*, 725–737. [\[CrossRef\]](#)
74. Dwivedi, P.; Spreen, T.; Goodrich-Schneider, R. Global Warming Impact of Florida's Not-From-Concentrate (NFC) Orange Juice. *Agric. Syst.* **2012**, *108*, 104–111. [\[CrossRef\]](#)
75. Girgenti, V.; Peano, C.; Baudino, C.; Tecco, N. From “Farm to Fork” Strawberry System: Current Realities and Potential Innovative Scenarios from Life Cycle Assessment of Non-Renewable Energy Use and Green House Gas Emissions. *Sci. Total Environ.* **2014**, *473*, 48–53. [\[CrossRef\]](#) [\[PubMed\]](#)
76. Lazzerini, G.; Lucchetti, S.; Nicese, F.P. Green House Gases(GHG) Emissions from the Ornamental Plant Nursery Industry: A Life Cycle Assessment(LCA) Approach in a Nursery District in Central Italy. *J. Clean. Prod.* **2016**, *112*, 4022–4030. [\[CrossRef\]](#)
77. Mohseni, P.; Borghei, A.M.; Khanali, M. Coupled Life Cycle Assessment and Data Envelopment Analysis for Mitigation of Environmental Impacts and Enhancement of Energy Efficiency in Grape Production. *J. Clean. Prod.* **2018**, *197*, 937–947. [\[CrossRef\]](#)
78. Milà i Canals, L.; Burnip, G.M.; Cowell, S.J. Evaluation of the Environmental Impacts of Apple Production Using Life Cycle Assessment (LCA): Case Study in New Zealand. *Agric. Ecosyst. Environ.* **2006**, *114*, 226–238. [\[CrossRef\]](#)
79. Steenwerth, K.L.; Strong, E.B.; Greenhut, R.F.; Williams, L.; Kendall, A. Life Cycle Greenhouse Gas, Energy, and Water Assessment of Wine Grape Production in California. *Int. J. Life Cycle Assess.* **2015**, *20*, 1243–1253. [\[CrossRef\]](#)
80. Langevin, B.; Basset-Mens, C.; Lardon, L. Inclusion of the Variability of Diffuse Pollutions in LCA for Agriculture: The Case of Slurry Application Techniques. *J. Clean. Prod.* **2010**, *18*, 747–755. [\[CrossRef\]](#)

81. Brentrup, F.; Küsters, J.; Kuhlmann, H.; Lammel, J. Application of the Life Cycle Assessment Methodology to Agricultural Production: An Example of Sugar Beet Production with Different Forms of Nitrogen Fertilisers. *Eur. J. Agron.* **2001**, *14*, 221–233. [\[CrossRef\]](#)
82. Nemecek, T.; Erzinger, S. Modelling Representative Life Cycle Inventories for Swiss Arable Crops (9 Pp). *Int. J. Life Cycle Assess.* **2005**, *10*, 68–76. [\[CrossRef\]](#)
83. Krzyżaniak, M.; Stolarski, M.J.; Warmiński, K. Life Cycle Assessment of Poplar Production: Environmental Impact of Different Soil Enrichment Methods. *J. Clean. Prod.* **2019**, *206*, 785–796. [\[CrossRef\]](#)
84. Lebailly, F.; Levasseur, A.; Samson, R.; Deschênes, L. Development of a Dynamic LCA Approach for the Freshwater Ecotoxicity Impact of Metals and Application to a Case Study Regarding Zinc Fertilization. *Int. J. Life Cycle Assess.* **2014**, *19*, 1745–1754. [\[CrossRef\]](#)
85. Mouron, P.; Nemecek, T.; Scholz, R.W.; Weber, O. Management Influence on Environmental Impacts in an Apple Production System on Swiss Fruit Farms: Combining Life Cycle Assessment with Statistical Risk Assessment. *Agric. Ecosyst. Environ.* **2006**, *114*, 311–322. [\[CrossRef\]](#)
86. Khanali, M.; Shahvarooghi Farahani, S.; Shojaei, H.; Elhami, B. Life Cycle Environmental Impacts of Saffron Production in Iran. *Environ. Sci. Pollut. Res. Int.* **2017**, *24*, 4812–4821. [\[CrossRef\]](#) [\[PubMed\]](#)
87. Andrianandraina; Ventura, A.; Senga Kiessé, T.; Cazacliu, B.; Idir, R.; Werf, H.M.G. Sensitivity Analysis of Environmental Process Modeling in a Life Cycle Context: A Case Study of Hemp Crop Production. *J. Ind. Ecol.* **2015**, *19*, 978–993. [\[CrossRef\]](#)
88. Abeliotis, K.; Barla, S.-A.; Detsis, V.; Malindretos, G. Life Cycle Assessment of Carnation Production in Greece. *J. Clean. Prod.* **2016**, *112*, 32–38. [\[CrossRef\]](#)
89. Deytieux, V.; Nemecek, T.; Freiermuth Knuchel, R.; Gaillard, G.; Munier-Jolain, N.M. Is Integrated Weed Management Efficient for Reducing Environmental Impacts of Cropping Systems? A Case Study Based on Life Cycle Assessment. *Eur. J. Agron.* **2012**, *36*, 55–65. [\[CrossRef\]](#)
90. Peña, N.; Knudsen, M.T.; Fantke, P.; Antón, A.; Hermansen, J.E. Freshwater Ecotoxicity Assessment of Pesticide Use in Crop Production: Testing the Influence of Modeling Choices. *J. Clean. Prod.* **2019**, *209*, 1332–1341. [\[CrossRef\]](#)
91. Xue, X.; Hawkins, T.R.; Ingwersen, W.W.; Smith, R.L. Demonstrating an Approach for Including Pesticide Use in Life-Cycle Assessment: Estimating Human and Ecosystem Toxicity of Pesticide Use in Midwest Corn Farming. *Int. J. Life Cycle Assess.* **2015**, *20*, 1117–1126. [\[CrossRef\]](#)
92. Juraske, R.; Sanjuán, N. Life Cycle Toxicity Assessment of Pesticides Used in Integrated and Organic Production of Oranges in the Comunidad Valenciana, Spain. *Chemosphere* **2011**, *82*, 956–962. [\[CrossRef\]](#)
93. Khoshnevisan, B.; Bolandnazar, E.; Shamshirband, S.; Shariati, H.M.; Anuar, N.B.; Mat Kiah, M.L. Decreasing Environmental Impacts of Cropping Systems Using Life Cycle Assessment (LCA) and Multi-Objective Genetic Algorithm. *J. Clean. Prod.* **2015**, *86*, 67–77. [\[CrossRef\]](#)
94. Schmidt Rivera, X.C.; Bacenetti, J.; Fusi, A.; Niero, M. The Influence of Fertiliser and Pesticide Emissions Model on Life Cycle Assessment of Agricultural Products: The Case of Danish and Italian Barley. *Sci. Total Environ.* **2017**, *592*, 745–757. [\[CrossRef\]](#) [\[PubMed\]](#)
95. Brodt, S.; Kramer, K.J.; Kendall, A.; Feenstra, G. Comparing Environmental Impacts of Regional and National-Scale Food Supply Chains: A Case Study of Processed Tomatoes. *Food Policy* **2013**, *42*, 106–114. [\[CrossRef\]](#)
96. Theurl, M.C.; Haberl, H.; Erb, K.-H.; Lindenthal, T. Contrasted Greenhouse Gas Emissions from Local versus Long-Range Tomato Production. *Agron. Sustain. Dev.* **2014**, *34*, 593–602. [\[CrossRef\]](#)
97. Iriarte, A.; Rieradevall, J.; Gabarrell, X. Life Cycle Assessment of Sunflower and Rapeseed as Energy Crops under Chilean Conditions. *J. Clean. Prod.* **2010**, *18*, 336–345. [\[CrossRef\]](#)
98. Schmidt, J.H. Comparative Life Cycle Assessment of Rapeseed Oil and Palm Oil. *Int. J. Life Cycle Assess.* **2010**, *15*, 183–197. [\[CrossRef\]](#)
99. Noya, I.; González-García, S.; Bacenetti, J.; Arroja, L.; Moreira, M.T. Comparative Life Cycle Assessment of Three Representative Feed Cereals Production in the Po Valley (Italy). *J. Clean. Prod.* **2015**, *99*, 250–265. [\[CrossRef\]](#)
100. Nemecek, T.; Kägi, T. Life Cycle Inventories of Agricultural Production Systems. *Final Rep. Ecoinvent* **2007**, 361. [\[CrossRef\]](#)
101. Antón, A.; Torrellas, M.; Núñez, M.; Seigné, E.; Amores, M.J.; Muñoz, P.; Montero, J.I. Improvement of Agricultural Life Cycle Assessment Studies through Spatial Differentiation and New Impact Categories: Case Study on Greenhouse Tomato Production. *Environ. Sci. Technol.* **2014**, *48*, 9454–9462. [\[CrossRef\]](#) [\[PubMed\]](#)
102. Shrestha, P.; Karim, R.A.; Sieverding, H.L.; Archer, D.W.; Kumar, S.; Nleya, T.; Graham, C.J.; Stone, J.J. Life Cycle Assessment of Wheat Production and Wheat-Based Crop Rotations. *J. Environ. Qual.* **2020**, *49*, 1515–1529. [\[CrossRef\]](#) [\[PubMed\]](#)
103. Gentil, C.; Basset-Mens, C.; Manteaux, S.; Mottes, C.; Maillard, E.; Biard, Y.; Fantke, P. Coupling Pesticide Emission and Toxicity Characterization Models for LCA: Application to Open-Field Tomato Production in Martinique. *J. Clean. Prod.* **2020**, *277*, 124099. [\[CrossRef\]](#)
104. Brentrup, F.; Küsters, J.; Kuhlmann, H.; Lammel, J. Environmental Impact Assessment of Agricultural Production Systems Using the Life Cycle Assessment Methodology. *Eur. J. Agron.* **2004**, *20*, 247–264. [\[CrossRef\]](#)
105. Salomone, R.; Ioppolo, G. Environmental Impacts of Olive Oil Production: A Life Cycle Assessment Case Study in the Province of Messina (Sicily). *J. Clean. Prod.* **2012**, *28*, 88–100. [\[CrossRef\]](#)
106. Peano, C.; Baudino, C.; Tecco, N.; Girgenti, V. Green Marketing Tools for Fruit Growers Associated Groups: Application of the Life Cycle Assessment (LCA) for Strawberries and Berry Fruits Ecobranding in Northern Italy. *J. Clean. Prod.* **2015**, *104*, 59–67. [\[CrossRef\]](#)