

Article

A New Best Proximity Point Result with an Application to Nonlinear Fredholm Integral Equations

Hakan Sahin 

Department of Mathematics, Faculty of Science and Arts, Amasya University, Amasya 05000, Turkey;
hakan.sahin@amasya.edu.tr

Abstract: In the current paper, we first introduce a new class of contractions via a new notion called p -cyclic contraction mapping by combining the ideas of cyclic contraction mapping and p -contraction mapping. Then, we give a new definition of a cyclically 0-complete pair to weaken the completeness condition on the partial metric spaces. Following that, we prove some best proximity point results for p -cyclic contraction mappings on $D \cup E$ where (D, E) is a cyclically 0-complete pair in the setting of partial metric spaces. Hence, we generalize and unify famous and well-known results in the literature of metric fixed point theory. Additionally, we present some nontrivial examples to compare our results with earlier. Finally, we investigate the sufficient conditions for the existence of a solution to nonlinear Fredholm integral equations by the results in the paper.

Keywords: best proximity point; p -cyclic contractions; nonlinear Fredholm integral equations

MSC: 54H25; 47H10; 45B05



Citation: Sahin, H. A New Best Proximity Point Result with an Application to Nonlinear Fredholm Integral Equations. *Mathematics* **2022**, *10*, 665. <https://doi.org/10.3390/math10040665>

Academic Editor: Simeon Reich

Received: 3 February 2022

Accepted: 18 February 2022

Published: 20 February 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Banach [1] proved a fixed point result, which is known as Banach contraction principle, in 1922. In this result, it has been shown that every self mapping $Y : \Lambda \rightarrow \Lambda$ on a complete metric space (Λ, ρ) such that there is $q \in [0, 1)$ such that

$$\rho(Y\kappa, Y\zeta) \leq q\rho(\kappa, \zeta)$$

for all $\kappa, \zeta \in \Lambda$ has a unique fixed point in Λ .

Banach contraction principle has been considered the beginning of metric fixed point theory. Then, many authors have generalized and improved it in different ways [2–4]. In this sense, Popescu [5] introduced the concept of p -contraction mapping and obtained a fixed point result for these mappings. According to this result, every p -contraction mapping $Y : \Lambda \rightarrow \Lambda$ on a complete metric space (Λ, ρ) , that is, there exists q in $[0, 1)$ such that

$$\rho(Y\kappa, Y\zeta) \leq q[\rho(\kappa, \zeta) + |\rho(\kappa, Y\kappa) - \rho(\zeta, Y\zeta)|]$$

for all $\kappa, \zeta \in \Lambda$ has a unique fixed point.

Recently, Kirk et al. [6] has obtained a new generalization of Banach contraction principle via a new concept of cyclic mapping. In their result, the cyclic mapping may not be continuous, unlike the Banach's result. This is the important feature of their result. Then, many researchers have studied to obtain some fixed point results for cyclic mappings [7,8].

On the other hand, one of the interesting generalizations of Banach contraction principle has been obtained by taking into account nonself mappings. Let $\emptyset \neq D, E$ be subsets of a metric space (Λ, ρ) and $Y : D \rightarrow E$ be a nonself mapping. If $D \cap E = \emptyset$, Y cannot have a fixed point. Then, since $\rho(\kappa, Y\kappa) \geq \rho(D, E)$ for all $\kappa \in D$, it is reasonable to search a point κ satisfying $\rho(\kappa, Y\kappa) = \rho(D, E)$. This point is said to be a best proximity point of Y . Note

that a best proximity point of Y is an optimal solution for the problem $\min_{x \in D} \rho(x, Yx)$. Additionally, a best proximity point turns into a fixed point in the case of $D = E = \Lambda$. Therefore, many authors have studied on this topic [9–15].

Taking into account the ideas of both best proximity point and cyclic mapping the famous concept of cyclic contraction mapping was introduced by Eldred and Veeramani [16]. In this way, these ideas were unified.

Until now, some properties such as bounded compactness have been used to guarantee the existence of best proximity points for cyclic contraction mappings. Recently, introducing a nice notion called cyclically completeness, Karpagam and Agrawal [17] show the existence of best proximity point of a cyclic contraction mapping without using the property of bounded compactness. Then, many authors have obtained some best proximity point results with the help of this concept [18,19].

In 1994, motivated by the experience of computer science, Matthews, Ref. [20], relaxed the condition $x = \zeta$ implies $\rho(x, \zeta) = 0$ in a metric space (Λ, ρ) by introducing the partial metric spaces. Following that, many authors obtained both various fixed point results and best proximity point results in the settings of partial metric spaces [21–23]. Very recently, Romaguera [24] introduced the concept of 0-complete partial metric space. Hence, a weaker form of completeness on partial metric spaces has been obtained.

In this paper, we aim to extend and unify some famous results in the literature of metric fixed point theory, such as the main results of Eldred-Veeramani [16] and Popescu [5]. Hence, we first introduce a new class of contractions via a new notion called p -cyclic contraction mapping by combining the ideas of cyclic contraction mapping and p -contraction mapping. Then, we give a new definition of a cyclically 0-complete pair to weaken the completeness condition on the partial metric spaces. Following, we prove some best proximity point results for p -cyclic contraction mappings on $D \cup E$ where (D, E) is a cyclically 0-complete pair in a partial metric space. Additionally, we present some nontrivial examples to show the effectiveness of our work. Finally, we investigate the sufficient conditions for the existence of a solution to nonlinear Fredholm integral equations by the results in the paper.

2. Preliminaries

In this section, we give some definitions, lemmas and theorems which are important in our main result. We begin this section with the following result for cyclic mappings, which was obtained by Kirk et al. [6].

Theorem 1 ([6]). *Let $\emptyset \neq D, E$ be closed subsets of a complete metric space (Λ, ρ) and $Y : D \cup E \rightarrow D \cup E$ be a cyclic mapping, that is, $Y(D) \subseteq E$ and $Y(E) \subseteq D$. Then, Y has a fixed point in $D \cap E$ if there is $q \in [0, 1)$ such that*

$$\rho(Yx, Y\zeta) \leq q\rho(x, \zeta)$$

for all $x \in D$ and $\zeta \in E$.

Taking into account $D \cap E = \emptyset$ in Theorem 1 the famous concept of cyclic contraction mapping was introduced by Eldred and Veeramani [16]. Then, they obtain a best proximity point result as follows:

Definition 1 ([16]). *Let $\emptyset \neq D, E$ be subsets of a metric space (Λ, ρ) and $Y : D \cup E \rightarrow D \cup E$ be a cyclic mapping. Then, $Y : D \cup E \rightarrow D \cup E$ is called cyclic contraction mapping if there exists q in $[0, 1)$ such that*

$$\rho(Yx, Y\zeta) \leq q\rho(x, \zeta) + (1 - q)\rho(D, E)$$

for all $x \in D$ and $\zeta \in E$.

Theorem 2 ([16]). Let (Λ, ρ) be a metric space, $\emptyset \neq D, E \subseteq \Lambda$ where D, E are closed and $Y : D \cup E \rightarrow D \cup E$ be a cyclic contraction mapping. If either D or E is boundedly compact, then Y has a best proximity point in $D \cup E$.

Next, we recall the concept of cyclically completeness.

Definition 2 ([17]). Let (Λ, ρ) be a metric space and $\emptyset \neq D, E \subseteq \Lambda$. A sequence $\{\varkappa_n\}$ in $D \cup E$ with $\{\varkappa_{2n}\} \subset D$ and $\{\varkappa_{2n+1}\} \subset E$ is called a cyclically Cauchy sequence if for each $\varepsilon > 0$ there is $n_0 \in \mathbb{N}$ satisfying

$$\rho(\varkappa_n, \varkappa_m) < \rho(D, E) + \varepsilon,$$

for all $n, m \geq n_0$ with m is odd, n is even.

Definition 3 ([17]). A pair (D, E) of subsets of a metric space is said to be cyclically complete if for every cyclically Cauchy sequence $\{\varkappa_n\}$ in $D \cup E$, either $\{\varkappa_{2n}\}$ or $\{\varkappa_{2n+1}\}$ are convergent.

Now, we give the definition of the partial metric space and its topological properties.

Definition 4 ([20]). Let $\Lambda \neq \emptyset$ and $\theta : \Lambda \times \Lambda \rightarrow [0, \infty)$ be a mapping satisfying the following conditions:

- (θ_1) $\theta(\varkappa, \varkappa) = \theta(\varkappa, \zeta) = \theta(\zeta, \zeta)$ if and only if $\varkappa = \zeta$,
- (θ_2) $\theta(\varkappa, \varkappa) \leq \theta(\varkappa, \zeta)$,
- (θ_3) $\theta(\varkappa, \zeta) = \theta(\zeta, \varkappa)$,
- (θ_4) $\theta(\varkappa, \eta) \leq \theta(\varkappa, \zeta) + \theta(\zeta, \eta) - \theta(\zeta, \zeta)$.

for all $\varkappa, \zeta, \eta \in \Lambda$. Then, θ is called a partial metric. Additionally, (Λ, θ) is called a partial metric space.

It can be easily seen that every metric space is a partial metric space, but the converse may not be true (see for more details [25–28]). Now, assume that θ is a partial metric on Λ . Then, there is an T_0 topology τ_θ on Λ . Additionally, the family open θ -balls

$$\{B_\theta(\varkappa, \varepsilon) : \varkappa \in \Lambda, \varepsilon > 0\}$$

is a base for the topology τ_θ where

$$B_\theta(\varkappa, \varepsilon) = \{\zeta \in \Lambda : \theta(\varkappa, \zeta) < \theta(\varkappa, \varkappa) + \varepsilon\}$$

for all $\varkappa \in \Lambda$ and $\varepsilon > 0$. If we take a sequence $\{\varkappa_n\} \subseteq \Lambda$ and $\varkappa \in \Lambda$, then it is clear that $\{\varkappa_n\}$ converges to \varkappa w.r.t. τ_θ if and only if

$$\lim_{n \rightarrow \infty} \theta(\varkappa_n, \varkappa) = \theta(\varkappa, \varkappa).$$

The sequence $\{\varkappa_n\}$ is said to be Cauchy sequence if $\lim_{n, m \rightarrow \infty} \theta(\varkappa_n, \varkappa_m)$ exists and is finite. Additionally, (Λ, θ) is said to be a complete partial metric space if every Cauchy sequence $\{\varkappa_n\}$ converges to a point \varkappa in Λ w.r.t. τ_θ such that

$$\lim_{n, m \rightarrow \infty} \theta(\varkappa_n, \varkappa_m) = \theta(\varkappa, \varkappa).$$

Definition 5 ([24]). Let (Λ, θ) be a partial metric space and $\{\varkappa_n\}$ be a sequence in Λ .

(i) $\{\varkappa_n\}$ is called 0-Cauchy sequence if

$$\lim_{n, m \rightarrow \infty} \theta(\varkappa_n, \varkappa_m) = 0.$$

- (ii) (Λ, θ) is called 0-complete partial metric space if every 0-Cauchy sequence converges to a point \varkappa in Λ w.r.t. τ_θ such that

$$\lim_{n,m \rightarrow \infty} \theta(\varkappa_n, \varkappa_m) = \theta(\varkappa, \varkappa).$$

If θ is a partial metric on Λ , then the mapping $\rho_\theta : \Lambda \times \Lambda \rightarrow [0, \infty)$ defined by

$$\rho_\theta(\varkappa, \zeta) = 2\theta(\varkappa, \zeta) - \theta(\varkappa, \varkappa) - \theta(\zeta, \zeta)$$

for all $\varkappa, \zeta \in \Lambda$ is an ordinary metric on Λ .

The following lemma shows the relation between a partial metric θ and ordinary metric ρ_θ .

Lemma 1. Let (Λ, θ) be a partial metric space.

- (i) $\{\varkappa_n\}$ is a Cauchy sequence in (Λ, θ) if and only if $\{\varkappa_n\}$ is a Cauchy sequence in (Λ, ρ_θ) ,
(ii) (Λ, θ) is a complete partial metric space if and only if (Λ, ρ_θ) is a complete metric space,
(iii) Given a sequence $\{\varkappa_n\}$ in Λ and $\varkappa \in \Lambda$. Then, we have

$$\lim_{n \rightarrow \infty} \rho_\theta(\varkappa_n, \varkappa) = 0 \iff \theta(\varkappa, \varkappa) = \lim_{n \rightarrow \infty} \theta(\varkappa_n, \varkappa) = \lim_{n,m \rightarrow \infty} \theta(\varkappa_n, \varkappa_m).$$

3. Main Results

We first give the definition of p -cyclic contraction mapping on partial metric spaces.

Definition 6. Let $\emptyset \neq D, E$ be subsets of a partial metric space (Λ, θ) and $Y : D \cup E \rightarrow D \cup E$ be a cyclic mapping. Then, Y is said to be p -cyclic contraction mapping if there is q in $[0, 1)$ such that

$$\theta(Y\varkappa, Y\zeta) \leq q\{\theta(\varkappa, \zeta) + |\theta(\varkappa, Y\varkappa) - \theta(\zeta, Y\zeta)|\} + (1 - q)\theta(D, E) \quad (1)$$

for all $\varkappa \in D$ and $\zeta \in E$ where $\theta(D, E) = \inf\{\theta(\varkappa, \zeta) : \varkappa \in D \text{ and } \zeta \in E\}$.

The following example shows that the classes of cyclic contractions and p -contractions are proper subsets of the class of p -cyclic contractions.

Example 1. Let $\Lambda = \{(\varkappa_1, \varkappa_2, \dots, \varkappa_n, \dots) \subseteq \mathbb{R} : \varkappa_1 \geq 0 \text{ and } \sup_{i \in \mathbb{N}} |\varkappa_i| < \infty\}$ and $\theta : \Lambda \times \Lambda \rightarrow \mathbb{R}$ be a function defined by

$$\theta(\varkappa, \zeta) = \max\{\varkappa_1, \zeta_1\} + \sup_{i \geq 2} |\varkappa_i - \zeta_i|$$

for $\varkappa = (\varkappa_1, \varkappa_2, \dots, \varkappa_n, \dots)$, $\zeta = (\zeta_1, \zeta_2, \dots, \zeta_n, \dots) \in \Lambda$. It is clear that (Λ, θ) is a partial metric space. Consider the following subsets

$$D = \{(t, 1, \dots, 1, \dots) : t \in [0, 1]\} \cup \{(1, -1, \dots, -1, \dots)\},$$

and

$$E = \{(t, 0, \dots, 0, \dots) : t \in [0, 1]\} \cup \{(0, 2, \dots, 2, \dots)\}.$$

Then, we have $\theta(D, E) = 1$. Let define a mapping $Y : D \cup E \rightarrow D \cup E$ by

$$Y\varkappa = \begin{cases} (\frac{t}{2}, 0, \dots, 0, \dots) & , \quad \varkappa = (t, 1, \dots, 1, \dots) \in D \\ (0, 0, \dots, 0, \dots) & , \quad \varkappa = (1, -1, \dots, -1, \dots) \\ (1, -1, \dots, -1, \dots) & , \quad \varkappa = (0, 2, \dots, 2, \dots) \\ (\frac{t}{2}, 1, \dots, 1, \dots) & , \quad \varkappa = (t, 0, \dots, 0, \dots) \in E \end{cases}.$$

Now, we shall show that Y is a p -cyclic contraction mapping. Then, we have the following four conditions:

Case 1: Let $\varkappa = (t, 1, \dots, 1, \dots) \in D$ and $\zeta = (s, 0, \dots, 0, \dots) \in E$. In this case, we have

$$\begin{aligned}\theta(Y\varkappa, Y\zeta) &= \max\left\{\frac{t}{2}, \frac{s}{2}\right\} + 1 \\ &= \frac{1}{2}(\max\{t, s\} + 1) + \frac{1}{2} \\ &\leq \frac{1}{2}\{\theta(\varkappa, \zeta) + |\theta(\varkappa, Y\varkappa) - \theta(\zeta, Y\zeta)|\} + \left(1 - \frac{1}{2}\right)\theta(D, E).\end{aligned}$$

Case 2: Let $\varkappa = (t, 1, \dots, 1, \dots) \in D$ and $\zeta = (0, 2, \dots, 2, \dots) \in E$. In this case, we have

$$\begin{aligned}\theta(Y\varkappa, Y\zeta) &= 2 \\ &\leq \frac{1}{2}(1 + t + 3 - t) + \frac{1}{2} \\ &= \frac{1}{2}\{\theta(\varkappa, \zeta) + |\theta(\varkappa, Y\varkappa) - \theta(\zeta, Y\zeta)|\} + \left(1 - \frac{1}{2}\right)\theta(D, E).\end{aligned}$$

Case 3: Let $\varkappa = (1, -1, \dots, -1, \dots) \in D$ and $\zeta = (t, 0, \dots, 0, \dots) \in E$. In this case, we have

$$\begin{aligned}\theta(Y\varkappa, Y\zeta) &= \frac{t}{2} + 1 \\ &\leq \frac{1}{2} \cdot 2 + \frac{1}{2} \\ &\leq \frac{1}{2}\{\theta(\varkappa, \zeta) + |\theta(\varkappa, Y\varkappa) - \theta(\zeta, Y\zeta)|\} + \left(1 - \frac{1}{2}\right)\theta(D, E).\end{aligned}$$

Case 4: Let $\varkappa = (1, -1, \dots, -1, \dots) \in D$ and $\zeta = (0, 2, \dots, 2, \dots) \in E$. In this case, we have

$$\begin{aligned}\theta(Y\varkappa, Y\zeta) &= 2 \\ &\leq \frac{1}{2} \cdot 4 + \frac{1}{2} \\ &\leq \frac{1}{2}\{\theta(\varkappa, \zeta) + |\theta(\varkappa, Y\varkappa) - \theta(\zeta, Y\zeta)|\} + \left(1 - \frac{1}{2}\right)\theta(D, E).\end{aligned}$$

Hence, Y is a p -cyclic contraction mapping for $q = \frac{1}{2}$. However, Y is not neither a cyclic contraction mapping nor a p -contraction mapping. If we take $\varkappa = (0, 1, \dots, 1, \dots) \in D$ and $\zeta = (0, 2, \dots, 2, \dots) \in E$, then we have

$$\theta(Y\varkappa, Y\zeta) = 2 > 1 = q\theta(\varkappa, \zeta) + (1 - q)\theta(D, E)$$

for all $q \in [0, 1)$ which implies that Y is not a cyclic contraction mapping. Additionally, if we take $\varkappa = (0, 1, \dots, 1, \dots) \in D$ and $\zeta = (0, 0, \dots, 0, \dots) \in E$, then we have

$$\theta(Y\varkappa, Y\zeta) = 1 > q = q\{\theta(\varkappa, \zeta) + |\theta(\varkappa, Y\varkappa) - \theta(\zeta, Y\zeta)|\}$$

for all $q \in [0, 1)$, which implies that Y is not a p -contraction mapping.

Then, we restate the definition of cyclically Cauchy sequence in the settings of partial metric spaces.

Definition 7. Let $\emptyset \neq D, E$ be subsets of a partial metric space (Λ, θ) . A sequence $\{\varkappa_n\}$ in $D \cup E$ with $\{\varkappa_{2n}\} \subseteq D$ and $\{\varkappa_{2n+1}\} \subseteq E$ is called a cyclically Cauchy sequence if for each $\varepsilon > 0$ there is $n_0 \in \mathbb{N}$ such that

$$\theta(\varkappa_n, \varkappa_m) < \theta(D, E) + \varepsilon$$

for all $n, m \geq n_0$ with m is odd, n is even.

Note that, if $\theta(D, E) = 0$ in Definition 7, then the definition of cyclically Cauchy sequence turns into the definition of 0-Cauchy sequence.

Now, we introduce the definition of cyclically 0-complete pair in a partial metric spaces.

Definition 8. Let $\emptyset \neq D, E$ be subsets of a partial metric space (Λ, θ) . A pair (D, E) is said to be cyclically 0-complete pair if for every cyclically Cauchy sequence $\{\varkappa_n\}$ in $D \cup E$, either the sequence $\{\varkappa_{2n}\}$ has a convergent subsequence $\{\varkappa_{2n_k}\}$ to a point $\varkappa^* \in D$ w.r.t. τ_θ such that

$$\lim_{k,l \rightarrow \infty} \theta(\varkappa_{2n_k}, \varkappa_{2n_l}) = \theta(\varkappa^*, \varkappa^*) = 0 \quad (2)$$

or $\{\varkappa_{2n+1}\}$ has a convergent subsequence $\{\varkappa_{2n_k+1}\}$ to a point $\zeta^* \in E$ w.r.t. τ_θ such that

$$\lim_{k,l \rightarrow \infty} \theta(\varkappa_{2n_k+1}, \varkappa_{2n_l+1}) = \theta(\zeta^*, \zeta^*) = 0. \quad (3)$$

Remark 1. If D or E is a closed subset of 0-complete partial metric space (Λ, θ) and $\theta(D, E) = 0$, then (D, E) is a cyclically 0-complete pair. However, if (D, E) is a cyclically 0-complete pair, then D and E are not necessarily 0-complete. The following example shows this fact.

Example 2. Let $\Lambda = [0, \infty) \times \mathbb{R}$ and $\theta : \Lambda \times \Lambda \rightarrow [0, \infty)$ be a function defined as

$$\theta(\varkappa, \zeta) = \varkappa_1 + \zeta_1 + |\varkappa_2 - \zeta_2|$$

for all $\varkappa = (\varkappa_1, \varkappa_2), \zeta = (\zeta_1, \zeta_2) \in \Lambda$. Then, (Λ, θ) is a partial metric space. Let's take the subsets $D = [0, 1) \times (-1, 0]$ and $E = [0, 1) \times [1, 2)$ of Λ . In this case, we have $\theta(D, E) = 1$. Now, we claim that (D, E) is a cyclically 0-complete pair. For this, let's take a cyclically Cauchy sequence $\{\varkappa_n\}$ in $D \cup E$ with $\{\varkappa_{2n}\} \subseteq D$ and $\{\varkappa_{2n+1}\} \subseteq E$. Then, for each $\varepsilon > 0$ there is $n_0 \in \mathbb{N}$ such that

$$\theta(\varkappa_n, \varkappa_m) < 1 + \varepsilon$$

for all $n, m \geq n_0$ with m is odd, n is even. Hence, we have that $\{\varkappa_{2n}\}$ converges to $(0, 0) \in D$ w.r.t. τ_θ and

$$\lim_{n,k \rightarrow \infty} \theta(\varkappa_{2n}, \varkappa_{2k}) = \theta((0, 0), (0, 0)) = 0,$$

that is, the sequence $\{\varkappa_{2n}\}$ has a subsequence satisfying (2). However, neither D nor E is 0-complete. Indeed, if we take a sequence $\{\varkappa_n\} = \left\{ \left(0, -1 + \frac{1}{n} \right) \right\}$ in D , then we have $\lim_{n,m \rightarrow \infty} \theta(\varkappa_n, \varkappa_m) = 0$. Hence, the sequence $\{\varkappa_n\}$ is a 0-Cauchy sequence in D , but it is not convergent in D . Hence, D is not 0-complete. Similarly, we can show that E is not 0-complete by considering the sequence $\{x_n\} = \left\{ \left(0, 2 - \frac{1}{n} \right) \right\}$ in E .

Now, we give a new definition in partial metric spaces.

Definition 9. Let $\emptyset \neq D$ be a subset of a partial metric space (Λ, θ) . Then, D is called 0-boundedly compact if every bounded sequence $\{\varkappa_n\}$ has a convergent subsequence $\{\varkappa_{n_k}\}$ to a point $\varkappa^* \in D$ w.r.t. τ_θ such that

$$\lim_{k,l \rightarrow \infty} \theta(\varkappa_{n_k}, \varkappa_{n_l}) = \theta(\varkappa^*, \varkappa^*) = 0.$$

Remark 2. Note that if either D or E is a 0-boundedly compact, then the pair (D, E) is a cyclically 0-complete pair. However, the converse may not be true. Example 2 can be given to show this fact.

Proposition 1. Let (Λ, θ) be a partial metric space and $\emptyset \neq D, E \subseteq \Lambda$. Suppose that $Y : D \cup E \rightarrow D \cup E$ is a p -cyclic contraction mapping. If for any sequence $\{x_n\}$ defined by $x_{n+1} = Yx_n$ with the initial point $x_0 \in D$, there is $n_0 \in \mathbb{N}$ such that

$$\theta(x_{n_0}, x_{n_0+1}) \leq \theta(x_{n_0+1}, x_{n_0+2}),$$

then Y has a best proximity point in $D \cup E$.

Proof. Assume that $\{x_n\}$ is an arbitrary sequence defined by $x_{n+1} = Yx_n$ with the initial point $x_0 \in D \cup E$. Since Y is a p -cyclic contraction mapping, there is q in $[0, 1)$ such that

$$\begin{aligned} \theta(x_n, x_{n+1}) &= \theta(Yx_{n-1}, Yx_n) \\ &\leq q\{\theta(x_{n-1}, x_n) + |\theta(x_{n-1}, Yx_{n-1}) - \theta(x_n, Yx_n)|\} \\ &\quad + (1-q)\theta(D, E) \end{aligned} \quad (4)$$

for all $n \in \mathbb{N}$. Now, if there is $n_0 \in \mathbb{N}$ such that

$$\theta(x_{n_0}, x_{n_0+1}) \leq \theta(x_{n_0+1}, x_{n_0+2}),$$

then from (4) we get

$$\begin{aligned} \theta(x_{n_0+1}, x_{n_0+2}) &= \theta(Yx_{n_0}, Yx_{n_0+1}) \\ &\leq q\{\theta(x_{n_0}, x_{n_0+1}) + |\theta(x_{n_0}, Yx_{n_0}) - \theta(x_{n_0+1}, Yx_{n_0+1})|\} \\ &\quad + (1-q)\theta(D, E) \\ &= q\{\theta(x_{n_0}, x_{n_0+1}) + |\theta(x_{n_0}, x_{n_0+1}) - \theta(x_{n_0+1}, x_{n_0+2})|\} \\ &\quad + (1-q)\theta(D, E) \\ &= q\theta(x_{n_0}, x_{n_0+1}) - q\theta(x_{n_0}, x_{n_0+1}) + q\theta(x_{n_0+1}, x_{n_0+2}) \\ &\quad + (1-q)\theta(D, E) \\ &= q\theta(x_{n_0+1}, x_{n_0+2}) + (1-q)\theta(D, E). \end{aligned}$$

Hence, we have

$$\theta(x_{n_0+1}, x_{n_0+2}) \leq \theta(D, E).$$

Additionally, since $\theta(D, E) \leq \theta(x_{n_0+1}, x_{n_0+2})$, we get

$$\theta(x_{n_0+1}, x_{n_0+2}) = \theta(D, E).$$

Then, we obtain

$$\theta(D, E) \leq \theta(x_{n_0}, Yx_{n_0}) \leq \theta(x_{n_0+1}, Yx_{n_0+1}) = \theta(x_{n_0+1}, x_{n_0+2}) = \theta(D, E)$$

Hence, x_{n_0} and x_{n_0+1} are best proximity points of Y . \square

Remark 3. If for the sequence $\{x_n\}$ mentioned in Proposition 1 there is $n_0 \in \mathbb{N}$ such that $\theta(x_{n_0}, x_{n_0+1}) \leq \theta(x_{n_0+1}, x_{n_0+2})$, then Y has a best proximity point in $D \cup E$. Therefore, we will investigate the condition $\theta(x_{n+1}, x_{n+2}) \leq \theta(x_n, x_{n+1})$ for all $n \in \mathbb{N}$ in the rest of paper.

Proposition 2. Let (Λ, θ) be a partial metric space and $\emptyset \neq D, E \subseteq \Lambda$. Assume that $Y : D \cup E \rightarrow D \cup E$ is a p -cyclic contraction mapping. Then, for every sequence $\{x_n\} \subseteq D \cup E$ created as in Proposition 1, we have $\theta(x_n, x_{n+1}) \rightarrow \theta(D, E)$ as $n \rightarrow \infty$.

Proof. Let $\{\mathcal{X}_n\} \subseteq D \cup E$ be a sequence constructed as in Proposition 1. Since Y is a p -cyclic contraction mapping, considering Remark 3 we have

$$\begin{aligned}\theta(\mathcal{X}_n, \mathcal{X}_{n+1}) &= \theta(Y\mathcal{X}_{n-1}, Y\mathcal{X}_n) \\ &\leq q\{\theta(\mathcal{X}_{n-1}, \mathcal{X}_n) + |\theta(\mathcal{X}_{n-1}, Y\mathcal{X}_{n-1}) - \theta(\mathcal{X}_n, Y\mathcal{X}_n)|\} + (1-q)\theta(D, E) \\ &= q\{\theta(\mathcal{X}_{n-1}, \mathcal{X}_n) + |\theta(\mathcal{X}_{n-1}, \mathcal{X}_n) - \theta(\mathcal{X}_n, \mathcal{X}_{n+1})|\} + (1-q)\theta(D, E) \\ &= q\{\theta(\mathcal{X}_{n-1}, \mathcal{X}_n) + \theta(\mathcal{X}_{n-1}, \mathcal{X}_n) - \theta(\mathcal{X}_n, \mathcal{X}_{n+1})\} + (1-q)\theta(D, E) \\ &= 2q\theta(\mathcal{X}_{n-1}, \mathcal{X}_n) - q\theta(\mathcal{X}_n, \mathcal{X}_{n+1}) + (1-q)\theta(D, E),\end{aligned}$$

and so we get

$$\theta(\mathcal{X}_n, \mathcal{X}_{n+1}) \leq \alpha\theta(\mathcal{X}_{n-1}, \mathcal{X}_n) + \beta\theta(D, E)$$

for all $n \in \mathbb{N}$, where $\alpha = \frac{2q}{1+q} < 1$ and $\beta = \frac{1-q}{1+q}$. By using the last inequality, we have

$$\begin{aligned}\theta(D, E) &\leq \theta(\mathcal{X}_n, \mathcal{X}_{n+1}) \\ &\leq \alpha\theta(\mathcal{X}_{n-1}, \mathcal{X}_n) + \beta\theta(D, E) \\ &\leq \alpha(\alpha\theta(\mathcal{X}_{n-2}, \mathcal{X}_{n-1}) + \beta\theta(D, E)) + \beta\theta(D, E) \\ &= \alpha^2\theta(\mathcal{X}_{n-2}, \mathcal{X}_{n-1}) + \beta\theta(D, E)(1 + \alpha) \\ &\vdots \\ &\leq \alpha^n\theta(\mathcal{X}_0, \mathcal{X}_1) + \beta\theta(D, E)(1 + \alpha + \dots + \alpha^{n-1}) \\ &= \alpha^n\theta(\mathcal{X}_0, \mathcal{X}_1) + \beta\theta(D, E)\left(\frac{1 - \alpha^n}{1 - \alpha}\right) \\ &= \alpha^n\theta(\mathcal{X}_0, \mathcal{X}_1) + \theta(D, E)(1 - \alpha^n)\end{aligned}$$

for all $n \in \mathbb{N}$. Hence, we get

$$\lim_{n \rightarrow \infty} \theta(\mathcal{X}_n, \mathcal{X}_{n+1}) = \theta(D, E). \quad (5)$$

□

The following proposition is crucial for our main result.

Proposition 3. Let $\emptyset \neq D, E$ be subsets of a partial metric space (Λ, θ) . Assume that $Y : D \cup E \rightarrow D \cup E$ is a p -cyclic contraction mapping. Then, every sequence $\{\mathcal{X}_n\} \subseteq D \cup E$ created as in Proposition 1 is bounded.

Proof. Let $\{\mathcal{X}_n\} \subseteq D \cup E$ be a sequence constructed as in Proposition 1. Hence, from Proposition 2 the sequence $\{\theta(\mathcal{X}_{2n}, \mathcal{X}_{2n+1})\}$ converges to $\theta(D, E)$ as $n \rightarrow \infty$, and so the sequence $\{\theta(\mathcal{X}_{2n}, \mathcal{X}_{2n+1})\}$ is bounded. Then, there exists $L > 0$ such that

$$\theta(\mathcal{X}_{2n}, \mathcal{X}_{2n+1}) \leq L$$

for all $n \in \mathbb{N}$. Since Y is a p -cyclic contraction mapping, considering Remark 3, we have

$$\begin{aligned}\theta(\mathcal{X}_{2n+1}, \mathcal{X}_1) &\leq \theta(\mathcal{X}_{2n+1}, \mathcal{X}_{2n+2}) + \theta(\mathcal{X}_{2n+2}, \mathcal{X}_1) \\ &\leq L + \theta(Y\mathcal{X}_{2n+1}, Y\mathcal{X}_0) \\ &\leq L + q\{\theta(\mathcal{X}_{2n+1}, \mathcal{X}_0) + |\theta(\mathcal{X}_{2n+1}, Y\mathcal{X}_{2n+1}) - \theta(\mathcal{X}_0, Y\mathcal{X}_0)|\} + (1-q)\theta(D, E) \\ &= L + q\{\theta(\mathcal{X}_{2n+1}, \mathcal{X}_0) + |\theta(\mathcal{X}_{2n+1}, \mathcal{X}_{2n+2}) - \theta(\mathcal{X}_0, \mathcal{X}_1)|\} + (1-q)\theta(D, E) \\ &= L + q\theta(\mathcal{X}_{2n+1}, \mathcal{X}_0) - q\theta(\mathcal{X}_{2n+1}, \mathcal{X}_{2n+2}) + q\theta(\mathcal{X}_0, \mathcal{X}_1) + (1-q)\theta(D, E) \\ &\leq L + q\theta(\mathcal{X}_{2n+1}, \mathcal{X}_1) + 2q\theta(\mathcal{X}_0, \mathcal{X}_1) + (1-q)\theta(D, E)\end{aligned}$$

for each $n \in \mathbb{N}$ which implies that

$$\theta(x_{2n+1}, x_1) \leq \frac{L}{1-q} + \frac{2q}{1-q} \theta(x_0, x_1) + \theta(D, E).$$

Let

$$M = \frac{L}{1-q} + \frac{2q}{1-q} \theta(x_0, x_1) + \theta(D, E).$$

Hence, $\{x_{2n+1}\}$ is bounded. Additionally, we get

$$\theta(x_{2n}, x_1) \leq \theta(x_{2n}, x_{2n+1}) + \theta(x_{2n+1}, x_1) \leq L + M.$$

Hence, $\{x_{2n}\}$ is bounded. Therefore, $\{x_n\}$ is bounded. \square

Theorem 3. Let $\emptyset \neq D, E$ be subsets of a partial metric space (Λ, θ) where (D, E) is a cyclically 0-complete pair. If $Y : D \cup E \rightarrow D \cup E$ is a p -cyclic contraction mapping, Y has a best proximity point.

Proof. Let $\{x_n\} \subseteq D \cup E$ be a sequence constructed as in Proposition 1 with the initial point $x_0 \in D$. If there is $n_0 \in \mathbb{N}$ such that

$$\theta(x_{n_0}, x_{n_0+1}) \leq \theta(x_{n_0+1}, x_{n_0+2}),$$

then, from Proposition 1 Y has a best proximity point. Now assume

$$\theta(x_{n+1}, x_{n+2}) \leq \theta(x_n, x_{n+1})$$

for all $n \in \mathbb{N}$. In this case, using Proposition 2, we have

$$\lim_{n \rightarrow \infty} \theta(x_n, x_{n+1}) = \theta(D, E). \quad (6)$$

Now, let us show that $\{x_n\}$ is a cyclically Cauchy sequence. Assume $n, m \in \mathbb{N}$ with $n \geq m$. Since Y is a p -cyclic contraction mapping, we get

$$\begin{aligned} \theta(x_n, x_m) &= \theta(Yx_{n-1}, Yx_{m-1}) \\ &\leq q\{\theta(x_{n-1}, x_{m-1}) + |\theta(x_{m-1}, x_m) - \theta(x_{n-1}, x_n)|\} + (1-q)\theta(D, E) \\ &= q\{\theta(x_{n-1}, x_{m-1}) + \theta(x_{m-1}, x_m) - \theta(x_{n-1}, x_n)\} + (1-q)\theta(D, E) \\ &\leq q\{\theta(x_{n-1}, x_{m-1}) + \theta(x_{m-1}, x_m) - \theta(D, E)\} + (1-q)\theta(D, E) \\ &= q\{\theta(x_{n-1}, x_{m-1}) + \theta(x_{m-1}, x_m)\} + (1-2q)\theta(D, E) \end{aligned}$$

for all $n, m \in \mathbb{N}$ with $n \geq m$. Additionally, from Proposition 2 we obtain

$$\theta(x_m, x_{m+1}) \leq \alpha \theta(x_{m-1}, x_m) + \beta \theta(D, E)$$

for all $m \in \mathbb{N}$ where $\alpha = \frac{2q}{1+q} < 1$ and $\beta = \frac{1-q}{1+q}$. Therefore, we have

$$\begin{aligned}
\theta(D, E) &\leq \theta(\mathcal{X}_n, \mathcal{X}_m) \\
&= \theta(Y\mathcal{X}_{n-1}, Y\mathcal{X}_{m-1}) \\
&\leq q\theta(\mathcal{X}_{n-1}, \mathcal{X}_{m-1}) + q\theta(\mathcal{X}_{m-1}, \mathcal{X}_m) + (1-2q)\theta(D, E) \\
&\leq q\{q\theta(\mathcal{X}_{n-2}, \mathcal{X}_{m-2}) + q\theta(\mathcal{X}_{m-2}, \mathcal{X}_{m-1}) + (1-2q)\theta(D, E)\} \\
&\quad + q\{\alpha\theta(\mathcal{X}_{m-2}, \mathcal{X}_{m-1}) + \beta\theta(D, E)\} + (1-2q)\theta(D, E) \\
&= q^2\theta(\mathcal{X}_{n-2}, \mathcal{X}_{m-2}) + (q^2 + \alpha q)\theta(\mathcal{X}_{m-2}, \mathcal{X}_{m-1}) + \beta q\theta(D, E) \\
&\quad + (1+q)(1-2q)\theta(D, E) \\
&\vdots \\
&\leq q^m\theta(\mathcal{X}_{n-m}, \mathcal{X}_0) \\
&\quad + q^m\left\{1 + \frac{\alpha}{q} + \left(\frac{\alpha}{q}\right)^2 + \left(\frac{\alpha}{q}\right)^3 + \cdots + \left(\frac{\alpha}{q}\right)^{m-1}\right\}\theta(\mathcal{X}_0, \mathcal{X}_1) \\
&\quad + \beta\left\{\begin{array}{c} \{q + q^2 + q^3 + \cdots + q^{m-2}\} \\ + \alpha\{q + q^2 + q^3 + \cdots + q^{m-1}\} \\ + \alpha^2\{q + q^2 + q^3 + \cdots + q^{m-2}\} \\ \vdots \\ + \alpha^{m-1}q \end{array}\right\}\theta(D, E) \\
&\quad + (1-2q)\{1 + q + q^2 + \cdots + q^{m-1}\}\theta(D, E) \\
&\leq q^m\theta(\mathcal{X}_0, \mathcal{X}_{n-m}) + q^m\sum_{i=0}^{m-1}\left(\frac{\alpha}{q}\right)^i\theta(\mathcal{X}_0, \mathcal{X}_1) \\
&\quad + \beta\left\{\sum_{i=1}^{\infty}q^i + \alpha\sum_{i=1}^{\infty}q^i + \alpha^2\sum_{i=1}^{\infty}q^i + \cdots + \alpha^{m-1}\sum_{i=1}^{\infty}q^i\right\}\theta(D, E) \\
&\quad + (1-2q)\sum_{i=0}^{m-1}q^i\theta(D, E) \\
&= q^m\theta(\mathcal{X}_{n-m}, \mathcal{X}_0) + q^m\frac{1 - \left(\frac{\alpha}{q}\right)^m}{1 - \frac{\alpha}{q}}\theta(\mathcal{X}_0, \mathcal{X}_1) \\
&\quad + \beta\frac{q}{1-q}\{1 + \alpha + \alpha^2 + \cdots + \alpha^{m-1}\}\theta(D, E) + (1-2q)\frac{1-q^m}{1-q}\theta(D, E)
\end{aligned}$$

Since $\{\mathcal{X}_n\}$ is a bounded sequence, considering the last inequality we obtain

$$\begin{aligned}
\theta(D, E) &\leq \theta(\mathcal{X}_n, \mathcal{X}_m) \\
&\leq q^m M + \frac{q^m - \alpha^m}{1 - \frac{\alpha}{q}}\theta(\mathcal{X}_0, \mathcal{X}_1) + \frac{q}{1-q}\theta(D, E) \\
&\quad + (1-2q)\frac{1-q^m}{1-q}\theta(D, E)
\end{aligned}$$

for all $n, m \in \mathbb{N}$ with $n \geq m$ and for some $M > 0$. Hence, we have

$$\lim_{n, m \rightarrow \infty} \theta(\mathcal{X}_n, \mathcal{X}_m) = \theta(D, E).$$

Now, since (D, E) is a cyclically 0-complete pair, without loss of the generality we can assume that $\{\mathcal{X}_{2n}\}$ has a subsequence $\{\mathcal{X}_{2n_i}\}$ such that

$$\lim_{i, j \rightarrow \infty} \theta(\mathcal{X}_{2n_i}, \mathcal{X}_{2n_j}) = \lim_{i \rightarrow \infty} \theta(\mathcal{X}_{2n_i}, \mathcal{X}^*) = \theta(\mathcal{X}^*, \mathcal{X}^*) = 0 \quad (7)$$

for some $\varkappa^* \in D$. Moreover, we have

$$\begin{aligned}\theta(D, E) &\leq \theta(\varkappa^*, \varkappa_{2n_i-1}) \\ &\leq \theta(\varkappa^*, \varkappa_{2n_i}) + \theta(\varkappa_{2n_i-1}, \varkappa_{2n_i}).\end{aligned}$$

Taking limit $i \rightarrow \infty$ in last inequality and using the equality (6) we have

$$\lim_{i \rightarrow \infty} \theta(\varkappa^*, \varkappa_{2n_i-1}) = \theta(D, E)$$

Additionally, we get

$$\begin{aligned}\theta(\varkappa_{2n_i}, Y\varkappa^*) &= \theta(Y\varkappa_{2n_i-1}, Y\varkappa^*) \\ &\leq q\{\theta(\varkappa_{2n_i-1}, \varkappa^*) + |\theta(\varkappa^*, Y\varkappa^*) - \theta(\varkappa_{2n_i-1}, Y\varkappa_{2n_i-1})|\} \\ &\quad + (1-q)\theta(D, E) \\ &= q\{\theta(\varkappa_{2n_i-1}, \varkappa^*) + |\theta(\varkappa^*, Y\varkappa^*) - \theta(\varkappa_{2n_i-1}, \varkappa_{2n_i})|\} \\ &\quad + (1-q)\theta(D, E).\end{aligned}$$

Taking limit $i \rightarrow \infty$ in last inequality, from (7) we get

$$\begin{aligned}\theta(\varkappa^*, Y\varkappa^*) &= \lim_{i \rightarrow \infty} \theta(\varkappa_{2n_i}, Y\varkappa^*) \\ &\leq q\theta(D, E) + q|\theta(\varkappa^*, Y\varkappa^*) - \theta(D, E)| + (1-q)\theta(D, E) \\ &= q\theta(D, E) + q\theta(\varkappa^*, Y\varkappa^*) - q\theta(D, E) + (1-q)\theta(D, E).\end{aligned}$$

Therefore, we have $\theta(\varkappa^*, Y\varkappa^*) \leq \theta(D, E)$, and so $\theta(\varkappa^*, Y\varkappa^*) = \theta(D, E)$. Hence, \varkappa^* is a best proximity point of Y in D . If $\{\varkappa_{2n+1}\}$ has a subsequence $\{\varkappa_{2n_i+1}\}$ such that

$$\lim_{i,j \rightarrow \infty} \theta(\varkappa_{2n_i+1}, \varkappa_{2n_j+1}) = \lim_{i \rightarrow \infty} \theta(\varkappa_{2n_i+1}, \zeta^*) = \theta(\zeta^*, \zeta^*) = 0$$

for some $\zeta^* \in E$. Then, by the similar way, it can be shown that ζ^* is a best proximity point of Y in E . \square

Example 3. Let $\Lambda = [0, \infty) \times [0, \infty)$ and $\theta : \Lambda \times \Lambda \rightarrow \mathbb{R}$ be a function defined by

$$\theta(\varkappa, \zeta) = \max\{\varkappa_1, \zeta_1\} + |\varkappa_2 - \zeta_2|.$$

for $\varkappa = (\varkappa_1, \varkappa_2), \zeta = (\zeta_1, \zeta_2) \in \Lambda$. It is clear that (Λ, θ) is a partial metric space. Let

$$D = \{(a, 1) : a \in [0, \infty)\},$$

and

$$E = \{(a, 2) : a \in [0, \infty)\},$$

then $\theta(D, E) = 1$. Now, we show that the pair (D, E) is a cyclically 0-complete pair. Let $\{\varkappa_n\}$ be a cyclically Cauchy sequence in $D \cup E$ with $\{\varkappa_{2n}\} \subseteq D$ and $\{\varkappa_{2n+1}\} \subseteq E$. Then, we have

$$\lim_{n,m \rightarrow \infty} \theta(\varkappa_n, \varkappa_m) = \theta(D, E).$$

Hence, we get

$$\lim_{n,m \rightarrow \infty} \max\{\varkappa_{2n}^1, \varkappa_{2m+1}^1\} = 0$$

which implies that $\lim_{n \rightarrow \infty} \varkappa_{2n}^1 = 0$ and $\lim_{m \rightarrow \infty} \varkappa_{2m+1}^1 = 0$. Then, we have

$$\lim_{n,k \rightarrow \infty} \theta(\varkappa_{2n}, \varkappa_{2k}) = \lim_{n \rightarrow \infty} \theta(\varkappa_{2n}, (0, 1)) = \theta((0, 1), (0, 1)) = 0,$$

that is, the sequence $\{\varkappa_{2n}\}$ has a subsequence satisfying (2). If we define a mapping $Y : D \cup E \rightarrow D \cup E$ by

$$Y\varkappa = \begin{cases} (\frac{a}{2}, 2) & , \quad \varkappa = (a, 1) \in D \\ (\frac{a}{2}, 1) & , \quad \varkappa = (a, 2) \in E \end{cases}$$

then, it is clear that Y is a p -cyclic contraction mapping for $q = \frac{1}{2}$. Hence, all conditions of Theorem 3 are satisfied, and so Y has a best proximity point \varkappa^* in $D \cup E$.

Corollary 1. Let $\emptyset \neq D, E$ be subsets of a partial metric space (Λ, θ) and $Y : D \cup E \rightarrow D \cup E$ is a p -cyclic contraction mapping. If D or E is 0-boundedly compact, then Y has a best proximity point.

Proof. From Remark 2, we know that if D or E is 0-boundedly compact, then the pair (D, E) is a cyclically 0-complete pair. Considering Theorem 3, we obtain that Y has a best proximity point. \square

Using Theorem 3 we obtain the following corollary which is a generalization of the main result of Popescu [5].

Corollary 2. Let (Λ, θ) be a 0-complete partial metric space and $Y : \Lambda \rightarrow \Lambda$ be a mapping. If there exists q in $[0, 1)$ such that

$$\theta(Y\varkappa, Y\zeta) \leq q\{\theta(\varkappa, \zeta) + |\theta(\varkappa, Y\varkappa) - \theta(\zeta, Y\zeta)|\} \quad (8)$$

then Y has a fixed point.

Proof. Let (Λ, θ) be a 0-complete partial metric space. If we take $D = E = \Lambda$ and $\theta(D, E) = 0$, taking into Remark 1 we can say that (D, E) is a cyclically 0-complete pair. Additionally, from inequality (8) Y is p -cyclic contraction mapping. Since all hypotheses of Theorem 3 are satisfied, we conclude that there exists a point $\varkappa^* \in \Lambda$ such that

$$\theta(\varkappa^*, Y\varkappa^*) = \theta(D, E) = 0$$

which implies that $\varkappa^* = Y\varkappa^*$. \square

4. Application

In this section, we will consider the following nonlinear Fredholm integral equation

$$u(t) = \varphi(t) + \int_0^1 K(t, s, u(s))ds \quad (9)$$

where the functions $\varphi : [0, 1] \rightarrow [0, \infty)$ and $K : [0, 1]^2 \times [0, \infty) \rightarrow [0, \infty)$ are continuous. In mathematics and other sciences such as physics, chemistry, biology, etc., some problems can be modeled by this kind of integral equations. In general, to find an exact solution to these integral equations may not be possible. Hence, it can be used the iterative methods as an alternative way to approach the solution [29–31]. We investigate the existence a solution of nonlinear Fredholm integral equations by taking into account Corollary 2. Now, we consider the space Λ as the positive cone of $C[0, 1]$, that is,

$$\Lambda = \{u \in C[0, 1] : u(t) \geq 0 \text{ for all } t \in [0, 1]\}.$$

Define a partial metric on Λ as

$$\theta(u, v) = \begin{cases} \sup_{t \in [0, 1]} u(t) & , \quad u = v \\ \sup_{t \in [0, 1]} \{u(t) + v(t)\} & , \quad u \neq v \end{cases}$$

Then, (Λ, θ) is a 0-complete partial metric space.

Theorem 4. Assume the following conditions hold:

(i) the mapping $Y : \Lambda \rightarrow \Lambda$ defined by

$$Yu(t) = \varphi(t) + \int_0^1 K(t, s, u(s)) ds$$

for all $u \in \Lambda$ and $t \in [0, 1]$ is continuous,

(ii) there is q in $[0, 1]$ such that

$$K(t, s, u(s)) + K(t, s, v(s)) \leq q \left\{ \begin{array}{c} u(s) + v(s) \\ \sup_{s \in [0, 1]} \{u(s) + Yu(s)\} \\ - \sup_{s \in [0, 1]} \{v(s) + Yv(s)\} \end{array} \right\} - 2\varphi(t)$$

for all $t, s \in [0, 1]$ and $u, v \in \Lambda$.

Then, the integral Equation (9) has a positive solution.

Proof. If we prove that Y has a fixed point, we show that Equation (9) has a solution. Now, for all $t \in [0, 1]$ and $u, v \in \Lambda$, we have

$$\begin{aligned} Yu(t) + Yv(t) &= \varphi(t) + \int_0^1 K(t, s, u(s)) ds + \varphi(t) + \int_0^1 K(t, s, v(s)) ds \\ &= 2\varphi(t) + \int_0^1 \{K(t, s, u(s)) + K(t, s, v(s))\} ds \\ &\leq 2\varphi(t) + \int_0^1 \left\{ q \left\{ \begin{array}{c} u(s) + v(s) \\ \sup_{s \in [0, 1]} \{u(s) + Yu(s)\} \\ - \sup_{s \in [0, 1]} \{v(s) + Yv(s)\} \end{array} \right\} - 2\varphi(t) \right\} ds \\ &= 2\varphi(t) + q \int_0^1 \sup_{s \in [0, 1]} \{u(s) + v(s)\} ds \\ &\quad + q \int_0^1 \left| \sup_{s \in [0, 1]} \{u(s) + Yu(s)\} - \sup_{s \in [0, 1]} \{v(s) + Yv(s)\} \right| ds \\ &\quad - 2\varphi(t) \int_0^1 ds \\ &= q \sup_{s \in [0, 1]} \{u(s) + v(s)\} \\ &\quad + q \left| \sup_{s \in [0, 1]} \{u(s) + Yu(s)\} - \sup_{s \in [0, 1]} \{v(s) + Yv(s)\} \right| \\ &= q \{\theta(u, v) + |\theta(u, Yu) - \theta(v, Yv)|\}. \end{aligned}$$

This implies that

$$\sup_{t \in [0, 1]} \{Yu(t) + Yv(t)\} \leq q \{\theta(u, v) + |\theta(u, Yu) - \theta(v, Yv)|\},$$

and so we have

$$\theta(Yu, Yv) \leq q \{\theta(u, v) + |\theta(u, Yu) - \theta(v, Yv)|\}.$$

Therefore, all conditions of Corollary 2 hold, and so the integral Equation (9) has a positive solution. \square

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: The author is thankful to the referees and editor for making valuable suggestions leading to the better presentations of the paper.

Conflicts of Interest: The author declares no conflict of interest.

References

- Banach, S. Sur les opérations dans les ensembles abstraits et leur applications aux équations intégrales. *Fund. Math.* **1922**, *3*, 133–181. [\[CrossRef\]](#)
- Kadelburg, Z.; Radenovic, S. Fixed point and tripled fixed point theorems under Pata-type conditions in ordered metric spaces. *Int. J. Anal. Appl.* **2014**, *6*, 113–122.
- Özeken, Ç.C.; Çevik, C. Ordered vectorial quasi and almost contractions on ordered vector metric spaces. *Mathematics* **2021**, *9*, 2443. [\[CrossRef\]](#)
- Reich, S. Fixed points of contractive functions. *Boll. Unione Mat. Ital.* **1972**, *5*, 26–42.
- Popescu, O. A new type of contractive mappings in complete metric spaces. *Bull. Transilv. Univ. Braşov Math. Inform. Phys.* **2008**, *1*, 479–482.
- Kirk, W.A.; Srinivasan, P.S.; Veeramani, P. Fixed points for mappings satisfying cyclical contractive conditions. *Fixed Point Theory* **2003**, *4*, 79–89.
- Karapinar, E. Best proximity points of cyclic mappings. *Appl. Math. Lett.* **2012**, *25*, 1761–1766. [\[CrossRef\]](#)
- Suzuki, T.; Kikkawa, M.; Vetro, C. The existence of best proximity points in metric spaces with the property UC. *Nonlinear Anal.* **2009**, *71*, 2918–2926. [\[CrossRef\]](#)
- Basha, S.S. Extensions of Banach’s contraction principle. *Numer. Funct. Anal. Optim.* **2010**, *31*, 569–576. [\[CrossRef\]](#)
- Espinola, R.; Kosuru, G.S.R.; Veeramani, P. Pythagorean property and best proximity theorems. *J. Optim. Theory Appl.* **2015**, *164*, 534–550. [\[CrossRef\]](#)
- Kim, W.K.; Lee, K.H. Existence of best proximity pairs and equilibrium pairs. *J. Math. Anal. Appl.* **2006**, *316*, 433–446. [\[CrossRef\]](#)
- Raj, V.S. A best proximity point theorem for weakly contractive non-self-mappings. *Nonlinear Anal. TMA* **2011**, *74*, 4804–4808.
- Reich, S. Approximate selections, best approximations, fixed points, and invariant sets. *J. Math. Anal. Appl.* **1978**, *62*, 104–113. [\[CrossRef\]](#)
- Aslantas, M. Some best proximity point results via a new family of F -contraction and an application to homotopy theory. *J. Fixed Point Theory Appl.* **2021**, *23*, 54. [\[CrossRef\]](#)
- Sahin, H.; Aslantas, M.; Altun, I. Best proximity and best periodic points for proximal nonunique contractions. *J. Fixed Point Theory Appl.* **2021**, *23*, 55. [\[CrossRef\]](#)
- Eldred, A.A.; Veeramani, P. Existence and convergence of best proximity points. *J. Math. Anal. Appl.* **2006**, *323*, 1001–1006. [\[CrossRef\]](#)
- Karpagam, S.; Agrawal, S. Best proximity points for cyclic contractions. 2012, *preprint*.
- Abkar, A.; Moezzifar, N.; Azizi, A.; Shahzad, N. Best proximity point theorems for cyclic generalized proximal contractions. *Fixed Point Theory Appl.* **2016**, *2016*, 66. [\[CrossRef\]](#)
- Basha, S.S.; Shahzad, N.; Vetro, C. Best proximity point theorems for proximal cyclic contractions. *J. Fixed Point Theory Appl.* **2017**, *19*, 2647–2661. [\[CrossRef\]](#)
- Matthews, S.G. *Partial Metric Topology*; Annals of the New York Academy of Sciences: New York, NY, USA, 1994; pp. 183–197.
- Aslantas, M.; Abed, A.Z.D.R. Some best proximity point results for multivalued mappings on partial metric spaces. *Math. Moravica* **2021**, *25*, 99–111. [\[CrossRef\]](#)
- Čirić, L.; Samet, B.; Aydi, H.; Vetro, C. Common fixed points of generalized contractions on partial metric spaces and an application. *Appl. Math. Comput.* **2011**, *218*, 2398–2406. [\[CrossRef\]](#)
- Karapinar, E. Generalizations of Caristi Kirk’s theorem on partial metric spaces. *Fixed Point Theory Appl.* **2011**, *2011*, 4. [\[CrossRef\]](#)
- Romaguera, S. A Kirk type characterization of completeness for partial metric spaces. *Fixed Point Theory Appl.* **2010**, *2009*, 493298. [\[CrossRef\]](#)
- Abbas, M.; Nazir, T. Fixed point of generalized weakly contractive mappings in ordered partial metric spaces. *Fixed Point Theory Appl.* **2012**, *2012*, 1. [\[CrossRef\]](#)
- Altun, I.; Sola, F.; Simsek, H. Generalized contractions on partial metric spaces. *Topol. Its Appl.* **2010**, *157*, 2778–2785. [\[CrossRef\]](#)
- Altun, I.; Simsek, H. Some fixed point theorems on dualistic partial metric spaces. *J. Adv. Math. Stud.* **2008**, *1*, 1–9.
- Haghi, R.H.; Rezapour, S.; Shahzad, N. Be careful on partial metric fixed point results. *Topol. Appl.* **2013**, *160*, 450–454. [\[CrossRef\]](#)
- Sidorov, N.A.; Sidorov, D.N. Solving the Hammerstein integral equation in the irregular case by successive approximations. *Sib. Math. J.* **2010**, *51*, 325–329. [\[CrossRef\]](#)
- Sidorov, D.N. Existence and blow-up of Kantorovich principal continuous solutions of nonlinear integral equations. *Diff. Equat.* **2014**, *50*, 1217–1224. [\[CrossRef\]](#)
- Trenogin, V.A. Locally Invertible Operators and the Method of Continuation with Respect to Parameter. *Funct. Anal. Appl.* **1996**, *30*, 147–148. [\[CrossRef\]](#)