

## Article

# Multiple Periodic Solutions for Odd Perturbations of the Discrete Relativistic Operator

Petru Jebelean \* and Călin Șerban

Department of Mathematics, West University of Timișoara, 4, Blvd. V. Pârvan, 300223 Timișoara, Romania; calin.serban@e-uvt.ro

\* Correspondence: petru.jebelean@e-uvt.ro

**Abstract:** We obtain the existence of multiple pairs of periodic solutions for difference equations of type  $-\Delta\left(\frac{\Delta u(n-1)}{\sqrt{1-|\Delta u(n-1)|^2}}\right) = \lambda g(u(n))$  ( $n \in \mathbb{Z}$ ), where  $g : \mathbb{R} \rightarrow \mathbb{R}$  is a continuous odd function with antiovercoercive primitive, and  $\lambda > 0$  is a real parameter. The approach is variational and relies on the critical point theory for convex, lower semicontinuous perturbations of  $C^1$ -functionals.

**Keywords:** discrete relativistic operator; periodic solution; critical point; genus

**MSC:** 39A23; 39A27; 47J20

## 1. Introduction

In this note, we are concerned with the multiplicity of solutions for difference equations with relativistic operator of type

$$-\Delta[\phi(\Delta u(n-1))] = \lambda g(u(n)), \quad u(n) = u(n+T) \quad (n \in \mathbb{Z}), \quad (1)$$

where  $\Delta u(n) = u(n+1) - u(n)$  is the usual forward difference operator,  $\lambda > 0$  is a real parameter,  $g : \mathbb{R} \rightarrow \mathbb{R}$  is a continuous odd function, and

$$\phi(y) = \frac{y}{\sqrt{1-y^2}} \quad (y \in (-1, 1)).$$

In recent years, special attention has been paid to the existence and multiplicity of  $T$ -periodic solutions for problems with a discrete relativistic operator. Thus, for instance, in [1,2], variational arguments were employed to prove the solvability of systems of difference equations having the form

$$\Delta[\phi_N(\Delta u(n-1))] = \nabla_u V(n, u(n)) + h(n) \quad (n \in \mathbb{Z}), \quad (2)$$

under various hypotheses upon  $V$  and  $h$  (coerciveness, growth restriction, convexity or periodicity conditions); here,  $\phi_N$  is the  $N$ -dimensional variant of  $\phi$ , i.e.,

$$\phi_N(y) = \frac{y}{\sqrt{1-|y|^2}} \quad (y \in \mathbb{R}^N, |y| < 1).$$

The existence of at least  $N+1$  geometrically distinct  $T$ -periodic solutions of (2) was proved in [3], under the assumptions that  $h$  is  $T$ -periodic,  $\sum_{j=1}^T h(j) = 0$ , and the mapping  $V(n, x)$  is  $T$ -periodic in  $n$  and  $\omega_i$ -periodic ( $\omega_i > 0$ ) with respect to each  $x_i$  ( $i = 1, \dots, N$ ). For the proof, using an idea from the differential case [4], the singular problem (2) was reduced to an equivalent non-singular one to which classical Ljusternik–Schnirelmann category methods can be applied. In addition, under some similar assumptions on  $V$  and  $h$ ,



**Citation:** Jebelean, P.; Șerban, C. Multiple Periodic Solutions for Odd Perturbations of the Discrete Relativistic Operator. *Mathematics* **2022**, *10*, 1595. <https://doi.org/10.3390/math10091595>

Academic Editor: Alessio Pomponio

Received: 2 April 2022

Accepted: 7 May 2022

Published: 8 May 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

were obtained in [5] using Morse theory, conditions under which system (2) has at least  $2^N$  geometrically distinct  $T$ -periodic solutions.

The motivation of the present study mainly comes from paper [6], where for problems involving Fisher-Kolmogorov nonlinearities of type

$$-\Delta[\phi(\Delta u(n-1))] = \lambda u(n)(1 - |u(n)|^q), \quad u(n) = u(n+T) \quad (n \in \mathbb{Z}), \quad (3)$$

with  $q > 0$  fixed and  $\lambda > 0$  a real parameter, it was proved that if  $\lambda > 8mT$  for some  $m \in \mathbb{N}$  with  $2 \leq m \leq T$ , then problem (3) has at least  $m$  distinct pairs of nontrivial solutions. We also refer the interested reader to [6] for a discussion concerning the origin and steps in the study of this type of nonlinearity. In this respect, we shall see in Example 1 below that a sharper result holds true, namely,

- (i) If  $\lambda > 8 \sin^2 \frac{m\pi}{T}$  with  $0 \leq m \leq \begin{cases} (T-1)/2 & \text{if } T \text{ is odd} \\ (T-2)/2 & \text{if } T \text{ is even} \end{cases}$ , then problem (3) has at least  $2m+1$  distinct pairs of nontrivial solutions.
- (ii) If  $T$  is even and  $\lambda > 8$ , then (3) has at least  $T$  distinct pairs of nontrivial solutions.

Moreover, we prove in Theorem 2 that the above statements (i) and (ii) still remain valid for a larger class of periodic problems.

As in [6], our approach to problem (1) is variational and combines a Clark-type abstract result for convex, lower semicontinuous perturbations of  $C^1$ -functionals, based on Krasnoselskii's genus. However, our technique here brings the novelty that it exploits the interference of the geometry of the energy functional with fine spectral properties of the operator  $-\Delta^2$ ; recall that

$$\Delta^2 u(n-1) := \Delta(\Delta u(n-1)) = u(n+1) - 2u(n) + u(n-1).$$

It is worth noting that in paper [7] analogous multiplicity results are obtained in the differential case for potential systems involving parametric odd perturbations of the relativistic operator. In addition, we mention the recent paper [8], where the authors obtain the existence and multiplicity of sign-changing solutions for a slightly modified parametric problem of type (1) using bifurcation techniques.

We conclude this introductory part by briefly recalling some topics in the frame of Szulkin's critical point theory [9], which is needed in the sequel. Let  $(Y, \|\cdot\|)$  be a real Banach space and  $\mathcal{I} : Y \rightarrow (-\infty, +\infty]$  be a functional having the following structure:

$$\mathcal{I} = \mathcal{F} + \psi, \quad (4)$$

where  $\mathcal{F} \in C^1(Y, \mathbb{R})$  and  $\psi : Y \rightarrow (-\infty, +\infty]$  is proper, convex and lower semicontinuous. A point  $u \in D(\psi)$  is said to be a *critical point* of  $\mathcal{I}$  if it satisfies the inequality

$$\langle \mathcal{F}'(u), v - u \rangle + \psi(v) - \psi(u) \geq 0 \quad \forall v \in D(\psi).$$

A sequence  $\{u_n\} \subset D(\psi)$  is called a (PS)-sequence if  $\mathcal{I}(u_n) \rightarrow c \in \mathbb{R}$  and

$$\langle \mathcal{F}'(u_n), v - u_n \rangle + \psi(v) - \psi(u_n) \geq -\varepsilon_n \|v - u_n\| \quad \forall v \in D(\psi),$$

where  $\varepsilon_n \rightarrow 0$ . The functional  $\mathcal{I}$  is said to *satisfy the (PS) condition* if any (PS)-sequence has a convergent subsequence in  $Y$ .

Let  $\Sigma$  be the collection of all symmetric subsets of  $Y \setminus \{0\}$  which are closed in  $Y$ . The *genus* of a nonempty set  $A \in \Sigma$  is defined as being the smallest integer  $k$  with the property that there exists an odd continuous mapping  $h : A \rightarrow \mathbb{R}^k \setminus \{0\}$ ; in this case, we write  $\gamma(A) = k$ . If such an integer does not exist, then  $\gamma(A) := +\infty$ . Notice that if  $A \in \Sigma$  is homeomorphic to  $S^{k-1}$  ( $k-1$  dimension unit sphere in the Euclidean space  $\mathbb{R}^k$ ) by an odd homeomorphism, then  $\gamma(A) = k$  ([10], Corollary 5.5). For other properties and more details on the notion of genus, we refer the reader to [10,11]. The following theorem is an immediate consequence of ([9], Theorem 4.3).

**Theorem 1.** Let  $\mathcal{I}$  be of type (4) with  $\mathcal{F}$  and  $\psi$  even. In addition, suppose that  $\mathcal{I}$  is bounded from below, satisfies the (PS) condition and  $\mathcal{I}(0) = 0$ . If there exists a nonempty compact symmetric subset  $A \subset Y \setminus \{0\}$  with  $\gamma(A) \geq k$ , such that

$$\sup_{v \in A} \mathcal{I}(v) < 0,$$

then the functional  $\mathcal{I}$  has at least  $k$  distinct pairs of nontrivial critical points.

## 2. Variational Approach and Preliminaries

To introduce the variational formulation for problem (1), let  $H_T$  be the space of all  $T$ -periodic  $\mathbb{Z}$ -sequences in  $\mathbb{R}$ , i.e., of mappings  $u : \mathbb{Z} \rightarrow \mathbb{R}$ , such that  $u(n) = u(n + T)$  for all  $n \in \mathbb{Z}$ . On  $H_T$ , we consider the following inner product and corresponding norm:

$$(u|v) := \sum_{j=1}^T u(j)v(j), \quad \|u\| = \left( \sum_{j=1}^T |u(j)|^2 \right)^{1/2},$$

which makes it a Hilbert space. In addition, for each  $u \in H_T$ , we set

$$\bar{u} := \frac{1}{T} \sum_{j=1}^T u(j), \quad \tilde{u} := u - \bar{u}.$$

It is not difficult to check that

$$|\tilde{u}(i)| \leq T^{1/2} \left( \sum_{j=1}^T |\Delta u(j)|^2 \right)^{1/2} \quad (i \in \{1, \dots, T\}). \quad (5)$$

Now, let the closed convex subset  $K$  of  $H_T$  be defined by

$$K := \{u \in H_T : |\Delta u|_\infty \leq 1\},$$

where  $|\Delta u|_\infty := \max_{i=1, \dots, T} |\Delta u(i)|$ . Then, from (5), one has

$$|\bar{u}| - T \leq |u(i)| \leq |\bar{u}| + T \quad (i \in \{1, \dots, T\}), \quad (6)$$

for all  $u \in K$ . We introduce the even functions

$$\Psi(u) = \begin{cases} \sum_{j=1}^T \Phi[\Delta u(j)], & \text{if } u \in K, \\ +\infty, & \text{otherwise,} \end{cases}$$

where  $\Phi(y) = 1 - \sqrt{1 - y^2}$  ( $y \in [-1, 1]$ ) and

$$\mathcal{G}_\lambda(u) = -\lambda \sum_{j=1}^T G(u(j)) \quad (u \in H_T),$$

with  $G$  the primitive

$$G(x) = \int_0^x g(\tau) d\tau \quad (x \in \mathbb{R}).$$

It is not difficult to see that  $\Psi$  is convex and lower semicontinuous, while  $\mathcal{G}_\lambda$  is of class  $C^1$ , its derivative being given by

$$\langle \mathcal{G}'_\lambda(u), v \rangle = -\lambda \sum_{j=1}^T g(u(j))v(j) \quad (u, v \in H_T).$$

Then, the functional  $I_\lambda : H_T \rightarrow (-\infty, +\infty]$  associated to (1) is

$$I_\lambda = \Psi + \mathcal{G}_\lambda$$

and it is clear that it has the structure required by Szulkin's critical point theory. A solution of problem (1) is an element  $u \in H_T$  such that  $|\Delta u(n)| < 1$ , for all  $n \in \mathbb{Z}$ , which satisfies the equation in (1). The following result reduces the search of solutions of problem (1) to finding critical points of  $I_\lambda$ .

**Proposition 1.** Any critical point of  $I_\lambda$  is a solution of problem (1).

**Proof.** Let  $e \in H_T$ . By virtue of Lemmas 5 and 6 in [1], the problem

$$\Delta[\phi(\Delta u(n-1))] = \bar{u} + e(n), \quad u(n) = u(n+T) \quad (n \in \mathbb{Z})$$

has a unique solution  $u_e$ , which is also the unique solution of the variational inequality

$$\sum_{j=1}^T \{\Phi[\Delta v(j)] - \Phi[\Delta u(j)] + \bar{u}(\bar{v} - \bar{u}) + e(j)(v(j) - u(j))\} \geq 0, \quad \forall v \in K \quad (7)$$

([6], Proposition 3.1). Next, let  $w \in K$  be a critical point of  $I_\lambda$ . Then, for any  $v \in K$ , one has

$$\sum_{j=1}^T \{\Phi[\Delta v(j)] - \Phi[\Delta w(j)] - \lambda g(w(j))(v(j) - w(j))\} \geq 0,$$

which can be written as

$$\sum_{j=1}^T \{\Phi[\Delta v(j)] - \Phi[\Delta w(j)] + \bar{w}(v(j) - w(j))\} - \sum_{j=1}^T [\lambda g(w(j)) + \bar{w}](v(j) - w(j)) \geq 0.$$

Hence,  $w$  is a solution of the variational inequality

$$\sum_{j=1}^T \{\Phi[\Delta v(j)] - \Phi[\Delta w(j)] + \bar{w}(\bar{v} - \bar{w}) + e_w(j)(v(j) - w(j))\} \geq 0, \quad \forall v \in K, \quad (8)$$

with  $e_w \in H_T$  being given by  $e_w(n) = -\lambda g(w(n)) - \bar{w}$  ( $n \in \mathbb{Z}$ ).

Therefore, by (8) and the uniqueness of the solution of (7), we obtain that, in fact,  $w$  solves problem (1).  $\square$

**Proposition 2.** If  $G$  is anticoercive, i.e.,

$$\lim_{|x| \rightarrow +\infty} G(x) = -\infty, \quad (9)$$

then  $I_\lambda$  is bounded from below and satisfies the (PS) condition.

**Proof.** From (9) we have that  $-G$ , hence  $\mathcal{G}_\lambda$ , are bounded from below on  $\mathbb{R}$ , respectively on  $H_T$ . This, together with the fact that  $\Psi$  is bounded from below, ensure that the same is true for  $I_\lambda$ .

To see that  $I_\lambda$  satisfies the (PS) condition, let  $\{u_n\} \subset K$  be a (PS)-sequence. Assuming by contradiction that  $\{|\bar{u}_n|\}$  is not bounded, we may suppose, going, if necessary, to a subsequence, that  $|\bar{u}_n| \rightarrow +\infty$ . Then, by virtue of (6) and (9), we deduce that  $I_\lambda(u_n) \rightarrow -\infty$ , contradicting the fact that  $\{I_\lambda(u_n)\}$  is convergent. Consequently,  $\{|\bar{u}_n|\}$  is bounded. This, together with  $|\bar{u}_n| \leq T$  shows that  $\{u_n\}$  is bounded in the finite-dimensional space  $H_T$ ; hence, it contains a convergent subsequence.  $\square$

**Remark 1.** Notice that until here in this section, no parity assumptions on the continuous function  $g : \mathbb{R} \rightarrow \mathbb{R}$  must be required.

We end this section by reviewing some spectral properties of the operator  $-\Delta^2$ , which is needed in the sequel. A real number  $\lambda \in \mathbb{R}$  is said to be an *eigenvalue* of  $-\Delta^2$  on  $H_T$ , if there is some  $u \in H_T \setminus \{0_{H_T}\}$  such that

$$-\Delta^2 u(n-1) = \lambda u(n), \quad (n \in \mathbb{Z}) \quad (10)$$

and in this case,  $u$  is called *eigensequence* corresponding to the eigenvalue  $\lambda$ . On account of the periodicity of  $u$ , relation (10) is equivalent to the system

$$\begin{cases} -u(2) + 2u(1) - u(T) = \lambda u(1) \\ -u(3) + 2u(2) - u(1) = \lambda u(2) \\ \vdots \\ -u(T) + 2u(T-1) - u(T-2) = \lambda u(T-1) \\ -u(1) + 2u(T) - u(T-1) = \lambda u(T). \end{cases} \quad (11)$$

If we consider the particular circulant matrix

$$M_T := \begin{pmatrix} 2 & -1 & 0 & \cdots & 0 & 0 & -1 \\ -1 & 2 & -1 & \cdots & 0 & 0 & 0 \\ \vdots & & & & & & \\ 0 & 0 & 0 & \cdots & -1 & 2 & -1 \\ -1 & 0 & 0 & \cdots & 0 & -1 & 2 \end{pmatrix}$$

then, having in view (11), the eigenvalues of  $-\Delta^2$  are precisely the characteristic roots of  $M_T$ . In addition, if  $y = (y_1, \dots, y_T) \in \mathbb{R}^T \setminus \{0_{\mathbb{R}^T}\}$  is an eigenvector corresponding to a characteristic root  $\lambda$ , then its *extension*  $u^y \in H_T$ , defined by  $u^y(i) = y_i$  for  $i = \overline{1, T}$ , is an eigensequence corresponding to the eigenvalue  $\lambda$ . This means that an orthonormal basis of eigensequences  $u^1, \dots, u^T$  can be constructed from an orthonormal basis of eigenvectors  $x^1, \dots, x^T$  of  $M_T$  by extending  $x^i$  in  $H_T$  ( $i = \overline{1, T}$ ) as above.

From ([12], p. 38), we know that the characteristic roots of  $M_T$ , hence the eigenvalues of  $-\Delta^2$ , are  $4 \sin^2 i\pi/T$  ( $i = \overline{0, T-1}$ ). We can label them according to the parity of  $T$  as follows:

Todd :

$$\lambda_0 = 0, \quad \lambda_{2k-1} = \lambda_{2k} = 4 \sin^2 \frac{k\pi}{T}, \quad k = 1, \dots, \frac{T-1}{2};$$

Even :

$$\lambda_0 = 0, \quad \lambda_{2k-1} = \lambda_{2k} = 4 \sin^2 \frac{k\pi}{T}, \quad k = 1, \dots, \frac{T-2}{2}, \quad \lambda_{T-1} = 4.$$

In both cases, we consider an orthonormal basis  $e^0, \dots, e^{T-1}$  in  $H_T$ , such that  $e^i$  is an eigensequence corresponding to  $\lambda_i$  ( $i = \overline{0, T-1}$ ). Observe that, by multiplying equality (10) by arbitrary  $v \in H_T$  and using summation by parts formula, one obtains that if  $u \in H_T$  and  $\lambda \in \mathbb{R}$  satisfy (10), then

$$\sum_{j=1}^T \Delta u(j) \Delta v(j) = \lambda(u|v).$$

This yields

$$\sum_{j=1}^T \Delta e^i(j) \Delta e^k(j) = \lambda_k \delta_{ik} \quad (i, k \in \{0, \dots, T-1\}), \quad (12)$$

where  $\delta_{ik}$  stands for the Kronecker delta function.

### 3. Main Result

Our main result is given in the following.

**Theorem 2.** Assume that  $g : \mathbb{R} \rightarrow \mathbb{R}$  is a continuous odd function and that  $G$  satisfies (9) together with

$$\liminf_{x \rightarrow 0} \frac{2G(x)}{x^2} \geq 1. \quad (13)$$

Then, the following hold true:

(i) If

$$\lambda > 8 \sin^2 \frac{m\pi}{T} (= 2\lambda_{2m}) \text{ with } 0 \leq m \leq \begin{cases} (T-1)/2 & \text{if } T \text{ is odd} \\ (T-2)/2 & \text{if } T \text{ is even} \end{cases}, \quad (14)$$

then problem (1) has at least  $2m+1$  distinct pairs of nontrivial solutions.

(ii) If  $T$  is even and

$$\lambda > 8 (= 2\lambda_{T-1}), \quad (15)$$

then (1) has at least  $T$  distinct pairs of nontrivial solutions.

**Proof.** We show (i) in the odd case because the even case follows by exactly the same arguments, and under assumption (15), a quite similar strategy works by simply replacing “ $2m$ ” with “ $T-1$ ”.

Thus, let  $0 \leq m \leq (T-1)/2$ . On account of Theorem 1 and Propositions 1 and 2, we have to prove that there exists a nonempty compact symmetric subset  $A_m \subset H_T \setminus \{0\}$  with  $\gamma(A_m) \geq 2m+1$ , such that

$$\sup_{v \in A_m} I_\lambda(v) < 0. \quad (16)$$

Since  $\lambda > 2\lambda_{2m}$ , we can choose  $\varepsilon \in (0, 1)$ , so that  $\lambda > 2\lambda_{2m}/(1-\varepsilon)$ . Then, by virtue of (13), there exists  $\delta > 0$  such that

$$2G(x) \geq (1-\varepsilon)x^2 \quad \text{as } |x| \leq \delta. \quad (17)$$

Next, we introduce the set

$$A_m := \left\{ \sum_{k=0}^{2m} \alpha_k e^k : \alpha_0^2 + \cdots + \alpha_{2m}^2 = \rho^2 \right\},$$

where  $\rho$  is a positive number, which is chosen  $\leq \min \left\{ \frac{1}{2\sqrt{2m+1}}, \delta \right\}$ .

Then, it is not difficult to see that the odd mapping  $H : A_m \rightarrow S^{2m}$  defined by

$$H \left( \sum_{k=0}^{2m} \alpha_k e^k \right) = \left( \frac{\alpha_0}{\rho}, \frac{\alpha_1}{\rho}, \dots, \frac{\alpha_{2m}}{\rho} \right)$$

is a homeomorphism between  $A_m$  and  $S^{2m}$ ; therefore,  $\gamma(A_m) = 2m+1$ .

We have that  $A_m \subset K$ . Indeed, let  $v = \sum_{k=0}^{2m} \alpha_k e^k \in A_m$ . Then, for all  $j \in \{1, \dots, T\}$ , we obtain

$$\begin{aligned} |\Delta v(j)| &\leq \sum_{k=0}^{2m} |\alpha_k e^k(j+1)| + \sum_{k=0}^{2m} |\alpha_k e^k(j)| \leq 2 \sum_{k=0}^{2m} |\alpha_k| \\ &\leq 2\sqrt{2m+1} \left( \sum_{k=0}^{2m} \alpha_k^2 \right)^{1/2} = 2\rho\sqrt{2m+1} \end{aligned} \quad (18)$$

and since  $\rho \leq 1/(2\sqrt{2m+1})$ , one has  $|\Delta v|_\infty \leq 1$ , which shows that  $v \in K$ . On the other hand, using (12), we obtain

$$\begin{aligned} \sum_{j=1}^T |\Delta v(j)|^2 &= \sum_{j=1}^T \left| \Delta \left( \sum_{k=0}^{2m} \alpha_k e^k(j) \right) \right|^2 = \sum_{j=1}^T \left( \sum_{k=0}^{2m} \alpha_k \Delta e^k(j) \right)^2 \\ &= \sum_{j=1}^T \left( \sum_{k=0}^{2m} \alpha_k^2 (\Delta e^k(j))^2 + \sum_{\substack{i,k=0 \\ i \neq k}}^{2m} \alpha_i \alpha_k \Delta e^k(j) \Delta e^i(j) \right) \\ &= \sum_{k=0}^{2m} \alpha_k^2 \sum_{j=1}^T (\Delta e^k(j))^2 + \sum_{\substack{i,k=0 \\ i \neq k}}^{2m} \alpha_i \alpha_k \sum_{j=1}^T \Delta e^k(j) \Delta e^i(j) \\ &= \sum_{k=0}^{2m} \lambda_k \alpha_k^2 \leq \lambda_{2m} \sum_{k=0}^{2m} \alpha_k^2 = \lambda_{2m} \rho^2. \end{aligned} \quad (19)$$

In addition, it is clear that

$$\sum_{j=1}^T |v(j)|^2 = \|v\|^2 = (v|v) = \sum_{k=0}^{2m} \alpha_k^2 = \rho^2. \quad (20)$$

Then, from (17), (19), (20) and  $|v(j)| \leq \rho \leq \delta$  ( $j \in \{1, \dots, T\}$ ), it follows that

$$\begin{aligned} I_\lambda(v) &= \Psi(v) + \mathcal{G}_\lambda(v) \leq \sum_{j=1}^T |\Delta v(j)|^2 - \frac{\lambda}{2}(1-\varepsilon) \sum_{j=1}^T |v(j)|^2 \\ &\leq \rho^2 \lambda_{2m} - \frac{\lambda}{2}(1-\varepsilon) \rho^2 = \rho^2 \frac{2\lambda_{2m} - \lambda(1-\varepsilon)}{2} < 0. \end{aligned}$$

Therefore, (16) holds true and the proof of (i) is complete.  $\square$

**Example 1.** If (14) holds true, then problem (3) has at least  $2m+1$  distinct pairs of nontrivial solutions. In addition, if  $T$  is even, under assumption (15), problem (3) has at least  $T$  distinct pairs of nontrivial solutions. Notice that besides the trivial solution, problem (3) always has the pair of constant solutions  $u \equiv \pm 1$ , and these are the only constant nontrivial solutions of (3). Therefore, problem (3) has at least  $2m$  (resp.  $T-1$ ) distinct pairs of nonconstant solutions if hypothesis (14) is satisfied (resp. (15) holds true).

Consider the eigenvalue type problem

$$-\Delta[\phi(\Delta u(n-1))] = \lambda u(n) + h(u(n)), \quad u(n) = u(n+T) \quad (n \in \mathbb{Z}) \quad (21)$$

and set  $H(x) = \int_0^x h(\tau) d\tau$  ( $x \in \mathbb{R}$ ).

**Corollary 1.** If the continuous function  $h : \mathbb{R} \rightarrow \mathbb{R}$  is odd and

$$\liminf_{x \rightarrow 0} \frac{H(x)}{x^2} \geq 0, \quad \lim_{x \rightarrow +\infty} \frac{H(x)}{x^2} = -\infty,$$

then the conclusions (i) and (ii) of Theorem 2 remain valid with (21) instead of (1).

**Proof.** Theorem 2 applies to the problem

$$-\Delta[\phi(\Delta u(n-1))] = \lambda \left( u(n) + \frac{h(u(n))}{\lambda} \right), \quad u(n) = u(n+T) \quad (n \in \mathbb{Z}).$$

□

Theorem 2 can be employed to derive the multiplicity of nontrivial solutions of autonomous non-parametric problems having the form

$$-\Delta[\phi(\Delta u(n-1))] = f(u(n)), \quad u(n) = u(n+T) \quad (n \in \mathbb{Z}). \quad (22)$$

Setting  $F(x) = \int_0^x f(\tau) d\tau$  ( $x \in \mathbb{R}$ ), we have the following.

**Corollary 2.** Assume that  $f : \mathbb{R} \rightarrow \mathbb{R}$  is a continuous odd function and that

$$\lim_{x \rightarrow +\infty} F(x) = -\infty. \quad (23)$$

Then, the following hold true:

(i) If

$$\liminf_{x \rightarrow 0} \frac{F(x)}{x^2} > 4 \sin^2 \frac{m\pi}{T} \text{ with } 0 \leq m \leq \begin{cases} (T-1)/2 & \text{if } T \text{ is odd} \\ (T-2)/2 & \text{if } T \text{ is even} \end{cases}, \quad (24)$$

then problem (22) has at least  $2m+1$  distinct pairs of nontrivial solutions.

(ii) If  $T$  is even and

$$\liminf_{x \rightarrow 0} \frac{F(x)}{x^2} > 4, \quad (25)$$

then (22) has at least  $T$  distinct pairs of nontrivial solutions.

**Proof.** From (24), there exists  $\bar{\lambda} > 0$  such that

$$\liminf_{x \rightarrow 0} \frac{2F(x)}{x^2} \geq \bar{\lambda} > 8 \sin^2 \frac{m\pi}{T}$$

and the result follows from Theorem 2 with  $g(x) = f(x)/\bar{\lambda}$ ; a similar argument works when (25) is fulfilled. □

**Example 2.** Let  $f_a : \mathbb{R} \rightarrow \mathbb{R}$  be given by

$$f_a(x) = 2x \sin |x|^{-\frac{1}{2}} - \frac{x|x|^{-\frac{1}{2}} \cos |x|^{-\frac{1}{2}}}{2} + 2ax - 4x^3 \quad (x \in \mathbb{R}).$$

Then,

$$F_a(x) = x^2 \left( \sin |x|^{-\frac{1}{2}} + a - x^2 \right) \quad (x \in \mathbb{R})$$

and by Corollary 2, we obtain that, if

$$a > 1 + 4 \sin^2 \frac{m\pi}{T} \text{ with } 0 \leq m \leq \begin{cases} (T-1)/2 & \text{if } T \text{ is odd} \\ (T-2)/2 & \text{if } T \text{ is even} \end{cases},$$

then the equation

$$-\Delta[\phi(\Delta u(n-1))] = f_a(u(n)) \quad (n \in \mathbb{Z}) \quad (26)$$

has at least  $2m+1$  distinct pairs of nontrivial  $T$ -periodic solutions, while if  $T$  is even and  $a > 5$ , then (26) has at least  $T$  distinct pairs of nontrivial  $T$ -periodic solutions.

**Remark 2.** A multiplicity result for odd perturbations of the discrete  $p$ -Laplacian operator is obtained in [13] using a Clark-type result in the frame of the classical critical point theory.

**Author Contributions:** Writing—original draft, P.J. and C.Ş.; Writing—review & editing, P.J. and C.Ş. All authors contributed equally to the manuscript. All authors have read and agreed to the published version of the manuscript.



**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Acknowledgments:** The authors are grateful to the anonymous referees for the very useful comments and suggestions, which helped them to improve the presentation of the paper.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Mawhin, J. Periodic solutions of second order nonlinear difference systems with  $\phi$ -Laplacian: A variational approach. *Nonlinear Anal.* **2012**, *45*, 4672–4687.
2. Mawhin, J. Periodic solutions of second order Lagrangian difference systems with bounded or singular  $\phi$ -Laplacian and periodic potential. *Discrete Contin. Dyn. Syst. Ser. S* **2013**, *6*, 1065–1076.
3. Mawhin, J. A simple proof of multiplicity for periodic solutions of Lagrangian difference systems with relativistic operator and periodic potential. *J. Difference Equ. Appl.* **2016**, *22*, 306–315.
4. Jebelean, P.; Mawhin, J.; Şerban, C. Multiple periodic solutions for perturbed relativistic pendulum systems. *Proc. Am. Math. Soc.* **2015**, *143*, 3029–3039.
5. Jebelean, P.; Mawhin, J.; Şerban, C. Morse theory and multiple periodic solutions of some quasilinear difference systems with periodic nonlinearities. *Georgian Math. J.* **2017**, *24*, 103–112.
6. Jebelean, P.; Şerban, C. Fisher-Kolmogorov type perturbations of the relativistic operator: differential vs. difference. *Proc. Am. Math. Soc.* **2018**, *144*, 2005–2014.
7. Jebelean, P.; Şerban, C. Relativistic type systems with parametric odd nonlinearities. **2022**, *submitted*.
8. Zhao, Z.; Ma, R.; Zhu, Y. Existence and multiplicity of sign-changing solutions for the discrete periodic problems with Minkowski-curvature operator. *J. Appl. Anal. Comput.* **2022**, *12*, 347–360.
9. Szulkin, A. Minimax principles for lower semicontinuous functions and applications to nonlinear boundary value problems. *Ann. l'Inst. Henri Poincaré C Anal. Non Linéaire* **1986**, *3*, 77–109.
10. Rabinowitz, P.H. *Some Aspects of Critical Point Theory*; MRC Tech. Rep. #2465; University of Wisconsin–Madison Mathematics Research Center: Madison, WI, USA, 1983.
11. Rabinowitz, P.H. Variational methods for nonlinear eigenvalue problems. In *Eigenvalues of Non-Linear Problems*; Prodi, G., Ed.; C.I.M.E. Summer Schools Series; Springer: Berlin/Heidelberg, Germany, 1974; Volume 67, pp. 139–195.
12. Elliott, J.F. The Characteristic Roots of Certain Real Symmetric Matrices. Master's Thesis, University of Tennessee, Knoxville, TN, USA, 1953.
13. Bian, L.-H.; Sun, H.-R.; Zhang, Q.-G. Solutions for discrete  $p$ -Laplacian periodic boundary value problems via critical point theory. *J. Differ. Equ. Appl.* **2012**, *18*, 345–355.