

# Article Pricing of Arithmetic Average Asian Option by Combining Variance Reduction and Quasi-Monte Carlo Method

Lingling Xu <sup>1</sup>, Hongjie Zhang <sup>2</sup> and Fu Lee Wang <sup>1,\*</sup>

Abstract: Financial derivatives have developed rapidly over the past few decades due to their riskaverse nature, with options being the preferred financial derivatives due to their flexible contractual mechanisms, particularly Asian options. The Black-Scholes stock option pricing model is often used in conjunction with Monte Carlo simulations for option pricing. However, the Black-Scholes model assumes that the volatility of asset returns is constant, which does not square with practical financial markets. Additionally, Monte Carlo simulation suffers from slow error convergence. To address these issues, we first correct the asset volatility in the Black-Scholes model using a GARCH model. Then, the low error convergence rate of the Monte Carlo method is improved using variance reduction techniques. Meanwhile, the quasi-Monte Carlo approach based on low discrepancy sequences is used to refine the error convergence rate. We also provide a simulation experiment and result analysis to validate the effectiveness of our proposed method.

Keywords: Asian option pricing; GARCH model; Monte Carlo method; variance reduction technique; low-discrepancy sequence

MSC: 91-10



Citation: Xu, L.; Zhang, H.; Wang, F.L. Pricing of Arithmetic Average Asian Option by Combining Variance Reduction and Quasi-Monte Carlo Method. Mathematics 2023, 11, 594. https://doi.org/10.3390/ math11030594

Academic Editors: Steven P. Clark, Alberto Bueno-Guerrero and Java Bishwal

Received: 7 December 2022 Revised: 19 January 2023 Accepted: 20 January 2023 Published: 23 January 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/).

# 1. Introduction

The complexity of financial markets has led to an increase in trading risk and has given rise to a number of financial derivative products to hedge risk. One such product is the Asian option [1,2], which is a path-dependent option whose return is determined by the average price of the underlying asset over a certain period of time. This feature also makes Asian options popular with investors as they can help hedge risk.

The Black–Scholes (B-S) option pricing model [3] further enhances the development of the options market. However, this model assumes that the volatility of asset returns is a constant, which differs significantly from the real financial market. To this end, the Constant Elasticity of Variance (CEV) model [4] and Heston stochastic volatility model [5] are designed. Although these two volatility models fit well with the actual financial data, a CEV model is not suitable for stock option [6], and the stochastic volatility model is too costly to implement and practically infeasible for empirical work [7]. Bollerslov [8] showed that the GARCH model performs well in fitting stock price return volatility, and studies from [9–14] further validated this and demonstrated that GRACH model helps to improve option pricing in the B-S model.

Initially, most approaches used Brownian motion to price Asian options with the help of probability theory. Subsequently, a number of studies have been conducted to investigate Asian option pricing, including Monte Carlo (MC) methods [15–18], Fourier transform [19], Laplace transform [20], and partial differential equation (PDE) approaches [21]. Notably, these Asian option pricing methods are all developed based on probability

<sup>1</sup> School of Science and Technology, Hong Kong Metropolitan University, Hong Kong 999077, China

Zhongtai Securities Institute for Financial Studies, Shandong University, Jinan 250100, China Correspondence: pwang@hkmu.edu.hk

theory. Moreover, several Asian option pricing methods [2,22] are devised for an uncertain financial market. The path-dependent nature of Asian options makes the MC approach a good choice for pricing Asian options. The MC method was introduced to option pricing by Boyle [15] and further studied by Joy et al. [16] to deal with Asian option pricing problems. However, the error convergence of the MC method is slow. To overcome these issues, Kemna and Vorst [23] applied the control variate variance reduction technique [24] to the pricing of the arithmetic average Asian option, improving the simulation error accuracy. On the other hand, [25–27] pointed out that the Quasi-Monte Carlo (QMC) method based on low-discrepancy sequences can enhance the error convergence rate of MC simulation.

In this paper, we first modify the B-S model using the GARCH volatility model and then consider combining variance reduction techniques and the QMC method to improve MC simulation. In particular, the QMC method based on low-discrepancy sequences can reduce the error convergence rate from  $O(1/\sqrt{N})$  to O(1/N). Specifically, the sum of the underlying assets price during the option period is regarded as a control variable for variance reduction, incorporating the QMC method based on the Sobol sequence [28]. Numerical simulation results indicate that the combination of these two techniques does improve the error accuracy of the simulated estimates of the Asian options. The main contribution of this paper can be summarized as follows:

- Combining the GARCH-based B-S model and the variance reduction technique, we develop a new numerical simulation method based on QMC to deal with the price of arithmetic average Asian options.
- The variance reduction technique and quasi-Monte Carlo method are combined to price the arithmetic average Asian options, significantly improving the error convergence rate of MC simulation.

The rest of the paper is organized as follows. Section 2 introduces some preliminary concepts of B-S models with the GARCH volatility model, Asian option pricing, and MC methods. Section 3 discusses strategies to improve the error convergence accuracy of MC simulation, i.e., variance reduction techniques and QMC methods. Then, we conduct numerical simulation experiments to demonstrate the effectiveness of our proposed method in Section 4. Finally, the conclusion and future work are made in Section 5.

#### 2. Preliminaries

# 2.1. Black–Scholes Model with GARCH Volatility

The Black–Scholes (B-S) model has received great popularity ever since it was proposed in 1973 [3]. Merton expanded on it by considering the impact of dividend payouts and proposed the Black–Scholes–Merton (B-S-M) model [29]. Assume  $S_t$  is the price of the underlying asset at time t. The B-S model assumes that, under the risk neutral probability measure, the change of underlying asset (stock)  $S_t$  follows a geometric Brownian motion, with the expression

$$dS_t = rS_t dt + \sigma S_t dW_t, \tag{1}$$

in which *r* (constant) is the risk-free interest rate,  $\sigma$  (constant) is the volatility of the underlying asset return, and  $W_t \sim N(0,t)$  follows standard Brownian motion. Then, the stock price change at any time step [t, T] follows a normal distribution with mean value of  $(r - \frac{\sigma^2}{2})(T - t)$  and variance of  $\sigma^2(T - t)$ , that is

$$S(T) = S(t) \exp\left[\left(r - \frac{\sigma^2}{2}\right)(T - t) + \sigma \epsilon \sqrt{T - t}\right], \ \epsilon \sim N(0, 1).$$
<sup>(2)</sup>

Next, we suppose *K* is the option expiry strike price and V(S, t) is the option price that relies on stock price *S* and time *t*; then, we can obtain Equation (3) with the condition  $V(S,T) = \mathbb{E}[\max(S(T) - K, 0)].$ 

$$\frac{\partial V(S,t)}{\partial t} + rS\frac{\partial V(S,t)}{\partial S} + \frac{1}{2}\sigma^2 S^2 \frac{\partial^2 V(S,t)}{\partial S^2} = rV(S,t)$$
(3)

Taking the European option as example, and supposing the price of the European call option is *c* and that of the European put option is *p*, we can obtain the expression of call option price as

$$c = e^{-r(T-t)} \mathbb{E}[\max(S(T) - K, 0)],$$
 (4)

while the put option price can be derived from Equation (5).

$$c + Ke^{-r(T-t)} = p + S \tag{5}$$

However, an ideal assumption of both the B-S model and B-S-M model is the volatility of asset return; i.e.,  $\sigma$  is a constant, which is not consistent with the real financial market. As a result, much work [12,13,30–32] has focused on investigating the return volatility of the B-S model, with [12,13] showing that fitting return volatility with a GARCH model further improves the B-S model.

The GARCH model is a classic model for depicting changes in the volatility of financial time series, in which models such as GJR [33] and EGARCH [34] have been proposed for studying the volatility of asset return. The GARCH model is an extension of the ARCH model [35], integrating moving average (MA) and auto-regressive (AR) models. When using the GARCH(1, 1) model to fit the asset return volatility of underlying assets in the options, it can be expressed in Equation (6),

$$\begin{cases} r_n = \mu + u_n, \\ u_n = \sigma_n \epsilon_n, \quad \epsilon_n \text{ i.i.d.} \sim N(0, 1), \\ \sigma_n^2 = \lambda + \alpha_1 u_{n-1}^2 + \beta_1 \sigma_{n-1}^2. \end{cases}$$
(6)

in which  $Var(u_n) = \frac{\lambda}{1-\alpha_1-\beta_1}$ ,  $\mu$ ,  $\lambda$ ,  $\alpha_1$ ,  $\beta_1$  are constants and  $\alpha_1 + \beta_1 < 1$ ,  $\lambda + \alpha_1 + \beta_1 < 1$ .  $\alpha_1$  denotes the correlation coefficient between the volatility at the current moment and the squared residual term at the previous moment, and  $\beta_1$  denotes the correlation coefficient between the volatility at the current moment and the volatility at the previous moment. Then, the first *l* steps of the GARCH(1, 1) model forecast can be expressed in Equation (7), so if we know the starting return volatility  $\sigma_h$ , we can obtain the return volatility at any time.

$$\sigma_h^2(l) = \lambda + (\alpha_1 + \beta_1)\sigma_h^2(l-1), l > 1.$$
(7)

#### 2.2. Asian Option Pricing

Asian options, also known as average price options, are derivative products of stock options. Unlike the regular European options, whose payoff depends on the price of the underlying asset at maturity, Asian options provide a payoff that relies on a certain average of the past prices of their underlying asset [36]. According to different settlement methods, Asian options can be classified into fixed strike price and floating strike price Asian options. Fixed strike price options replace the underlying asset price at maturity of a European option S(T) with the average price of the underlying asset during [0, T] A(T), removing the volatility risk associated with frequent asset trading. Meanwhile, floating strike options replace the option's maturity strike price K with A(T), ensuring that the purchaser's average purchase price is less than the final price.

The unique average characteristics of Asian options make them not only cheaper than European options but also more actively traded and favored by investors in the financial derivatives market. Asian options, on the other hand, can be divided into two categories based on the form of the average: geometric and arithmetic averages. In addition, Asian options can be separated into discrete arithmetic average, continuous arithmetic average, discrete geometric average, and continuous geometric average depending on how the average is calculated. In this paper, we focus on the pricing of discrete arithmetic average Asian options. Assume that the price of the underlying asset (stock) at moment  $t_n$  is  $S_{t_n}$  during the option period [0, T]  $(0 = t_0 < t_1 < \cdots < t_{M-1} = T)$ . The average value of underlying assets at maturity  $A(T) = \frac{1}{M} \sum_{n=0}^{M-1} S_{t_n}$ , and the return of arithmetic average Asian option at maturity, can be expressed in Table 1. *K* is the option expiry strike price, and S(T) is the asset (stock) price at maturity.

Table 1. Maturity returns of arithmetic average Asian options.

Туре	Call Option	Put Option
Fixed strike price	$\mathbb{E}[\max(A(T) - K, 0)]$	$\mathbb{E}[\max(K - A(T), 0)]$
Floating strike price	$\mathbb{E}[\max(S(T) - A(T), 0)]$	$\mathbb{E}[\max(A(T) - S(T), 0]]$

Following the B-S model with GARCH(1, 1) volatility in Section 2.1, the average value of underlying assets at maturity A(T) can be rewritten as following

$$\begin{cases} A(T) = \frac{1}{M} \sum_{n=0}^{M-1} S_{t_n}, \\ S_{t_n} = S_{t_{n-1}} \exp\left[(r - \frac{\sigma_{n-1}^2}{2})\frac{M}{T} + \sigma_{n-1}\epsilon \sqrt{\frac{T}{M}}\right]. \end{cases}$$
(8)

Thus, the price of the arithmetic average Asian option can be shown in Table 2.

**Table 2.** Price of arithmetic average Asian options.

Туре	Call Option	Put Option
Fixed strike price Floating strike price	$e^{-rT} \mathbb{E}[\max(A(T) - K, 0)]$ $e^{-rT} \mathbb{E}[\max(S(T) - A(T), 0)]$	$e^{-rT}\mathbb{E}[\max(K - A(T), 0)]$ $e^{-rT}\mathbb{E}[\max(A(T) - S(T), 0]$

#### 2.3. Monte Carlo Method

The Monte Carlo (MC) approach is often used to solve for the expected value of a random variable. The method relies on a large number of computer simulations to approximate the probability or mathematical characteristics of a random variable.

We use the B-S model with GARCH volatility to calculate the price of the underlying asset in the option. Taking the arithmetic average Asian options as the research object, we can obtain a simulated estimate of the price of Asian options using MC simulations. The specific algorithm details can be seen in Algorithm 1.

In particular, for a random variable h(x), its expectation calculated by MC simulation can be approximated as  $\mathbb{E}[h(x)] = \bar{h}(x) = \frac{1}{N} \sum_{n=1}^{N} h(x_n)$ . According to the law of large numbers [37] and central limit theorem [38], when *N* is sufficiently large, at the significance level of  $1 - \alpha$ , we have

$$\lim_{N \to +\infty} P(\frac{|\bar{h}(x) - \nu|}{\omega/\sqrt{N}} \le u_{\alpha/2}) = 1 - \alpha.$$
(9)

Then, we can know that the standard error of the MC simulation estimate is  $O(\frac{\omega}{\sqrt{N}})$ . Thus, we can reduce the error by reducing the variance *w* of variable *X*. On the other hand, we observe that the error convergence order of the MC simulation is  $O(1/\sqrt{N})$  and the convergence rate is not efficient. To further improve the MC approach, we will investigate these two aspects in Section 3.

# Algorithm 1 MC algorithm to estimate expected present value of payoff

1:  $\lambda$ ,  $\alpha_1$ ,  $\beta_1$ ,  $\sigma_0$ , r and T are known constants.  $Z_j$  represents the random data that follow the standard normal distribution.

2: for  $i \leftarrow 1, 2, \cdots, N$  do generate  $Z_i$ 3: for j = 1; j < M - 1; j + + do 4: if j = 1 then 5:  $\sigma_1^2 = \lambda + \alpha_1 u_0^2 + \beta_1 \sigma_0^2$ 6: else  $\sigma_j^2 = \lambda + (\alpha_1 + \beta_1)\sigma_{j-1}^2$ 7: 8: 9:  $S_i(j) = S_i(j-1) \exp \left[ (r - \frac{\sigma_{j-1}^2}{2}) \frac{T}{M} + \sigma_{j-1} \sqrt{\frac{T}{M}} Z_i \right];$ 10: end for 11:  $A_i(T) = \sum_{i=0}^{M-1} S_i(j) / M$ 12:  $C_i = e^{-rT} (A_i(T) - K)^+;$ 13: 14: end for 15:  $\ddot{C}_N = (C_1 + \dots + C_N)/N$ 

# 3. Quasi-Monte Carlo Pricing of Arithmetic Average Asian Options With Control Variate Technique

In this section, we focus on two approaches for improving the error convergence rate of the MC method, the variance reduction technique and the QMC method. Then, we combine the two to perform MC simulation pricing of arithmetic average Asian option.

#### 3.1. Control Variate Variance Reduction Technique

The variance reduction techniques are strategies designed to improve the error convergence rate of MC calculations without modifying their expectation values, i.e., they aim to reduce relative statistical uncertainty. Common variance reduction techniques include antithetic variable variance reduction, control variate variance reduction and importance sampling variance reduction [39,40].

According to the research from [18], using the sum of the prices of the underlying asset over the option period [0, T] and the analytical solution of the geometric average Asian option as control variates can usually achieve good results for arithmetic average Asian options. However, it is very difficult to obtain the analytical solution of the geometric mean Asian option when the volatility of the underlying asset's return follows a GARCH(1, 1) model. Therefore, we choose the sum of the prices of the underlying asset over the option period [0, T] as the control variate to reduce the variance.

The basic idea of the control variate technique is to reduce the variance of an unknown estimate by using information about the known variable, and the higher the correlation between the known variable and the unknown estimate, the better the variance reduction. For random variables  $X_i$  ( $i = 1, 2, \dots, N$ ) and random variables  $Y_i$  ( $i = 1, 2, \dots, N$ ), E(Y) = v is known. For a fixed constant  $\alpha$ , let  $X_i(\alpha) = X_i - \alpha(Y_i - v)$ ; then, we can obtain  $E(X_i(\alpha)) = E(X_i - \alpha(Y_i - v)) = E(X_i) = E(X)$ , and the control variable estimator of the expected value E(X) can be expressed as:

$$CV = \frac{1}{N} \sum_{i=1}^{N} (X - i - \alpha (Y_i - v)).$$
(10)

Calculate the variance of the random variable  $X_i(\alpha) = X_i - \alpha(Y_i - \nu)$ ; then, we have  $Var(X_i(\alpha)) = Var(X_i) + \alpha^2 Var(Y_i) - 2\alpha Cov(X_i, Y_i)$ , and when  $\alpha = \frac{Cov(X_i, Y_i)}{Var(Y_i)}$ , the minimum value of  $Var(X_i(\alpha))$  is equal to  $Var(X_i)(1 - \rho_{XY}^2)$ , where  $\rho_{XY}$  is the correlation coefficient of variables X and Y. The higher the value of  $\rho_{XY}$ , the smaller the variance of  $Var(X_i(\alpha))$ .

Under risk-neutral conditions [41], we assume that the MC simulation value of the *i*-th arithmetic average Asian call option is

$$V(i) = e^{-rT} \left(\frac{1}{M} \sum_{j=1}^{M} S_i(j) - K\right)^+,$$
(11)

then the control variable, the sum of the prices of the underlying asset over the option period [0, T], can be written as  $S(i) = S_i(0) + S_i(2) + \cdots + S_i(M-1)$ . The estimated option price for the arithmetic average Asian option can be expressed as

$$\hat{V}_{CV} = \frac{1}{N} \sum_{i=1}^{N} (V(i) - \alpha(S(i) - S^{\star})).$$
(12)

Here,  $\alpha$  is the correlation coefficient between V(i) and S(i), and  $S^*$  is the expectation of underlying asset price.

#### 3.2. Quasi-Monte Carlo Method

The Quasi-Monte Carlo (QMC) method, also known as the low-discrepancy sequence method, is a method similar to the MC method. The main difference between these two methods lies in the generation of random sequences. MC simulations use pseudo-random sequences, which are aggregated and converge slowly. The QMC method, on the other hand, employs a more evenly distributed random sequence with less discrepancy, which is less aggregated and converges more quickly.

#### 3.2.1. Error Convergence

The QMC method is similar to the MC method, whose numerical integration is approximated by the value of a measurable function f at some points. For example, to find the approximation of the QMC integral over the unit volume, take *N* points  $x_1, x_2, \dots, x_N$ ; then, we have

$$\int_{[0,1]^s} f(u) du \approx \frac{1}{N} \sum_{i=1}^N f(x_i).$$
(13)

 $x_i$  ( $i = 1, 2, \dots, N$ ) is *s*-dimensional vector.

The approximated error estimation of the QMC method can be the upper bound of the difference degree of points  $x_1, x_2, \dots, x_N$ . Specifically, the Koksma–Hlawka inequality [42] is one of its classical results, and the error upper bound of QMC integration can be expressed as

$$\left|\frac{1}{N}\sum_{i=1}^{N}f(x_{i})-\int_{I^{s}}f(x)dx\right| \leq V(f)D_{N}^{*}(x_{1},\cdots,x_{N}).$$
(14)

 $x_1, \dots, x_N \in I^s$ , V(f) are bounded variations of the function f on the interval  $I^s$ ; then, the error estimate of the QMC method depends on the difference degree  $D_N^*$ . The convergence order of the deviation of the point set of the quasi-random sequence with N points is  $O(N^{-1}logN^{s-1})$ , so the error convergence rate of the QMC integral is not only related to the number of simulations N but also related to the random sequence dimension s. More importantly, the convergence rate of  $O(N^{-1})$  further improves the MC error convergence rate of  $O(N^{-1/2})$ .

#### 3.2.2. Low-Discrepancy Sequence

The Halton, Faure, and Sobol sequences are three commonly used low-discrepancy sequences. The generation of the Halton sequence depends on a prime number related to the dimension s. If the dimension s = 1, the smallest prime number 2 is taken as the number base. If the dimension s = 2, the second smallest prime number 3 is taken as the number base. Analogously, an s-dimensional Halton sequence would select the top s smallest prime numbers as the number base and then calculate the decimal decimals

with the top *s* prime number as the base for any integer *n*. Meanwhile, the Faure sequence selects the smallest prime number greater than or equal to dimension *s* as the number base for each dimension. As a result, each dimensional sequence in the Faure sequence takes 2 as the number base and is a reordering of the first-dimensional sequence. Different from the Halton and Faure sequence, each dimension of the Sobol sequence consists of a radical inversion with base 2, but each dimension has a different matrix for the radical inversion. Due to each dimension taking 2 as the base, the Sobol sequence can be generated directly using bitwise operations to achieve radical inversion, which is very efficient.

To better demonstrate the difference between pseudo-random and low-discrepancy sequences, we randomly sampled 1000 points on a two-dimensional (first and second dimensions) space to observe their distributions. As shown in Figure 1, the Halton, Faure, and Sobol sequences are more evenly distributed than the pseudo-random sequence. In particular, we observe that the distribution of Faure and Sobol sequences in two-dimensional is the same, since they both take the prime 2 as the base, and the second dimension is a reordering of the data in the first dimension. It is worth noting that as the dimensionality increases, the low-discrepancy sequences degenerate to varying degrees. When the dimension is set to 30, the random data in the two adjacent dimensions (dimensions 29 and 30) of the Halton sequence show a high correlation, and the data are not evenly distributed in U[0, 1]. When the dimension is set to 60, the random data in the two adjacent dimensions (dimension 59 and 60) of the Faure sequence suffer from a similar problem. However, the Sobol sequence does not have this problem, and when the number of dimensions reaches 250, the data in the two adjacent dimensions (dimension 249 and 250) are still well distributed. Taking 1000 data points as examples, results in Halton, Faure and Sobol sequences can be seen in Figure 2.



Figure 1. Two-dimensional distribution of pseudo-random and three low discrepancy sequences.

Furthermore, given that the fact that the Asian options are usually valid for 1 to 9 months, and the change in the price of the underlying asset is usually calculated in days, the pricing of Asian options is a high-dimensional problem (the dimension is usually greater than 30). Given that the Sobol sequence performs better in low-dimension and high-dimension scenarios, we choose the Sobol low-discrepancy sequence to generate random



points for our numerical experiments. The construction details of the Sobol sequence can be seen in Appendix A.

**Figure 2.** Two-dimensional distribution of adjacent low-discrepancy sequences (high-dimensional case). The Halton and Faure sequences show varying degrees of degeneracy, while the Sobol sequence still shows good uniformity.

#### 3.2.3. Inverse Transformation Algorithm

Notably, the above three low discrepancy sequences all obey the U[0, 1] distribution, but the B-S formula requires a normally distributed sequence, so it is necessary to convert the uniformly distributed sequences into standard normally distributed sequences. Although Box–Muller [43–45] is the classical algorithm for solving this problem, it disturbs the uniformity of the low-discrepancy sequences. We will adopt the Moro algorithm [46] for transformation. Since the value of the *Y*-axis in the standard normal distribution obeys the U[0, 1] distribution, an intuitive idea is to inverse-normally transform the value of the *Y*-axis. The Moro algorithm divides the *Y*-axis of the standard normal distribution into two parts: the peak part ( $0.08 \le y \le 0.92$ ) and the fat tail part (y < 0.08 and y > 0.92).

The Moro algorithm uses different algorithms for different parts. Let y be a random number in the low-discrepancy sequence and u = y - 0.5.

• If  $|u| \le 0.42$ , then we use the Beasley algorithm [47] to calculate the estimate of the inverse function

$$F^{-1}(y) = u \frac{\sum_{n=0}^{3} a_n u^{2n}}{\sum_{n=0}^{4} b_n u^{2n}}.$$
(15)

• If |u| > 0.42, then we employ the truncated Chebyshev sequence to calculate the estimate of the inverse function

$$F^{-1}(y) = \begin{cases} \sum_{n=0}^{8} c_n T_n(z) - c_0/2, & 0.42 < u < 0.5\\ c_0/2 - \sum_{n=0}^{8} c_n T_n(z), & -0.5 < u < -0.42 \end{cases}$$
(16)

in which  $z = k_1 \{2 \ln(-\ln(0.5 - |u|)) - k_2\}$ , and the parameter value of  $a_n, b_n, c_n$  can be seen in Appendix B.

Next, we employ the Moro algorithm to transform the Sobol sequence into a normal distribution sequence. We consider the standard normal distribution for the two adjacent dimensions of 29 and 30, as well as the distribution of random data from the Sobol sequence in two adjacent dimensions after Moro normalization when the dimensions are set to 30, 90, and 180, respectively. As illustrated in Figure 3, the normal distribution of the Moro-normalized Sobol sequence of adjacent two-dimensional data holds well as the number of dimensions increases.



Figure 3. Two-dimensional distribution of adjacent Moro-normarlized Sobol sequence with dimensions set to 30, 90, and 180.

# 4. Numerical Results and Analysis

Although the option information is publicly available, we choose to set appropriate data manually for numerical simulation pricing of arithmetic average Asian options for convenience. This is due to the objective of this paper to improve the Monte Carlo simulation method. As a result, the numerical simulation we implement is to verify whether our proposed method improves the accuracy of error convergence in Asian option prices. We choose the B-S model with the GARCH(1, 1) volatility to model the change of underlying assets during the option period. Additionally, we select the sum of the underlying asset prices within the option valid period as the control variable for variance reduction and combine it with the QMC method based on the Sobol sequence. All the numerical simulation experiments in this paper are implemented in Python, and the code is available at https://github.com/Linda230/Option\_pricing (accessed on 1 December 2022).

#### 4.1. Main Results

Taking the arithmetic mean Asian call option with a fixed strike price *K* as an example, the steps to simulate and solve the option price are as follows:

- 1. Generate Sobol sequence  $Z_n$ , then utilize the Moro algorithm to normalize  $Z_n$ , i.e., find  $F^{-1}(Z_n)$ .
- 2. Substitute  $F^{-1}(Z_n)$  into the option underlying asset pricing formula

$$S(t + \Delta t) = S(t) \exp[(r - \frac{\sigma_t^2}{2})\Delta t + \sigma \sqrt{\Delta t} F^{-1}(Z_n)],$$

in which  $\sigma_t$  follows GARCH(1, 1) model, and  $\Delta t = \frac{T}{M}$ .

3. Calculate the maturity return of the arithmetic average Asian option and discount it with the risk-free rate *r* to obtain the estimated option price of the *i*-th simulation path.

$$C(i) = e^{-rT} \left(\frac{1}{M} \sum_{j=0}^{M-1} S_i(j) - K\right)^+.$$

4. Use the price sum of the underlying asset of the option over the option validity [0, *T*] as the control variable to reduce the variance; then, the new variable can be expressed as:

$$\hat{C}(i) = C(i) - \alpha(S_i - S^*),$$

in which  $S_i = S_i(0) + S_i(1) + \cdots + S_i(M-1)$ ,  $\alpha = \frac{cov(C(i),S_i)}{var(S_i)}$ , and  $S^*$  is the expectation value of the underlying asset price sum over the period of [0, T].

$$S^* = E(S_i(0) + S_i(1) + \dots + S_i(M-1))$$
  
=  $S_i(0)(1 + e^{r\Delta t} + e^{2r\Delta t} + \dots + e^{(M-1)r\Delta t})$   
=  $S_i(0)\frac{(1 - e^{rT})}{1 - e^{r\Delta t}}.$ 

5. Simulate *N* times to obtain the estimated mean value of the price of the arithmetic average Asian option.

$$\hat{C} = \frac{\hat{C}(1) + \hat{C}(2) + \dots + \hat{C}(N)}{N}$$

Assume the initial price of the underlying asset (stock) in the Asian option  $S_i(0) = 50$ , a fixed strike price of K = 50, and a risk-free daily return of r = 0.0005. The volatility model for the stock price return is

$$\begin{cases} r_n = 0.0002 + u_n, \\ u_n = \sigma_n \epsilon_n, \quad \epsilon_n \text{ i.i.d. } \sim N(0, 1), \\ \sigma_{n+1}^2 = 0.0006 + 0.1u_n^2 + 0.85\sigma_n^2. \end{cases}$$

and the initial daily volatility of the stock price return is  $\sigma_0 = 0.02$ . Here, the GARCH(1, 1) volatility model considered is an overnight volatility change, and the time interval is 1 day. That is, if the option is valid for 30 days, then the number of dimensions in the simulation is 30. We use Python for programming to calculate the simulated estimated price and standard error of arithmetic average Asian option. As shown in Table 3, the price of the fixed-strike price arithmetic average Asian call option increases with the effective period of the option, while the standard error of the option price decreases as the number of simulations increases.

**Table 3.** Price of arithmetic average Asian options (fixed-strike call option). The values in brackets denote the standard error of the MC simulation estimates.

Number of Simulations	T = 30	T = 90	T = 180
N = 1000	2.6498(0.0027)	8.6223(0.0062)	14.2003(0.0092)
N = 10,000	2.8358(0.0003)	8.7017(0.0006)	13.8482(0.0009)
N = 100,000	2.8757(0.00003)	8.7498(0.00006)	13.9332(0.00009)

# 4.2. Ablation Study

Our proposed method incorporates the control variate technique and QMC method. To verify their effectiveness, we conduct an ablation study for each of the two components with a simulation number of N = 10,000 and an option validity of T = 30/90/180. As shown

in Table 4, removing the control variable or the low-discrepancy sequence will result in a degradation of the standard error and a small change in option price. In addition, we investigate the effect of volatility on the option price. We find the removal of GARCH(1,1) volatility leads to a dramatic decrease in the option prices. The reason for this is that the greater the volatility of the underlying asset's return during the term of the option, the greater the price change of the underlying asset price, and the higher the corresponding option price.

Model	T = 30	T = 90	T = 180
Ours	2.8358(0.0003)	8.7017(0.0006)	14.2003(0.0092)
w/o Control Variate	3.8336(0.0007)	9.0521(0.0019)	13.6596(0.0038)
w/o Sobol Sequence	2.8392(0.0278)	8.7498(0.00006)	13.9332(0.00009)
w/o GARCH(1,1)	1.3968(0.00009)	2.6469(0.0002)	4.0149(0.0002)
w/o ALL	1.3871(0.0201)	2.6589(0.0373)	4.0166(0.0546)

Table 4. Ablation study on the control variate and QMC method based on Sobol sequence.

#### 5. Conclusions

This paper takes arithmetic average Asian option pricing as the research object and focuses on improving the error convergence rate of Monte Carlo simulation. To begin with, the GARCH model is used to fit the volatility of asset returns to modify the constant volatility assumption of the B-S model. To address the problem of low accuracy of error estimation in the MC method, we consider combining variance reduction techniques with the QMC method to improve the accuracy. Specifically, we use the sum of the underlying asset prices within the option valid period as the control variable for variance reduction and combine it with the QMC method based on the Sobol sequence. The empirical results show a significant improvement in the standard error convergence rate of arithmetic average Asian options, demonstrating the effectiveness of our proposed method.

**Author Contributions:** Conceptualization, L.X. and H.Z.; methodology, L.X.; software, H.Z.; validation, L.X., H.Z. and F.L.W.; formal analysis, L.X. and F.L.W.; investigation, L.X.; writing—original draft preparation, L.X., H.Z. and F.L.W.; writing—review and editing, H.Z. and F.L.W.; supervision, F.L.W.; project administration, F.L.W. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

**Acknowledgments:** This article is partially translated and adapted from Lingling Xu's unpublished master thesis entitled "Monte Carlo simulation improvement study of arithmetic average Asian option pricing" (In Chinese) submitted to Shangdong University, China, in 2018.

Conflicts of Interest: The authors declare no conflict of interest.

### Abbreviations

The following abbreviations are used in this manuscript:

- B-S Black–Scholes
- MC Monte Carlo
- QMC Quasi-Monte Carlo
- B-S-M Black–Scholes–Merton
- CEV Constant Elasticity of Variance
- PDE Partial Differential Equation

# Appendix A

Sobol sequences are generated based on a set of sequences called direction numbers  $d_i$ , and the number base of the sequence is fixed to 2. Set  $d_1$  as a positive odd number less than  $2^i$ ,

$$d_i = \frac{n_i}{2^i}$$

Construction method for Sobol sequence:

1. Let f(x) be a polynomial with coefficients of 0 or 1.

$$f(z) = x^p + c_1 x^{p-1} + \dots + c_{p-1} x + c_p.$$

When i > p, we have a recursive formula that

$$d_i = c_1 d_{i-1} \oplus c_2 d_{i-2} \oplus \cdots \oplus c_p d_{i-p} \oplus [d_{i-p}/2^p],$$

in which  $\oplus$  represents the iso-or operation symbol in binary (same to be 0, different to be 1). That is,  $1 \oplus 0 = 1$ ;  $0 \oplus 0 = 0$ ;  $1 \oplus 1 = 0$ ).

2. Calculate  $n_i$ ; then, we have

$$n_i = 2c_1n_{i-1} \oplus 2^2c_2n_{i-2} \oplus \cdots \oplus 2^pc_pn_{i-p} \oplus n_{i-p}.$$

The following example illustrates how the Sobol sequence is generated by taking the simple polynomial to be  $x^3 + x^2 + 1$ ; then, we take the corresponding recursive polynomial to be

$$n_i = 2n_{i-1} \oplus 8n_{i-3} \oplus n_{i-3}.$$

If we take the initial values of  $n_1 = 1$ ,  $n_2 = 1$  and  $n_3 = 3$ , then we have

$$n_4 = 2n_3 \oplus 8n_1 \oplus n_1$$
  
= 6 \overline 8 \overline 1  
= (0110)\_2 \overline (1000)\_2 (0001)\_2  
= (1111)\_2  
= 15.

Then, we can obtain the *i*-th number in the Sobol sequence

$$s_i = b_1 d_1 \oplus b_2 d_2 \oplus \cdots, \tag{A1}$$

where( $\cdots b_2 b_1$ ) represents the binary expression of number *i*, i.e.,  $i = (\cdots b_2 b_1)_2$ .

# Appendix **B**

The specific parameter settings of the Moro algorithm can be seen in Table A1.

Table A1.	The specific	parameter	settings of	of More	o algorithi	m.
-----------	--------------	-----------	-------------	---------	-------------	----

a <sub>n</sub>	$b_n$	n	<i>c</i> <sub>n</sub>
2.506 628 238 84	1.00	0	7.710 887 070 548 789 5
$-18.615\ 000\ 625\ 29$	$-8.473\ 510\ 930\ 90$	1	2.777 201 353 368 516 9
41.391 197 735 34	23.083 367 437 43	2	0.361 496 412 926 100 2
$-25.441\ 060\ 496\ 37$	$-21.062\ 241\ 018\ 26$	3	0.037 341 823 343 455 4
	3.130 829 098 33	4	0.002 829 714 303 696 7
		5	0.000 162 571 691 792 2
		6	0.000 008 017 330 474 0
$k_1$	$k_2$	7	0.000 000 384 091 986 5
$0.417\ 988\ 642\ 492\ 643\ 1$	4.245 468 688 137 656 9	8	0.000 000 012 970 717 0

# References

- 1. Rogers, L.C.G.; Shi, Z. The value of an Asian option. J. Appl. Probab. 1995, 32, 1077–1088.
- 2. Sun, J.; Chen, X. Asian option pricing formula for uncertain financial market. J. Uncertain. Anal. Appl. 2015, 3, 11.
- 3. Black, F.; Scholes, M. The pricing of options and corporate liabilities. J. Political Econ. 1973, 81, 637–654.
- 4. Cox, J.; Ross, A. The valuation of option for alternative stochastic processes. J. Financ. Econ. 1976, 3, 145–166.
- 5. Heston, S. A closed—Form solution for options with stochastic volatility with applications to bond and currency options. *Rev. Financ. Stud.* **1993**, *6*, 327–343.
- 6. Bates, D.S. 20 testing option pricing models. *Handb. Stat.* 1996, 14, 567–611.
- 7. Chen, R.R.; Lee, C.F.; Lee, H.H. Empirical performance of the constant elasticity variance option pricing model. *Rev. Pac. Basin Financ. Mark. Policies* **2009**, *12*, 177–217.
- 8. Bollerslev, T. Generalized autoregressive conditional heteroskedasticity. J. Econom. 1986, 31, 307–327.
- 9. Duan, J.C. The GARCH option pricing model. Math. Financ. 1995, 5, 13–32.
- 10. Kallsen, J.; Taqqu, M. Option pricing in ARCH-type models. Math. Financ. 1998, 8, 13-26.
- 11. Heston, S.; Nandi, S. A closed-form GARCH option pricing model. Rev. Financ. Stud. 2000, 13, 586-625.
- 12. Harikumar, T.; De Boyrie, M.E.; Pak, S.J. Evaluation of Black-Scholes and GARCH models using currency call options data. *Rev. Quant. Financ. Account.* 2004, 23, 299–312.
- 13. Gong, H.; Thavaneswaran, A.; Singh, J. A black-scholes model with garch volatility. Math. Sci. 2010, 35, 37–42.
- 14. Sheraz, M.; Preda, V. Implied volatility in black-scholes model with GARCH volatility. Procedia Econ. Financ. 2014, 8, 658-663.
- 15. Boyle, P.P. Options: A monte carlo approach. J. Financ. Econ. 1977, 4, 323–338.
- 16. Joy, C.; Boyle, P.P.; Tan, K.S. Quasi-Monte Carlo methods in numerical finance. Manag. Sci. 1996, 42, 926–938.
- 17. Lapeyre, B.; Temam, E. Competitive Monte Carlo methods for the pricing of Asian options. J. Comput. Financ. 2001, 5, 39-58.
- 18. Mehrdoust, F. A new hybrid Monte Carlo simulation for Asian options pricing. J. Stat. Comput. Simul. 2015, 85, 507-516.
- 19. Fusai, G.; Meucci, A. Pricing discretely monitored Asian options under Lévy processes. J. Bank. Financ. 2008, 32, 2076–2088.
- 20. Cai, N.; Song, Y.; Kou, S. A general framework for pricing Asian options under Markov processes. Oper. Res. 2015, 63, 540-554.
- 21. Vecer, J. A new PDE approach for pricing arithmetic average Asian options. J. Comput. Financ. 2001, 4, 105–113.
- 22. Lu, Z.; Zhu, Y.; Li, B. Critical value-based Asian option pricing model for uncertain financial markets. Physica A 2019, 525, 694–703.
- 23. Kemna, A.; Vorst, A. A pricing method for option based on average asset values. J. Bank. Financ. 1990, 14, 113–129.
- 24. Kleijnen, J.P.; Ridder, A.; Rubinstein, R. Variance Reduction Techniques in Monte Carlo Methods; Springer: Berlin/Heidelberg, Germany, 2010.
- 25. Galanti, S.; Jung, A. Low-discrepancy sequences: Monte Carlo simulation of option prices. J. Deriv. 1997, 5, 63–83.
- Kocis, L.; Whiten, W.J. Computational investigations of low-discrepancy sequences. *ACM Trans. Math. Softw.* 1997, 23, 266–294.
   De Rainville, F.M.; Gagné, C.; Teytaud, O.; Laurendeau, D. Evolutionary optimization of low-discrepancy sequences. *ACM Trans.*
- *Model. Comput. Simul.* 2012, 22, 1–25.
  28. Sobol, I.M. The distribution of points in a cube and the approximate evaluation of integrals. *Zhurnal Vychislitelnoi Mat. Mat. Fiz.* 1967, 7, 784–802.
- 29. Merton, R.C. Theory of rational option pricing. Bell J. Econ. Manag. Sci. 1973, 4, 141-183.
- 30. Bishwal, J.P. Sequential Monte Carlo methods for stochastic volatility models: A review. J. Interdiscip. Math. 2010, 13, 619–635.
- 31. Duan, J.C. Pricing foreign currency and cross-currency options under GARCH. J. Deriv. 1999, 7, 51-63.
- 32. Clark, S.P.; Dickson, M. Performance expectations of basic options strategies may be different than you think. *J. Asset Manag.* **2019**, *20*, 91–102.
- 33. Glosten, L.R.; Jagannathan, R.; Runkle, D.E. On the relation between the expected value and the volatility of the nominal excess return on stocks. *J. Financ.* **1993**, *48*, 1779–1801.
- 34. Nelson, D.B. Conditional heteroskedasticity in asset returns: A new approach. Econom. J. Econom. Soc. 1991, 59, 347–370.
- 35. Engle, R.F. Autoregressive conditional heteroscedasticity with estimates of the variance of United Kingdom inflation. *Econom. J. Econom. Soc.* **1982**, *50*, 987–1007.
- 36. Bueno-Guerrero, A.; Moreno, M.; Navas, J.F. Hedging Asian bond options with Malliavin calculus under stochastic string models. In *New Methods in Fixed Income Modeling*; Springer: Cham, Switzerland, 2018; pp. 169–180
- 37. Dekking, F.M.; Kraaikamp, C.; Lopuhaä, H.P.; Meester, L.E. A Modern Introduction to Probability and Statistics: Understanding Why and How; Springer: London, UK, 2005; Volume 488, pp. 181–190, 333
- 38. Montgomery, D.C.; Runger, G.C. Applied Statistics and Probability for Engineers; John Wiley & Sons: Hoboken, NJ, USA, 2010.
- 39. Law, A.M.; Kelton, W.D.; Kelton, W.D. Simulation Modeling and Analysis; Mcgraw-Hill: New York, NY, USA, 2007; Volume 3.
- 40. Rubinstein, R.Y.; Kroese, D.P. Simulation and the Monte Carlo Method; John Wiley & Sons: Hoboken, NJ, USA, 2016.
- 41. Cox, J.C.; Ross, S.A. A survey of some new results in financial option pricing theory. J. Financ. 1976, 31, 383–402.
- 42. Hickernell, F.J. Koksma-Hlawka Inequality. In *Wiley StatsRef: Statistics Reference Online;* John Wiley & Sons: Hoboken, NJ, USA, 2014.
- 43. Box, G. A Note on the Generation of Random Normal Deviates. *Ann. Math. Stat.* **1958**, *29*, 610–611.
- 44. Scott, D. W. Box–muller transformation. *Wiley Interdiscip. Rev. Comput. Stat.* **2011**, *3*, 177–179.
- Ökten, G.; Göncü, A. Generating low-discrepancy sequences from the normal distribution: Box–Muller or inverse transform? *Math. Comput. Model.* 2011, 53, 1268–1281.

46. Glasserman, P. Monte Carlo Methods in Financial Engineering; Springer: New York, NY, USA, 2004; Volume 53.

47. Beasley, J.D.; Springer, S.G. The percentage points of the normal distribution. *Appl. Stat.* **1977**, *26*, 118–121.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.